

Supporting Information

Some Considerations about the Anodic Limit of Ionic Liquids Obtained by Means of DFT Calculations

Annalisa Paolone ^{1,*}, Simone Di Muzio ^{1,2}, Oriele Palumbo ¹ and Sergio Brutti ^{1,3}

¹ Consiglio Nazionale delle Ricerche, Istituto dei Sistemi Complessi, Piazzale Aldo Moro 5, 00185 Rome, Italy; simone.dimuzio1@graduate.univaq.it (S.D.M.); oriele.palumbo@roma1.infn.it (O.P.); sergio.brutti@uniroma1.it (S.B.)

² Department of Physical and Chemical Sciences, University of L'Aquila, Via Vetoio, 67100 L'Aquila, Italy

³ Department of Chemistry, Sapienza University of Rome, Piazzale Aldo Moro 5, 00185 Rome, Italy

* Correspondence: annalisa.paolone@roma1.infn.it

Citation: Paolone, A.; Di Muzio, S.; Palumbo, O.; Brutti, S. Some Considerations about the Anodic Limit of Ionic Liquids Obtained by Means of DFT Calculations. *Entropy* **2023**, *25*, 793. <https://doi.org/10.3390/e25050793>

Academic Editor:
Antonio M. Scarfone

Received: 4 April 2023
Revised: 5 May 2023
Accepted: 11 May 2023
Published: 12 May 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).




Figure S1. Experimental anodic limit versus the energy (in atomic units), LUMO and HOMO levels (in eV) and the minimum of the ionization potential (in eV) calculated at the Hartree-Fock level using the 6-31G** basis set. The red curves are the best fit lines obtained by a linear regression.

