

Article

Expeditious Entry to Novel 2-Methylene-2,3-dihydrofuro[3,2-*c*]chromen-2-ones from 6-Chloro-4-hydroxychromen-2-one and Propargylic Alcohols

Noel Nebra, Alba E. Díaz-Álvarez, Josefina Díez and Victorio Cadierno *

Departamento de Química Orgánica e Inorgánica, IUQOEM-CSIC, Facultad de Química, Universidad de Oviedo, E-33071 Oviedo, Spain

* Author to whom correspondence should be addressed; E-Mail: vcm@uniovi.es;
Tel.: +34-985-103-453; Fax: +34-985-103-446.

Received: 14 July 2011; in revised form: 27 July 2011 / Accepted: 28 July 2011 /

Published: 2 August 2011

Abstract: A catalytic system consisting of the ruthenium(II) complex $[\text{Ru}(\eta^3\text{-C}_3\text{H}_4\text{Me})(\text{CO})(\text{dppf})][\text{SbF}_6]$ (dppf = 1,1'-bis(diphenylphosphino)ferrocene) and trifluoroacetic acid has been used to promote the coupling of secondary propargylic alcohols with 6-chloro-4-hydroxychromen-2-one. The reactions afforded unusual 2-methylene-2,3-dihydrofuro[3,2-*c*]chromen-2-ones in good yields.


Keywords: chromen-2-ones; furochromen-2-ones; propargylic alcohols; ruthenium catalysts; propargylation; cycloisomerization

1. Introduction

Chromen-2-ones (coumarins) constitute an important family of heterocyclic compounds of natural origin which have attracted considerable attention for many years due to their versatile applications [1–4]. Among them, furochromen-2-ones (furocoumarins), tricyclic systems in which a furan ring is fused to the chromen-2-one unit, are of particular interest since they exhibit potent biological and pharmacological activity [5–8]. Although several methods of synthesis are presently known [7,8], new approaches for the rapid and selective construction of furochromen-2-one scaffolds are still highly desirable. In this context, recent efforts by different groups have been focused in the *one-pot* synthesis of furo[3,2-*c*]chromen-2-ones (Figure 1) from readily available starting materials, with successful examples


including CAN-mediated cycloaddition of 4-hydroxychromen-2-one with terminal alkynes [9,10], rhodium(II)-catalyzed heterocyclization of 3-diazo-2,4-chromenediones with terminal alkynes [11,12], cascade addition/cyclization/oxidation of 3-alkynyl-chromones [13,14], Sonogashira-acetylide coupling/demethylation/cyclization of 3-iodo-4-methoxychromen-2-ones [15,16] and alkynylation/hydroalkoxylation of 3-bromo-4-acetoxychromen-2-ones [17].

Figure 1. The furo[3,2-*c*]chromen-2-one skeleton.



In the course of our studies focused on the application of ruthenium catalysts for the construction of furan- and pyrrole-ring frameworks [18–25], we disclosed a straightforward approach of tetra-substituted furans from readily accessible secondary propargylic alcohols and 1,3-dicarbonyl compounds (Scheme 1) [18]. The process, which proceeds in a *one-pot* manner, involves the initial trifluoroacetic acid-promoted propargylation of the 1,3-dicarbonyl compound, and subsequent cycloisomerization of the resulting γ -ketoalkyne **A** (5-*exo* cyclization + aromatization) catalyzed by the 16-electron allyl-ruthenium(II) complex $[\text{Ru}(\eta^3\text{-2-C}_3\text{H}_4\text{Me})(\text{CO})(\text{dppf})][\text{SbF}_6]$ (**1**).


Scheme 1. Direct synthesis of furans from alkynols and 1,3-dicarbonyl compounds.



By applying this synthetic route a large variety of furans containing carbonyl functionalities on the aromatic ring, could be prepared in good yields starting from both terminal and internal secondary alkynols, and β -diketones or β -keto esters [18,21]. In addition, we also demonstrated that furo[3,2-*c*]chromen-2-ones are also accessible by this route using 4-hydroxychromen-2-one as substrate, representing an appealing *one-pot* method of synthesis for this type of heterocycles [21]. Related work by Zhou and co-workers also confirmed the utility of this propargylation-cycloisomerization sequence for the construction of furochromen-2-one skeletons [26].

Following with these studies, herein we would like to communicate that related C–C coupling processes involving 6-chloro-4-hydroxychromen-2-one and terminal propargylic alcohols $\text{HC}\equiv\text{CC}(\text{OH})\text{HR}$ result in the selective formation of the 2-methylene-2,3-dihydrofuro[3,2-*c*]chromen-2-one derivatives **4** (Figure 2), instead of the expected 8-chloro-substituted furo[3,2-*c*]chromen-2-ones **5**, due to the reluctance of the former to undergo aromatization of the five-membered ring.


Figure 2. Structures of compounds **4** and **5**.



2. Results and Discussion

Initially, the coupling of the secondary propargylic alcohol 1-(4-methoxyphenyl)-2-propyn-1-ol (**2a**) with 6-chloro-4-hydroxychromen-2-one (**3**) was investigated under the same reaction conditions previously employed by us in the synthesis of furo[3,2-*c*]chromen-2-one derivatives starting from 4-hydroxychromen-2-one [21], that is, heating a THF solution of both reactants (equimolar mixture) at 75 °C in the presence of 50 mol% of trifluoroacetic acid and 5 mol% of the allyl-ruthenium(II) complex **1** (Scheme 2). Almost complete disappearance of the starting materials, accompanied by the selective formation of a single reaction product **4a**, was observed by GC after 8 hours of heating.

Scheme 2. Catalytic synthesis of compound **4a** under classical thermal conditions.



Appropriate chromatographic workup allowed the isolation of **4a** as a crystalline yellow solid in 83% yield. NMR spectroscopic data obtained for **4a** clearly revealed the selective formation of a 2-methylene-2,3-dihydrofuran unit, instead of the expected aromatic 2-methylfuran one (details are given in the Experimental), a fact that was unambiguously confirmed by means of a single-crystal X-ray diffraction study (an ORTEP view of the molecule is shown in Figure 3; selected bonding parameters are listed in Table 2). The bond distance C10–C11 (1.321(3) Å) showed the expected value for a C=C bond, while that observed for C10–C12 (1.523(3) Å) falls within the expected range for a C(sp²)-C(sp³) single bond.

Figure 3. ORTEP-type view of the structure of compound **4a** showing the crystallographic labelling scheme. Thermal ellipsoids are drawn at the 20% probability level.





Table 2. Selected bond distances (Å) and angles (°) for compound **4a**.


Distances	
C8-C9	1.338(3)
C9-O3	1.360(2)
O3-C10	1.417(2)
C10-C11	1.321(3)
C10-C12	1.523(3)
C12-C8	1.507(3)
C7-O1	1.393(3)
C7-O2	1.211(3)
C1-C11	1.738(2)
Angles	
C8-C9-O3	114.13(18)
C9-O3-C10	106.39(16)
O3-C10-C11	118.8(2)
C11-C10-C12	131.1(2)
O3-C10-C12	110.07(16)
C10-C12-C8	99.46(17)
C12-C8-C9	109.77(18)

The use of microwave (MW) irradiation represents a convenient alternative to the conventional thermal heating in organic synthesis since a more effective energy transfer to the system takes place, thus shortening considerably the reaction times and improving in many cases the product yields [27–29]. Accordingly, we have found that, performing the same coupling reaction of alkynol **2a** with **3** under controlled microwave heating at 75 °C, selective and almost quantitative formation of 2-methylene-2,3-dihydrofuro[3,2-*c*]chromen-2-one (**4a**, 99% GC-yield; 89% isolated yield) takes place after only 10 min. As shown in Scheme 3, the process is general since the related heterocycles **4b–d** could also be synthesized in good yields (77%–92%) by reacting **3** with the secondary propargylic alcohols 1-(2-methoxyphenyl)-2-propyn-1-ol (**2b**), 1-(1-naphthyl)-2-propyn-1-ol (**2c**) and 1-(2-thienyl)-2-propyn-1-ol (**2d**) under the same MW conditions.

Scheme 3. Catalytic synthesis of compound **4a–d** under MW-irradiation.

The presence of a 2-methylene-2,3-dihydrofuran moiety in the structure of these compounds was readily identified by the appearance of a high-field CH carbon resonance at 43–49 ppm (CHR unit) and a CH_2 signal at *ca.* 92 ppm, typical of a terminal olefinic $=\text{CH}_2$ unit, in their $^{13}\text{C}\{^1\text{H}\}$ -NMR spectra (DEPT experiments). Characteristic ^1H -NMR peaks for these units were also observed at 4.5–5.5 ppm (details can be found in the Experimental).

At this point, it is worthy of note that occurrence of 2-methylene-2,3-dihydrofuro[3,2-*c*]chromen-2-ones has been scarcely documented in the literature [30–35], with most of the known examples being disubstituted at the C-3 position of the 2-methylene-2,3-dihydrofuran ring which prevents their tautomerization into the corresponding 2-methyl-furo[3,2-*c*]chromen-2-ones. In this sense, the reluctance shown by compounds **4a–d** to aromatize under the acidic conditions employed merits to be highlighted. In fact, only in the case of **4b** such aromatization process could be observed after prolonged MW irradiation (3 h) of the reaction mixture at 100 °C. Under these conditions, the novel 8-chloro-substituted furo[3,2-*c*]chromen-2-one **5b** could be synthesized with an acceptable 63% yield and fully characterized (Scheme 4).

Scheme 4. Synthesis of the furo[3,2-*c*]chromen-2-one **5b**.

3. Experimental

3.1. General

Solvents were dried by standard methods and distilled under nitrogen before use. The complex $[\text{Ru}(\eta^3\text{-2-C}_3\text{H}_4\text{Me})(\text{CO})(\text{dppf})][\text{SbF}_6]$ (**1**) [36] and propargylic alcohols **2a–d** [37] were prepared by following the methods reported in the literature. Flash chromatography was performed using Merck silica gel 60 (230–400 mesh). Melting points were determined in a Gallenkamp apparatus and are uncorrected. Infrared spectra were recorded on a Perkin-Elmer 1720-XFT spectrometer. NMR spectra were recorded on a Bruker DPX-300 instrument at 300 MHz (^1H) or 75.4 MHz (^{13}C). The chemical shift values (δ) are given in parts per million and are referred to the residual peak of the deuterated solvent used (CDCl_3). High-resolution mass spectra (HRMS) were provided by the Mass Spectrometry Service of the Instituto de Investigaciones Químicas (IIQ-CSIC, Seville). CCDC 831021 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.

3.2. Synthesis of the 8-chloro-2-methylene-2,3-dihydrofuro[3,2-*c*]chromen-2-ones **4a–d**

Under a nitrogen atmosphere, a pressure-resistant septum-sealed glass vial was charged with the corresponding propargylic alcohol **2a–d** (1 mmol), 6-chloro-4-hydroxychromen-2-one (**3**) (0.197 g, 1 mmol), THF (0.5 mL), $[\text{Ru}(\eta^3\text{-2-C}_3\text{H}_4\text{Me})(\text{CO})(\text{dppf})][\text{SbF}_6]$ (**1**) (0.049 g, 0.05 mmol), $\text{CF}_3\text{CO}_2\text{H}$ (37 μL , 0.5 mmol) and a magnetic stirring bar. The vial was then placed inside the cavity of a CEM Discover[®] S-Class microwave synthesizer and power was held at 300 W until the desired temperature was reached (75 °C). Microwave power was automatically regulated for the remainder of the experiment (10 min) to maintain the temperature (monitored by a built-in infrared sensor). Then, the vial was cooled to room temperature, the volatiles removed under vacuum, and the residue purified by flash chromatography (silica gel) using a mixture EtOAc/hexanes (1:20) as eluent. Characterization data for the novel 8-chloro-2-methylene-2,3-dihydrofuro[3,2-*c*]chromen-2-ones **4a–d** are as follows:

8-Chloro-3-(4-methoxyphenyl)-2-methylene-2,3-dihydrofuro[3,2-*c*]chromen-2-one (4a). Yellow solid (0.303 g, 89%); m.p. 125–127 °C; IR (Nujol) $\nu = 1721$ (C=O) cm^{-1} ; ^1H -NMR (CDCl_3) $\delta = 3.78$ (s, 3H), 4.52 (dd, 1H, $J = 3.4$ and 2.8 Hz), 5.11 (m, 2H), 6.87 (d, 2H, $J = 8.5$ Hz), 7.23 (d, 2H, $J = 8.5$ Hz), 7.32 (d, 1H, $J = 8.8$ Hz), 7.53 (dd, 1H, $J = 8.8$ and 2.2 Hz), 7.75 (d, 1H, $J = 2.2$ Hz) ppm; ^{13}C -NMR (CDCl_3) $\delta = 47.7, 55.2, 91.7, 107.4, 112.5, 114.2, 118.5, 122.2, 128.8, 129.6, 131.2, 132.7, 153.5, 159.1, 157.8, 162.9, 164.8$ ppm; HRMS (EI) $m/z = 340.0501$, $\text{C}_{19}\text{H}_{13}\text{O}_4\text{Cl}$ requires 340.0502.

8-Chloro-3-(2-methoxyphenyl)-2-methylene-2,3-dihydrofuro[3,2-*c*]chromen-2-one (4b). Yellow solid (0.262 g, 77%); m.p. 122–124 °C; IR (Nujol) $\nu = 1748$ (C=O) cm^{-1} ; ^1H -NMR (CDCl_3) $\delta = 3.79$ (s, 3H), 4.55 (dd, 1H, $J = 3.0$ and 2.5 Hz), 5.12 (m, 2H), 6.82–6.91 (m, 3H), 7.25 (m, 1H), 7.34 (d, 1H, $J = 8.9$ Hz), 7.55 (dd, 1H, $J = 8.9$ and 2.4 Hz), 7.76 (d, 1H, $J = 2.2$ Hz) ppm; ^{13}C -NMR (CDCl_3) $\delta = 48.5, 55.3, 92.1, 107.2, 112.6, 113.0, 113.8, 118.6, 120.1, 122.3, 129.8, 130.0, 132.9, 140.7, 153.6, 157.9, 160.0, 163.4, 164.3$ ppm; HRMS (EI) $m/z = 340.0513$, $\text{C}_{19}\text{H}_{13}\text{O}_4\text{Cl}$ requires 340.0502.

8-Chloro-3-(1-naphthyl)-2-methylene-2,3-dihydrofuro[3,2-c]chromen-2-one (4c). Yellow solid (0.331 g, 92%); m.p. 140–142 °C; IR (Nujol) ν = 1719 (C=O) cm^{-1} ; $^1\text{H-NMR}$ (CDCl_3) δ = 4.55 (dd, 1H, J = 3.3 and 2.5 Hz), 5.15 (dd, 1H, J = 3.3 and 2.5 Hz), 5.33 (dd, 1H, J = 3.3 and 3.3 Hz), 7.33–7.49 (m, 5H), 7.81–7.85 (m, 5H) ppm; $^{13}\text{C-NMR}$ (CDCl_3) δ = 48.8, 92.3, 107.3, 112.6, 118.7, 122.4, 125.3, 126.2, 126.4, 127.0, 127.7, 127.9, 129.0, 129.8, 132.9, 133.0, 133.4, 136.5, 153.6, 157.9, 163.4, 164.5 ppm; HRMS (EI) m/z = 360.0557, $\text{C}_{22}\text{H}_{13}\text{O}_3\text{Cl}$ requires 360.0553.

8-Chloro-3-(2-thienyl)-2-methylene-2,3-dihydrofuro[3,2-c]chromen-2-one (4d). Orange solid (0.253 g, 80%); m.p. 131–133 °C; IR (Nujol) ν = 1733 (C=O) cm^{-1} ; $^1\text{H-NMR}$ (CDCl_3) δ = 4.72 (dd, 1H, J = 3.5 and 2.4 Hz), 5.18 (dd, 1H, J = 3.5 and 2.7 Hz), 5.48 (dd, 1H, J = 2.7 and 2.4 Hz), 6.98 (dd, 1H, J = 5.2 and 3.6 Hz), 7.12 (dd, 1H, J = 3.6 and 1.1 Hz), 7.25 (dd, 1H, J = 5.2 and 1.1 Hz), 7.35 (d, 1H, J = 9.0 Hz), 7.56 (dd, 1H, J = 9.0 and 2.5 Hz), 7.75 (d, 1H, J = 2.5 Hz) ppm; $^{13}\text{C-NMR}$ (CDCl_3) δ = 43.0, 92.3, 106.3, 112.1, 118.2, 122.0, 124.9, 125.6, 126.7, 129.4, 132.7, 141.2, 153.1, 157.3, 162.9, 163.0 ppm; HRMS (EI) m/z = 315.9971, $\text{C}_{16}\text{H}_9\text{O}_3\text{ClS}$ requires 315.9961.

3.3. Synthesis of 8-chloro-3-(2-methoxyphenyl)-2-methyl-furo[3,2-c]chromen-2-one (5b)

Under nitrogen atmosphere, a pressure-resistant septum-sealed glass vial was charged with 1-(2-methoxyphenyl)-2-propyn-1-ol (**2b**, 0.162 g, 1 mmol), 6-chloro-4-hydroxychromen-2-one (**3**, 0.197 g, 1 mmol), THF (0.5 mL), $[\text{Ru}(\eta^3\text{-2-C}_3\text{H}_4\text{Me})(\text{CO})(\text{dppf})][\text{SbF}_6]$ (**1**, 0.049 g, 0.05 mmol), $\text{CF}_3\text{CO}_2\text{H}$ (37 μL , 0.5 mmol) and a magnetic stirring bar. The vial was then placed inside the cavity of a CEM Discover® S-Class microwave synthesizer and power was held at 300 W until the desired temperature was reached (100 °C). Microwave power was automatically regulated for the remainder of the experiment (3 h) to maintain the temperature (monitored by a built-in infrared sensor). Then, the vial was cooled to room temperature, the volatiles removed under vacuum, and the residue purified by flash chromatography (silica gel) using a mixture EtOAc/hexanes (1:20) as eluent to give **5b**. Yellow solid (0.214 g, 63%); m.p. 120–122 °C; IR (Nujol) ν = 1749 (C=O) cm^{-1} ; $^1\text{H-NMR}$ (CDCl_3) δ = 2.42 (s, 3H), 3.82 (s, 3H), 7.01–7.08 (m, 2H), 7.26–7.43 (m, 4H), 7.85 (d, 1H, J = 2.5 Hz) ppm; $^{13}\text{C-NMR}$ (CDCl_3) δ = 12.1, 55.1, 110.6, 113.7, 116.2, 118.0, 118.3, 119.6, 120.0, 129.2, 129.4, 131.0, 132.9, 136.5, 150.1, 152.5, 156.9, 162.5, 163.0 ppm; HRMS (EI) m/z = 340.0496, $\text{C}_{19}\text{H}_{13}\text{O}_4\text{Cl}$ requires 340.0502.

3.4. X-ray Crystal Structure Determination of Compound 4a

The most relevant crystal and refinement data are collected in Table 1. Data collection was performed on a Oxford Diffraction Xcalibur Nova single crystal diffractometer, using Cu-K α radiation. Images were collected at a 65 mm fixed crystal-to-detector distance using the oscillation method, with 1° oscillation and a 5 s exposure time per image. Data collection strategy was calculated with the program CrysAlis Pro CCD [38]. Data reduction and cell refinement were performed with the program CrysAlis Pro RED [38]. An empirical absorption correction was applied using the SCALE3 ABSPACK algorithm as implemented in the program CrysAlis Pro RED [38]. The software package WinGX was used for space group determination, structure solution and refinement [39]. The structure was solved by direct methods using SIR92 [40]. Isotropic least-squares refinement on F^2 using

SHELXL97 was performed [41]. During the final stages of the refinements, all the positional parameters and the anisotropic temperature factors of all the non-H atoms were refined. The coordinates of the H atoms were found from different Fourier maps and included in a refinement with isotropic parameters. The function minimized was $[\sum w(F_o^2 - F_c^2)/\sum w(F_o^2)]^{1/2}$ where $w = 1/[\sigma^2(F_o^2) + (aP)^2 + bP]$ ($a = 0.0902$; $b = 0.0000$) with $\sigma^2(F_o^2)$ from counting statistics and $P = (\text{Max}(F_o^2) + 2F_c^2)/3$. Atomic scattering factors were taken from the International Tables for X-ray Crystallography [42]. The crystallographic plot was made with PLATON [43].

Table 1. Crystal data and structure refinement parameters for compound **4a**.

Empirical formula	C ₁₉ H ₁₃ O ₄ Cl
Formula weight	340.74
Temperature	150(2) K
Wavelength	1.5418 Å
Crystal system, space group	<i>triclinic, P-1</i>
Unit cell dimensions	$a = 4.8366(2)$ Å $\alpha = 94.822(4)^\circ$ $b = 11.0016(5)$ Å $\beta = 90.363(4)^\circ$ $c = 14.6466(7)$ Å $\gamma = 94.200(4)^\circ$
Volume	774.45(6) Å ³
Z, calculated density	2, 1.461 mg/m ³
Absorption coefficient	2.369 mm ⁻¹
$F(000)$	352
Crystal size	0.37 × 0.03 × 0.02 mm
Theta range for data collection	3.03 to 73.76°
Limiting indices	−4 ≤ h ≤ 6, −13 ≤ k ≤ 12, −17 ≤ l ≤ 17
Reflections collected / unique	7403/2919 ($R_{int} = 0.0214$)
Completeness to theta = 73.76°	93.4%
Refinement method	Full-matrix least-squares on F^2
Data / restraints / parameters	2919/0/269
Goodness-of-fit on F^2	1.166
Final R indices [$I > 2\sigma(I)$]	$R_1 = 0.0411$, $wR_2 = 0.1177$
R indices (all data)	$R_1 = 0.0517$, $wR_2 = 0.1354$
Largest diff. peak and hole	0.334 and −0.267 e [−] Å ³

4. Conclusions

In summary, an efficient synthesis of unusual and remarkably stable 2-methylene-2,3-dihydrofuro[3,2-*c*]chromen-2-one derivatives, by coupling of secondary propargylic alcohols with commercially available 6-chloro-4-hydroxychromen-2-one, has been developed with the aid of the catalytic system $[\text{Ru}(\eta^3\text{-2-C}_3\text{H}_4\text{Me})(\text{CO})(\text{dppf})][\text{SbF}_6]/\text{CF}_3\text{CO}_2\text{H}$. Apparently, the presence of the electron-withdrawing Cl substituent on the 4-hydroxychromen-2-one skeleton exerts a marked influence on the behavior of these species since, as previously described by us [21], the same reactions performed with its non-substituted counterpart leads to the selective formation of isomeric furo[3,2-*c*]chromen-2-ones by aromatization of the five-membered ring. Overall, the results reported herein

represent a new example of the utility of the allyl-ruthenium(II) complex **1** in synthetic organic chemistry [25].

Acknowledgments

This work was supported by the Spanish MICINN (Projects CTQ2006-08485/BQU, CSD2007-00006 and CTQ2010-14796/BQU). N.N. thanks MEC of Spain and the European Social Fund (FPU program) for the award of a Ph.D. grant.

Conflict of Interest

The authors declare no conflict of interest.

References and Notes

1. Sethna, S.M.; Shah, N.M. The chemistry of coumarins. *Chem. Rev.* **1945**, *36*, 1-62.
2. Murray, R.D.; Méndez, J.; Brown, S.A. *The Natural Coumarins: Occurrence, Chemistry & Biochemistry*; John Wiley & Sons: New York, NY, USA, 1982.
3. Hoult, J.R.S.; Payá, M. Pharmacological and biochemical actions of simple coumarins: Natural products with therapeutic potential. *Gen. Pharmacol.* **1996**, *27*, 713-722.
4. *Coumarins: Biology, Applications and Mode of Action*; O'Kennedy, R., Thornes, R.D., Eds.; John Wiley & Sons: Chichester, UK, 1997.
5. Gambari, R.; Lampronti, I.; Bianchi, N.; Zuccato, C.; Viola, G.; Vedaldi, D.; Dall'Acqua, F. Structure and biological activity of furocoumarins. *Top. Heterocycl. Chem.* **2007**, *9*, 265-276.
6. Conforti, F.; Marrelli, M.; Menichini, F.; Bonesi, M.; Statti, G.; Provenzano, E.; Menichini, F. Natural and synthetic furanocoumarins as treatment for vitiligo and psoriasis. *Curr. Drug Ther.* **2009**, *4*, 38-58.
7. Santana, L.; Uriarte, E.; Roleira, F.; Milhazes, N.; Borges, F. Furocoumarins in medicinal chemistry. Synthesis, applications, and biological activity. *Curr. Med. Chem.* **2004**, *11*, 3239-3261.
8. Traven, V.F. New synthetic routes to furocoumarins and their analogs: A review. *Molecules* **2004**, *9*, 50-66.
9. Lee, Y.R.; Byun, M.W.; Kim, B.S. Efficient one-step synthesis of 2-arylfurans by ceric ammonium nitrate (CAN)-mediated cycloaddition of 1,3-dicarbonyl compounds to alkynes. *Bull. Korean Chem. Soc.* **1998**, *19*, 1080-1083.
10. Kobayashi, K.; Sakashita, K.; Akamatsu, H.; Tanaka, K.; Uchida, M.; Uneda, T.; Kitamura, T.; Morikawa, O.; Konishi, H. CAN-mediated formation of furopyranones and furoquinolinones. *Heterocycles* **1999**, *51*, 2881-2892.
11. Lee, Y.R.; Suk, J.Y.; Kim, B.S. Rhodium(II)-catalyzed reactions of 3-diazo-2,4-chromenediones. First one-step synthesis of pterophyllin 2. *Tetrahedron Lett.* **1999**, *40*, 6603-6607.
12. Tollari, S.; Palmisano, G.; Cenini, S.; Crovotto, G.; Giovenzana, G.B.; Penoni, A. Synthesis of furocoumarins via rhodium(II)-catalysed heterocyclisation of 3-diazobenzopyran-2,4(3H)-dione with terminal alkynes. *Synthesis* **2001**, 735-740.
13. Cheng, G.; Hu, Y. One-pot synthesis of furocoumarins through cascade addition-cyclization-oxidation. *Chem. Commun.* **2007**, 3285-3287.

14. Cheng, G.; Hu, Y. Two efficient cascade reactions to synthesize substituted furocoumarins. *J. Org. Chem.* **2008**, *73*, 4732-4735.
15. Conreux, D.; Belot, S.; Desbordes, P.; Monteiro, N.; Balme, G. Et₃N-Induced demethylation-annulation of 3-alkynyl-4-methoxy-2-pyridones and structurally related compounds in the synthesis of furan-fused heterocycles. *J. Org. Chem.* **2008**, *73*, 8619-8622.
16. Raffa, G.; Rusch, M.; Balme, G.; Monteiro, N. A Pd-catalyzed heteroannulation approach to 2,3-disubstituted furo[3,2-*c*]coumarins. *Org. Lett.* **2009**, *11*, 5254-5257.
17. Chen, L.; Li, Y.; Xu, M.-H. One-pot synthesis of furocoumarins via sequential Pd/Cu-catalyzed alkynylation and intramolecular hydroalkoxylation. *Org. Biomol. Chem.* **2010**, *8*, 3073-3077.
18. Cadierno, V.; Gimeno, J.; Nebra, N. A novel propargylation/cycloisomerization tandem process catalyzed by a ruthenium(II)/trifluoroacetic acid system: One-pot entry to fully substituted furans from readily available secondary propargylic alcohols and 1,3-dicarbonyl compounds. *Adv. Synth. Catal.* **2007**, *349*, 382-394.
19. Albers, J.; Cadierno, V.; Crochet, P.; García-Garrido, S.E.; Gimeno, J. Octahedral ruthenium(II) complexes *cis,cis*-[RuX₂(CNR)(CO)(P[^]P)] and *cis,cis,cis*-[RuX₂(CO)₂(P[^]P)] (X = Cl, Br; P[^]P = 1,1'-bis(diphenylphosphino)ferrocene, 1,1'-bis(diisopropylphosphino)ferrocene): Synthesis and catalytic applications in transfer hydrogenation of acetophenone and cycloisomerization of (*Z*)-3-methylpent-2-en-4-yn-1-ol. *J. Organomet. Chem.* **2007**, *692*, 5234-5244.
20. Cadierno, V.; Gimeno, J.; Nebra, N. One-pot three-component catalytic synthesis of fully substituted pyrroles from readily available propargylic alcohols, 1,3-dicarbonyl compounds and primary amines. *Chem. Eur. J.* **2007**, *13*, 9973-9981.
21. Cadierno, V.; Díez, J.; Gimeno, J.; Nebra, N. Ruthenium/TFA catalyzed coupling of activated secondary propargylic alcohols with cyclic 1,3-diones: Furan vs. pyran ring formation. *J. Org. Chem.* **2008**, *73*, 5852-5858.
22. Cadierno, V.; Crochet, P. Ruthenium-catalyzed furan- and pyrrole-ring formation. *Curr. Org. Synth.* **2008**, *5*, 343-364.
23. Cadierno, V.; Gimeno, J.; Nebra, N. One-pot three-component synthesis of tetrasubstituted N-H pyrroles from secondary propargylic alcohols, 1,3-dicarbonyl compounds and *tert*-butyl carbamate. *J. Heterocycl. Chem.* **2010**, *47*, 233-236.
24. García-Garrido, S.E.; Francos, J.; Cadierno, V.; Basset, J.M.; Polshettiwar, V. Chemistry by nanocatalysis: First example of a solid-supported RAPTA complex for organic reactions in aqueous medium. *ChemSusChem* **2011**, *4*, 104-111.
25. Cadierno, V.; García-Garrido, S.E.; Gimeno, J.; Nebra, N. Atom-economic transformations of propargylic alcohols catalyzed by the 16-electron allyl-ruthenium(II) complex [Ru(η^3 -2-C₃H₄Me)(CO)(dppf)][SbF₆] (dppf = 1,1'-bis(diphenylphosphino)ferrocene). *Inorg. Chim. Acta* **2010**, *363*, 1912-1934.
26. Huang, W.; Wang, J.; Shen, Q.; Zhou, X. Yb(OTf)₃-catalyzed propargylation and allenylation of 1,3-dicarbonyl derivatives with propargylic alcohols: One-pot synthesis of multisubstituted furocoumarin. *Tetrahedron* **2007**, *63*, 11636-11643.
27. *Microwaves in Organic Synthesis*; Loupy, A., Ed.; Wiley-VCH: Weinheim, Germany, 2006.
28. Kappe, C.O.; Dallinger, D.; Murphee, S.S. *Practical Microwave Synthesis for Organic Chemists*; Wiley-VCH: Weinheim, Germany, 2009.

29. *Microwave Heating as a Tool for Sustainable Chemistry*; Leadbeater, N.E., Ed.; CRC Press: Boca Raton, FL, USA, 2011.
30. Sawhney, K.N.; Mathur, K.B.L. Studies on some structural aspects of 4-hydroxycoumarin: Further extension of Meerwein's diazo reaction and a substitution reaction involving 2-chloro-2-methylbut-3-yne. *Indian J. Chem. B: Org. Chem.* **1976**, *14*, 518-521.
31. Chênevert, R.; Pagé, J.; Plante, R.; Beaucage, D. Synthesis of 4,4-dimethyl-5-methylene-4,5-dihydrofurans. *Synthesis* **1982**, 75-77.
32. Chênevert, R.; Pagé, J.; Voyer, N. Synthesis of ($\pm$)-dehydroxyglaupalol and analogs. *Synth. Commun.* **1984**, *14*, 737-742.
33. Reisch, J.; Dharmaratne, H.R.W. A convenient synthesis of the 2-dimethyl-2H-chromene system. *Z. Naturforsch. B* **1985**, *40*, 636-638.
34. Mitra, J.; Mitra, A.K. Palladium(II) assisted cyclization of hydroxyallylcoumarins. *Indian J. Chem. B: Org. Chem.* **1994**, *33*, 276-279.
35. Berger, S.; Haak, E. Ruthenium-catalyzed addition of carboxylic acids or cyclic 1,3-dicarbonyl compounds to propargylic alcohols. *Tetrahedron Lett.* **2010**, *51*, 6630-6634.
36. Cadierno, V.; Díez, J.; García-Garrido, S.E.; Gimeno, J. $[\text{Ru}(\eta^3\text{-2-C}_3\text{H}_4\text{Me})(\text{CO})(\text{dppf})][\text{SbF}_6]$: A mononuclear $16e^-$ ruthenium(II) catalyst for propargylic substitution and isomerization of $\text{HCCPh}_2(\text{OH})$. *Chem. Commun.* **2004**, 2716-2717.
37. Midland, M.M. Preparation of monolithium acetylide in tetrahydrofuran. Reaction with aldehydes and ketones. *J. Org. Chem.* **1975**, *40*, 2250-2252.
38. *CrysAlis^{Pro} CCD & CrysAlis^{Pro} RED*; Oxford Diffraction Ltd.: Oxford, UK, 2008.
39. Farrugia, L.J. *WinGX* suite for small-molecule single-crystal crystallography. *J. Appl. Crystallogr.* **1999**, *32*, 837-838.
40. Altomare, A.; Cascarano, G.; Giacovazzo, C.; Guagliardi, A. Completion and refinement of crystal structures with *SIR92*. *J. Appl. Crystallogr.* **1993**, *26*, 343-350.
41. Sheldrick, G.M. *SHELXL97: Program for the Refinement of Crystal Structures*; University of Göttingen: Göttingen, Germany, 1997.
42. *International Tables for X-Ray Crystallography*; Kynoch Press: Birmingham, UK, 1974, Volume IV (present distributor: Kluwer Academic Publishers: Dordrecht, The Netherlands).
43. Spek, A.L. *PLATON: A multipurpose Crystallographic Tool*; University of Utrecht: Utrecht, The Netherlands, 2006.

Sample Availability: Samples of the compounds **4a–d** and **5b** are available from the authors.