

Article

Chiral Zn(II)-Bisamidine Complex as a Lewis-Brønsted Combined Acid Catalyst: Application to Asymmetric Mukaiyama Aldol Reactions of α -Ketoesters

Ryo Gotoh and Masahiro Yamanaka *

Department of Chemistry and Research Center for Smart Molecules, Rikkyo University, Toshima-Ku, Tokyo 171-8501, Japan

* Author to whom correspondence should be addressed; E-Mail: myamanak@rikkyo.ac.jp; Tel./Fax: +81-3-3985-2395.

Received: 11 July 2012; in revised form: 21 July 2012 / Accepted: 21 July 2012 /

Published: 30 July 2012

Abstract: Focusing on the steric and electronic properties of the resonance-stabilized amidine framework, a cationic metal-bisamidine complex was designed as a conjugated combined Lewis-Brønsted acid catalyst. The chiral Zn(II)-bisamidine catalyst prepared from the 2,2'-bipyridyl derived bisamidine ligand, ZnCl₂, and AgSbF₆ promoted asymmetric Mukaiyama aldol reaction of α -ketoester and α,α -disubstituted silyl enol ether to afford the α -hydroxyester having sequential quarternary carbons in good yield, albeit with low enantioselectivity. Addition of 1.0 equivalent of the fluoroalcohol having suitable acidity and bulkiness dramatically increased the enantioselectivity (up to 68% *ee*). DFT calculations suggested that this additive effect would be caused by self-assembly of the fluoroalcohol on the Zn(II)-bisamidine catalyst.


Keywords: Lewis-Brønsted combined acid; bisamidine; asymmetric Mukaiyama aldol reaction; α -ketoester

1. Introduction

Lewis-Brønsted combined acid catalysts promote higher catalytic activity and stereoselectivity than the individual acid catalysts through the dual effects of enhancement of Lewis acidity and the intramolecular hydrogen-bonding interaction of additional Brønsted acid sites (Figure 1a) [1]. Since the late 1990s, various types of Brønsted acid-assisted chiral Lewis acid catalysts have been developed


and are regarded as an efficient catalyst design for asymmetric reactions [2–14]. Based on Yamamoto's combined acid concept, we proposed double activation phenomena of carbonyl compounds caused by simultaneous coordination of Lewis acid and Brønsted acid sites of the combined acid catalyst. In contrast to the previously reported combined acid catalysts, in which doubly coordination of both acid sites would form a labile four-membered complex (Figure 1b), the three-centered conjugated scaffold would form a more stable six-membered complex (Figure 1c).

Figure 1. (a) Schematic concept of Lewis-Brønsted combined acid catalyst; (b) Double activation by the normal combined acid; (c) Double activation by the conjugated combined acid.



Focusing on the steric and electronic properties of the resonance-stabilized amidine framework, we designed a novel conjugated Lewis-Brønsted combined acid catalyst. Amidinium cation is significantly stabilized by delocalization of positive charge on two N atoms. The electronic flexibility of the amidine framework should induce similar electronic delocalization in the cationic metal amidine complex. When coordinating of cationic metal ions (M^{n+}) instead of proton (H^+), positive charge in the metal amidine complex could be delocalized to enhance Brønsted acidity of the NH moiety in the parent amidine. We eventually designed the cationic metal bisamidine catalyst, which should be thermodynamically stabilized by bidentate coordination of two amidine moieties (Figure 2).

Figure 2. Design of the metal bisamidine complex as a Lewis-Brønsted combined acid catalyst.




Various chiral bisamidines have been recently developed as chiral Brønsted acid catalysts [15–19] and chiral ligands [20]. To our best knowledge, however, there has been no report of bisamidine-based combined acid catalysis. For our initial studies, our designed cationic Zn(II)-bisamidine catalyst was applied in the reaction of an α -ketoester and an α,α -disubstituted silyl enol ether to overcome the present limitations of the asymmetric Mukaiyama aldol reaction [21]. The catalytic asymmetric Mukaiyama aldol reaction of α -ketoesters has been recently identified as an efficient synthetic tool for

making a chiral quaternary carbon center. Evans has reported pioneering studies of asymmetric addition of thioester-derived enolsilanes to pyruvate esters catalyzed by the chiral Cu(II)-bisoxazoline complexes [22–24]. Bolm has also reported highly enantioselective additions of various ketone-derived enolsilanes to methyl pyruvate catalyzed by a chiral Cu(II)-oxazolanyl sulfoximine [25,26]. However, the use of aryl substituted α -ketoesters or ester-derived enolsilanes is still rather limited. Hoveyda developed the chiral AgF₂-pyridyl Schiff base catalyzed reaction of alkyl/aryl substituted α -ketoesters [27]. Pagenkopf reported a significant expansion in substrate scope to include ester-derived enolsilanes and alkyl/aryl substituted α -ketoesters [28]. Despite the synthetic value of α -hydroxyesters having sequential quaternary carbons, the reactions of α -ketoesters and α,α -disubstituted silyl enol ethers have been quite rare [29,30].

2. Results and Discussion

As a preliminary computational study, we investigated the dual effects of Lewis and Brønsted acids on the cationic metal bisamidine complex by DFT calculations (Figure 3) [31]. In spite of the structurally stable aromatic ring in the 2,2'-bipyridine framework, the difference of the bond lengths between *a* and *b* in the amidine moiety becomes smaller upon coordination of the Zn(II) cation. The natural charge of the NH moiety in **Zn²⁺-L** (−0.171) was more positive than in **L** (−0.206). Furthermore, **Zn²⁺-Bipy** exhibited a greater degree of the positive charge on the Zn(II) center (+1.545) than **Zn²⁺-L** (+1.433). These natural charge differences suggested delocalization of the positive charge via the amidine framework to increase Brønsted acidity of the NH proton. In a manner similar to the amidinium complex, the efficient charge delocalization in the amidine moiety was also induced in **Zn²⁺-L**. These DFT calculations predicted that the cationic Zn(II)-bisamidine complex such as **Zn²⁺-L** would have a unique catalytic ability incorporating Lewis acid and Brønsted acid functions.


Figure 3. DFT calculations of bisamidine ligand (**L**), Zn(II)-bisamidine (**Zn²⁺-L**), and Zn(II)-bipyridine (**Zn²⁺-Bipy**) complexes (B3LYP/LANL2DZ for Zn, 6-31G* for the rest). Bond lengths are shown in Å. NBO charges are shown in *italics*.



First, we compared the catalytic activities of the cationic metal complex prepared with **L1** and **L2**, ZnCl₂, and 2.0 equivalents of AgSbF₆ under stoichiometric conditions (Table 1). In the Zn(II), Cu(II), and Fe(II)-catalyzed Mukaiyama aldol reaction of methyl benzoylformate (**1a**) and dimethylketene methyl trimethylsilyl acetal (**2**), **L1** derived catalysts gave the product **3a** in better yields than the corresponding **L2** derived catalyst. This indicates that the presence of the amidine unit is essential for

the higher catalytic activity, which would be caused by the Brønsted acid function at the NH moiety. Based on the promising results of the Zn(II)-**L1** catalyst, various counteranions, SbF_6^- , BF_4^- , and TfO^- were explored under catalytic conditions (the reaction was carried out in CH_2Cl_2 at -78°C for 15 h in the presence of 10 mol% Zn(II)-**L1**). The highest yield of 89% was achieved by use of SbF_6^- (BF_4^- : 68%, TfO^- : 57%).

Table 1. Metal-bisamidine catalyzed reaction of **1a** and **2** under stoichiometric condition.



Notes: ^a Determined by $^1\text{H-NMR}$ analysis of crude mixture; ^b 10 mol% of Zn(II)-**L1** catalyst was used for 15 h.

The (*R*)-DABN derived chiral bisamidine ligand **L3** was developed for the catalytic asymmetric version (Table 2). The asymmetric Mukaiyama aldol reaction of **1a** and **2** was catalyzed by 10 mol% of Zn(II)-**L3** to afford **3a** in moderate yield with 26% *ee* (Entry 1). On the other hand, both Zn(II)-**L4** and Zn(II)-**L5** catalysts proved much less selective (Entries 2 and 3). The NMe_2 group in the DABN moiety performed better than either smaller or larger amino groups. In some cases, it has been recognized that achiral additives can be beneficial for catalytic activity and the stereocontrol ability of chiral catalysts [32]. Fluoroalcohol, in particular, has emerged as an efficient additive for promoting catalyst turnover in addition reactions of enolsilanes [26,33–35]. These reports encouraged us to examine the additive effect of trifluoroethanol (TFE), which acts as a proton source to regenerate the active catalyst or coordinates on the active catalyst to form a more efficient chiral environment at the reaction centre. Addition of 20 mol% of TFE slightly improved yield (71%) and enantioselectivity (35% *ee*, Entry 4). The enantioselectivity was increased with increasing of the amount of the additional TFE. Addition of 1.0 equivalent of TFE provided the optimal enantioselectivity of up to 50% *ee* (Entry 5), while an excess amount of TFE would affect solvent polarity to decrease the enantioselectivity (Entry 6). We predicted that the present additive effect would be caused by self-assembly of the fluoroalcohol on the Zn(II)-**L3** catalyst. To investigate our prediction, DFT calculations for the reactive complex models of the Zn(II)-**L3** and α -ketoester with/without TFE were carried out (Figures 4 and 5).

Figure 4. Schematic geometries for the assembled complex models (A–F) of Zn(II)-L3 and α -ketoester with TFE and the relative energies (B3LYP/LANL2DAZ for Zn, 6-31G* for the rest).






Figure 5. DFT calculations for the TFE addition effect in the complex of Zn(II)-L3 and α -ketoester (B3LYP/LANL2DAZ for Zn, 6-31G* for the rest). Bond lengths are shown in Å.



Focusing on the TFE coordination to Zn(II)-**L3** and α -ketoester, methyl pyruvate was used in the chemical models to reduce the computational costs. Six assembled complex models (**A–F**) were optimized (B3LYP/LANL2DAZ for Zn, 6-31G* for the rest). The assembled complex **A** found to be most stable (Figure 4). The structural properties of the parent complex **G** were dramatically changed by coordination with two molecules of TFE (Figure 5). Whereas **G** has the tetrahedral Zn(II) centre, an octahedral hexacoordinated Zn(II) centre was formed in **A**. The hydrogen-bonding between F and binaphthyl hydrogen atoms (2.42 Å, 2.47 Å) and the Zn-O interaction (2.24 Å, 2.27 Å) stabilized **A** by 7.4 kcal/mol in Gibbs free energy. These structural changes would enhance enantiofacial differentiation at the carbonyl group (dotted purple circles in Figure 5). There is no enantiofacial differentiation at the carbonyl group in **G**. In contrast, *Si*-facial attack of **2** in **A** would be prevented by the steric hindrance of TFE to increase the enantioselectivity. This computational prediction was supported by the stoichiometric reaction with 2.0 equivalents of TFE yielding **3a** with 46% *ee* (almost the same enantioselectivity under catalytic condition, Entry 5 in Table 2). To further improve the enantioselectivity, other alcohol additives were examined (Scheme 1).

Table 2. Metal-bisamidine catalyzed reaction of **1a** and **2** under stoichiometric conditions.




Entry	Ligand	Y eq. of TFE	Yield (%) ^a	ee (%) ^b
1	L3	-	60	26
2	L4	-	81	1
3	L5	-	53	0
4	L3	0.2	71	35
5	L3	1.0	69	50
6	L3	5.0	68	33

Notes: ^a Determined by ¹H-NMR analysis of the reaction crude mixtures. ^b Determined by chiral HPLC analysis.


Whereas ethanol (**b**), trichloroethanol (**c**), and phenol derivatives (**d** and **e**) exhibited no positive impact on the enantioselectivity, 2,2,3,3,3-pentafluoro-1-propanol (**f**) slightly increased the enantioselectivity (40% *ee*). Aliphatic fluorinated alcohols yielded promising results in the initial screening. After investigation of various aliphatic fluorinated alcohols, hexafluoroisopropyl alcohol (HFIP, **g**) was found to be the most efficient alcohol additive and the highest enantioselectivity was achieved (68% *ee*). A more sterically demanding alcohol, 1,1,1,3,3,3-hexafluoro-2-methyl-2-propanol

(**h**), exhibited lower enantioselectivity than HFIP (34% *ee*). The more bulky and acidic alcohol tris(trifluoromethyl)methanol (**i**), promoted the achiral pathway to exhibit no enantioselectivity. Therefore, aliphatic fluorinated alcohols having suitable acidity and bulkiness exerted a significant impact on the enantiofacial control of Zn(II)-**L3**. A similar additive effect was observed when methyl pyruvate (**1b**) was used as electrophile. The Zn(II)-**L3** catalyzed reaction of **1b** and **2** readily proceeds at $-78\text{ }^{\circ}\text{C}$ to afford **3b** in 95% yield with 11% *ee*. Addition of 1.0 equivalent of HFIP led to the same yield (95%) and an increase in the enantioselectivity to 41% *ee* (Scheme 2). The lower *ee* value of **3b** than **3a** indicates that Me group would be too small to achieve the enantiofacial discrimination controlled by the Zn(II)-**L3** catalyst.

Scheme 1. Screening of alcohol additives.



Scheme 2. Zn(II)-**L3** catalyzed reaction of **1b** and **2** with or without HFIP.



3. Experimental

3.1. General

All reactions were performed under nitrogen atmosphere with magnetic stirring in dried glassware. AgOTf, AgBF₄, ZnCl₂, CuCl₂, FeCl₂ 4H₂O, were purchased from Sigma-Aldrich Japan K. K. (Tokyo,

Japan). Dimethylketene methyl trimethylsilyl acetal was purchased from Tokyo Chemical Industry Co., Ltd. (Tokyo, Japan). (*R*)-1,1'-binaphthyl-2,2'-diamine ((*R*)-DABN), methyl phenylglyoxylate, methyl pyruvate were purchased from Wako Pure Chemical Industries, Ltd. (Tokyo, Japan) All dehydrated solvents were purchased from Kanto Chemical Co., Inc. (Tokyo, Japan) Enantiomeric excess was determined on a Shimadzu HPLC system with UV detection or a Shimadzu GC system. ^1H -NMR and ^{13}C -NMR spectra were recorded on a JEOL JNM-ECX400 spectrometer. Chemical shifts in CDCl_3 were reported in ppm from tetramethylsilane as the internal standard (CDCl_3 : $\delta = 0$) for ^1H -NMR and from the solvent resonance (CDCl_3 : $\delta = 77.0$) for ^{13}C -NMR. Infrared (IR) spectra were recorded on a JASCO FT/IR-230 Fourier transform infrared spectrophotometer. High resolution mass was performed by JEOL JMS-GC mate II. Purification of the products was performed by column chromatography on silica gel 60N (spherical, neutral, Kanto Chemical Co., Inc), Wakogel 50NH₂ (Wako Pure Chemical Industries, Ltd.) or activated alumina (about 75 μm).

3.2. Synthesis of the 2,2'-Bipyridyl Derived Bisamidine Ligands

3.2.1. Synthesis of 6,6'-Dibromo-2,2'-dipyridyl

To a solution of 2,6-dibromopyridine (1.17 g, 5.0 mmol) in Et_2O (10 mL), *n*-BuLi (1.67 M solution in hexane, 6.6 mL, 11 mmol) was added dropwise at -78°C over 1 h, and then CuCl_2 (1.01 g, 7.5 mmol) was added to the reaction mixture. After stirring for 30 min, the reaction mixture was bubbled by O_2 for 24 h at -78°C . To the reaction mixture 6 N HCl aq. (15 mL) was added, and then filtered through Büchner funnel. The residue was purified by recrystallization from AcOEt gave 6,6'-dibromo-2,2'-dipyridyl as a white powder in 28% yield.

3.2.2. Synthesis of the (*R*)-DABN Derivatives

(*R*)-*N,N*-Dimethyl-1,1'-binaphthalene-2,2'-diamine [(*R*)-NMe₂-DABN] was prepared according to the previously reported method [36]. To a solution of (*R*)-1,1'-binaphthalene-2,2'-diamine [(*R*)-DABN] (516.0 mg, 2.0 mmol) and AcOH (1.2 mL, 20 mmol) in CH_2Cl_2 (10 mL) was added acetic anhydride (0.2 mL, 2.0 mmol) at 0°C under N_2 . The reaction mixture was stirred for 20 h at room temperature, and then 1 N NaOH aq. was added to adjust the solution to $\text{pH} > 7$. The solution was extracted by CH_2Cl_2 and the combined organic layers were washed with brine, dried over Na_2SO_4 , and then concentrated under reduced pressure. The crude mixture was purified by column chromatography on silica gel 60N (AcOEt/hexane = 2/1) to afford *N*-[(1*R*)-2'-amino-1,1'-binaphthalen-2-yl]acetamide [(*R*)-Ac-DABN] as a colorless oil in 75% yield. To a solution of (*R*)-Ac-DABN (0.49 g, 1.5 mmol) and aqueous formaldehyde (37%, 1.48 mL, 17.5 mmol) in THF (19 mL) was added NaBH_3CN (0.19 mg, 10.3 mmol). The reaction mixture was stirred for 15 min at room temperature followed by addition of AcOH (1.9 mL). After stirring for 4 h at room temperature, 1 N NaOH aq. was added to adjust the solution to $\text{pH} > 7$. The solution was extracted by CH_2Cl_2 and the combined organic layers were washed with brine, dried over Na_2SO_4 , and then concentrated under reduced pressure. The crude mixture was purified by column chromatography on silica gel 60N (AcOEt/hexane = 1/5) to afford the *N*-[(1*R*)-2'-(dimethylamino)-1,1'-binaphthalen-2-yl]acetamide [(*R*)-Ac-NMe₂-DABN] as a brown powder in quantitative yield. To a solution of (*R*)-Ac-NMe₂-DABN (529 mg, 1.5 mmol) in EtOH

(5 mL) was added 4 N HCl (5 mL). The reaction mixture was stirred for 24 h at 75 °C, and then 1 N NaOH aq. was added to adjust the solution to pH > 7. The solution was extracted by CH₂Cl₂ and the combined organic layers were washed with brine, dried over Na₂SO₄, and then concentrated under reduced pressure. The crude mixture was purified by column chromatography on silica gel 60N (AcOEt/hexane = 1/10) to afford (1*R*)-*N,N*-dimethyl-1,1'-binaphthalene-2,2'-diamine [(*R*)-NMe₂-DABN] as a white powder in 90% yield.

To a solution of (*R*)-Ac-DABN (93 mg, 0.29 mmol) in DMF (1.5 mL), 1,5-dibromopentane (345 mg, 1.5 mmol) and Et₃N (0.3 mL, 3.0 mmol) were added. The reaction mixture was stirred for 2 ddy at 150 °C followed by addition of water (5 mL). The solution was extracted by CH₂Cl₂ and the combined organic layers were washed with brine, dried over Na₂SO₄, and then concentrated under reduced pressure. To a solution of the crude residue in EtOH (2 mL) was added 6 N HCl aq. (2 mL). After stirring for 24 h at 85 °C, 1 N NaOH aq. was added to adjust the solution to pH > 7. The solution was extracted by CH₂Cl₂ and the combined organic layers were washed with brine, dried over Na₂SO₄, and then concentrated under reduced pressure. The crude mixture was purified by column chromatography on silica gel 60N (AcOEt/hexane = 1/10) to afford (1*R*)-2'-(1-piperidiny)-1,1'-binaphthalen-2-amine [(*R*)-C₅H₁₀N-DABN] as a white powder in 66% yield.

3.2.3. Synthesis of **L1**, **L3**, **L4**, and **L5**

2,2'-Bipyridyl linked bisamidines **L1**, **L3**, **L4**, and **L5** were prepared according to the C-N coupling method reported by Buchwald [37]. A solution of Pd(dba)₂ (6.8 mg, 0.012 mmol) and racemic-BINAP (14.8 mg, 0.024 mmol) in toluene (2 mL) under N₂ was stirred for 20 min at room temperature. 6,6'-Dibromo-2,2'-dipyridyl (153.5 mg 0.49 mmol), ^tBuONa (119.5 mg, 1.25 mmol) and the primary amine (**L1**: BnNH₂, **L3**: (*R*)-NMe₂-DABN, **L4**: (*R*)-Ac-DABN, **L5**: (*R*)-C₅H₁₀N-DABN, 1.0 mmol) were added to the solution at room temperature followed by stirring or 24 h at 110 °C. The solution was filtered through Celite and concentrated under reduced pressure.

For **L1**, **L3**, and **L5**: The crude mixture was purified by column chromatography on activated alumina (**L1**: AcOEt/CH₂Cl₂ = 1/1, **L3**: AcOEt/hexane = 1/3) or silica gel 60N (**L5**: AcOEt/hexane = 1/15) to afford the 2,2'-bipyridyl linked bisamidine as a white powder (**L1**: 99%, **L3**: 99%, **L5**: 51%).

For **L4**: To a solution of the crude residue in EtOH (2 mL) was added 6 N HCl aq. (2 mL). After stirring for 24 h at 85 °C, 1 N NaOH aq. was added to adjust the solution to pH > 7. The solution was extracted by CH₂Cl₂ and the combined organic layers were washed with brine and dried over Na₂SO₄. The crude mixture was purified by column chromatography on silica gel 60N (AcOEt/ hexane = 1/3) to afford **L4** as a white powder in 99% yield.

L1: ¹H-NMR (400 MHz, CDCl₃): δ 7.64 (d, *J* = 7.6 Hz, 2H), 7.50 (m, 2H) 7.40 (d, *J* = 7.2 Hz, 4H), 7.34 (m, 4H), 7.28 (m, 2H), 6.38 (d, *J* = 7.6 Hz, 2H), 4.97 (br, 2H), 4.60 (s, 4H); ¹³C-NMR (100 MHz, CDCl₃): δ 158.3, 139.9, 138.5, 128.9, 127.8, 127.8, 127.4, 110.8, 107.2, 46.7; IR ν 3392, 3055, 2831, 2785, 1572, 1500, 1429, 1335 cm⁻¹; HRMS Calcd for C₂₄H₂₃N₄ [M+H]: 367.1923; Found: 367.1948.

L3: ¹H-NMR (400 MHz, CDCl₃): δ 8.28 (d, *J* = 9.2 Hz 2H), 7.90 (m, 6H), 7.81 (d, *J* = 8.4 Hz, 2H), 7.67 (m, 2H), 7.47 (m, 4H), 7.33(m, 2H), 7.27 (m, 2H), 7.20 (m, 2H), 7.12 (m, 6H), 6.68 (s, 2H), 6.63 (d, *J* = 8.0 Hz, 2H), 2.51 (s, 12H); ¹³C-NMR (100 MHz, CDCl₃): δ 155.6, 155.0, 150.4, 138.3, 137.2,

134.4, 134.2, 130.5, 130.0, 129.9, 128.4, 128.3, 128.2, 126.8, 126.6, 126.4, 125.6, 124.3, 124.1, 124.0, 122.6, 121.7, 119.6, 112.6, 109.5, 43.6; IR ν 3395, 3054, 2930, 2867, 2831, 2784, 1571, 1501, 1442, 1428, 1334 cm^{-1} ; HRMS Calcd for $\text{C}_{54}\text{H}_{45}\text{N}_6$ $[\text{M}+\text{H}]$: 777.3706; Found: 777.3377.

L4: ^1H -NMR (400 MHz, CDCl_3): δ 8.57 (d, J = 8.8 Hz 2H), 7.98 (d, J = 8.8 Hz, 2H) 7.88 (d, J = 8.0 Hz, 2H), 7.80 (m, 4H), 7.74 (m, 2H), 7.48 (m, 2H), 7.34 (m, 2H), 7.24–7.04 (m, 12H), 6.53 (d, J = 7.6 Hz, 2H), 6.36 (s, 2H), 3.67 (s, 4H); ^{13}C -NMR (100 MHz, CDCl_3): δ 154.8, 154.6, 142.9, 138.2, 137.9, 133.9, 133.3, 130.2, 129.9, 128.7, 128.3, 128.2, 128.1, 127.0, 126.7, 125.0, 124.0, 123.9, 122.5, 120.5, 119.0, 118.2, 113.0, 111.6, 110.5; IR ν 3470, 3381, 3051, 1617, 1463, 1335, 1310, 1270 cm^{-1} ; HRMS Calcd for $\text{C}_{50}\text{H}_{37}\text{N}_6$ $[\text{M}+\text{H}]$: 721.3080; Found: 721.3436.

L5: ^1H -NMR (400 MHz, CDCl_3): δ 8.30 (d, J = 8.8 Hz, 2H), 7.93 (m, 4H) 7.84 (m, 4H), 7.69 (d, J = 7.6 Hz, 2H), 7.48 (m, 4H), 7.32 (m, 4H), 7.20–7.05 (m, 8H), 7.00 (d, J = 4.0 Hz, 2H), 6.65 (d, J = 8.0 Hz, 2H), 2.85 (m, 8H), 1.26–1.17 (m, 12H); ^{13}C -NMR (100 MHz, CDCl_3): δ 155.1, 154.8, 150.4, 138.0, 136.6, 134.1, 133.8, 130.2, 130.1, 129.6, 127.9, 127.9, 126.3, 126.1, 126.1, 125.7, 125.1, 124.0, 123.6, 123.3, 121.1, 121.1, 120.0, 112.1, 109.3, 52.9, 26.1, 24.1; IR ν 3393, 3054, 2929, 2850, 2796, 1572, 1501, 1443, 1429, 1337 cm^{-1} ; HRMS Calcd for $\text{C}_{60}\text{H}_{53}\text{N}_6$ $[\text{M}+\text{H}]$: 857.4332; Found: 857.4039.

3.3. General Procedure for Asymmetric Mukaiyama Aldol Reaction of **1a** and **2** (Scheme 1)

Under a N_2 atmosphere, a solution of ZnCl_2 (1.4 mg, 0.01 mmol) and **L3** (7.8 mg, 0.01 mmol) in CH_3CN was stirred at room temperature for 20 min. The solvent was removed under reduced pressure and AgSbF_6 (6.9 mg, 0.02 mmol) was added under N_2 . The residue was stirred in CH_2Cl_2 (0.5 mL) at room temperature for 30 min followed by addition of HFIP (16.8 mg, 0.1 mmol). After stirring at room temperature for 20 min, **1a** (16.4 mg, 0.1 mmol), **2** (17.9 mg, 0.1 mmol) were added to the solution and the mixture was stirred at -45°C . The reaction was quenched by filtration through silica gel 60N. After concentration under reduced pressure, the residue was dissolved in THF (0.5 mL). 6 N HCl aq. was added and the solution was stirred at 65°C . The solution was extracted by CH_2Cl_2 and the combined organic layers were washed with brine, dried over Na_2SO_4 , and then concentration under reduced pressure. The crude mixture was purified by column chromatography on silica gel 60N (AcOEt/Hexane = 1/15) to afford **3a** as white crystals (63% yield).

Dimethyl-2,2-dimethyl-3-hydroxy-3-phenylbutanedioate (3a) [38]. **3a** is a known compound. white crystals: IR ν 3474, 2955, 1706, 1442, 1356 cm^{-1} ; ^1H -NMR (400 MHz, CDCl_3): δ 7.57 (d, J = 6.4 Hz, 2H), 7.37–7.30 (m, 3H), 4.70 (s, 1H), 3.81 (s, 3H), 3.74 (s, 3H), 1.36 (s, 3H), 1.13 (s, 3H); ^{13}C -NMR (100 MHz, CDCl_3) δ = 178.7, 174.3, 136.9, 128.0, 127.5, 127.3, 81.9, 52.9, 52.4, 50.4, 22.7, 21.5; EI-MS m/z = 266 $[\text{M}]^+$ for $\text{C}_{14}\text{H}_{18}\text{O}_5$; Enantiomeric excess was determined by chiral HPLC analysis (68% *ee*, DAICEL CHIRALPAK OD-H, 2-propanol/*n*-hexane = 5/95, flow rate 1.0 mL/min, wavelength = 220 nm, t_{minor} = 5.9 min and t_{major} = 7.5 min).

Dimethyl-3-hydroxy-2,2,3-trimethylbutanedioate (3b) [38]. **3b** is a known compound. pale yellow oil.: IR ν 3511, 2954, 1728, 1439, 1374 cm^{-1} ; ^1H -NMR (400 MHz, CDCl_3): δ 3.94 (s, 1H), 3.77 (s, 3H),

3.70 (s, 3H), 1.43 (s, 3H), 1.28 (s, 6H). ^{13}C -NMR (100 MHz, CDCl_3) δ = 176.8, 175.9, 78.0, 52.7, 52.2, 49.3, 21.6, 20.9, 20.6; EI-MS m/z = 205 $[\text{M}+\text{H}]^+$ for $\text{C}_9\text{H}_{16}\text{O}_5$; Enantiomeric excess was determined by chiral GC analysis (41% *ee*, Cyclodex- β column, 100 $^\circ\text{C}$, t_{minor} = 37.5 min and t_{major} = 38.4 min).

4. Conclusions

In summary, we developed asymmetric Mukaiyama aldol reaction of α,α -disubstituted silyl enol ether and α -ketoester promoted by the rationally designed chiral Zn(II)-bisamidine catalyst, which exhibited an unique catalytic ability incorporating Lewis acid and Brønsted acid functions. Aliphatic fluorinated alcohols having suitable acidity and bulkiness dramatically enhanced the enantioselectivity. DFT calculations revealed self-assembly of the fluorinated alcohol exerted a significant impact on the active catalyst structure to enhance enantiofacial control. Although DFT calculations predicted *Re*-facial attack leading to the *R* configured product would be preferred, the absolute configuration of the product has not been determined yet. Studies on further improvement of the cationic metal-bisamidine catalyst system and determination of the absolute configuration of the product are currently underway.

Supplementary Materials

Supplementary materials can be accessed at: <http://www.mdpi.com/1420-3049/17/8//9010/s1>.

Acknowledgments

This work was supported by a Grant-in-Aid for Young Scientists (B) from JSPS (Grant No. 20750082).

References

1. Yamamoto, H.; Futatsugi, K. “Designer acids”: Combined acid catalysis for asymmetric synthesis. *Angew. Chem. Int. Ed. Engl.* **2005**, *44*, 1924–1942.
2. Maruoka, K.; Sakurai, M.; Fujiwara, J.; Yamamoto, H. Asymmetric Diels-Alder reaction directed toward chiral anthracycline intermediates. *Tetrahedron* **1986**, *27*, 4895–4898.
3. Furuta, K.; Miwa, Y.; Iwanaga, K.; Yamamoto, H. Acyloxyborane: An activating device for carboxylic acids. *J. Am. Chem. Soc.* **1988**, *110*, 6254–6255.
4. Ishihara, K.; Miyata, M.; Hattori, K.; Tada, T.; Yamamoto, H.Y. A new chiral BLA promoter for asymmetric aza Diels-Alder and Aldol-type reactions of imines. *J. Am. Chem. Soc.* **1994**, *116*, 10520–10524.
5. Ishihara, K.; Yamamoto, H. Brønsted acid assisted chiral Lewis acid (BLA) catalyst for asymmetric Diels-Alder reaction. *J. Am. Chem. Soc.* **1994**, *116*, 1561–1562.
6. Ishihara, K.; Kurihara, H.; Yamamoto, H. A new powerful and practical BLA catalyst for highly enantioselective Diels-Alder reaction: An extreme acceleration of reaction rate by Brønsted acid. *J. Am. Chem. Soc.* **1996**, *118*, 3049–3050.
7. Ishihara, K.; Kurihara, H.; Matsumoto, M.; Yamamoto, H. Design of Brønsted acid-assisted chiral Lewis acid (BLA) catalysts for highly enantioselective Diels-Alder reactions. *J. Am. Chem. Soc.* **1998**, *120*, 6920–6930.

8. Corey, E.J.; Shibata, T.; Lee, T.W. Asymmetric Diels-Alder reactions catalyzed by a triflic acid activated chiral oxazaborolidine. *J. Am. Chem. Soc.* **2002**, *124*, 3808–3809.
9. Ryu, D.H.; Lee, T.W.; Corey, E.J. Broad-spectrum enantioselective Diels-Alder catalysis by chiral, cationic oxazaborolidines. *J. Am. Chem. Soc.* **2002**, *124*, 9992–9993.
10. Ryu, D.H.; Corey, E.J. Triflimide activation of a chiral oxazaborolidine leads to a more general catalytic system for enantioselective Diels-Alder addition. *J. Am. Chem. Soc.* **2003**, *125*, 6388–6390.
11. Payette, J.N.; Yamamoto, H. Regioselective and asymmetric Diels-Alder reaction of 1- and 2-substituted cyclopentadienes catalyzed by a Brønsted acid activated chiral oxazaborolidine. *J. Am. Chem. Soc.* **2007**, *129*, 9536–9537.
12. Hu, G.; Huang, L.; Huang, R.H.; Wulff, W.D. Evidence for a boroxinate based Brønsted acid derivative of VAPOL as the active catalyst in the catalytic asymmetric aziridination reaction. *J. Am. Chem. Soc.* **2009**, *131*, 15615–15617.
13. Rauniyar, V.; Hall, D.G. Rationally improved chiral Brønsted acid for catalytic enantioselective allylboration of aldehydes with an expanded reagent scope. *J. Org. Chem.* **2009**, *74*, 4236–4241.
14. Lv, J.; Li, X.; Zhong, L.; Luo, S.; Cheng, J.-P. Asymmetric binary-acid catalysis with chiral phosphoric acid and MgF₂: catalytic enantioselective Friedel-Crafts reactions of β,γ -unsaturated α -ketoesters. *Org. Lett.* **2010**, *12*, 1096–1099.
15. Nugent, B.M.; Yoder, R.A.; Johnston, J.N. Chiral proton catalysis: A catalytic enantioselective direct Aza-Henry reaction. *J. Am. Chem. Soc.* **2004**, *126*, 3418–3419.
16. Singh, A.; Yoder, R.A.; Shen, B.; Johnston, J.N. Chiral proton catalysis: Enantioselective Brønsted acid catalyzed additions of nitroacetic acid derivatives as glycine equivalents. *J. Am. Chem. Soc.* **2007**, *129*, 3466–3467.
17. Davis, T.A.; Wilt, J.C.; Johnston, J.N. Bifunctional asymmetric catalysis: Amplification of Brønsted basicity can orthogonally increase the reactivity of a chiral Brønsted acid. *J. Am. Chem. Soc.* **2010**, *132*, 2880–2882.
18. Akalay, D.; Dürner, G.; Bats, J.W.; Bolte, M.; Göbel, M.W. Synthesis of C₂-Symmetric Bisamidines: A New Type of Chiral Metal-Free Lewis Acid Analogue Interacting with Carbonyl Groups. *J. Org. Chem.* **2007**, *72*, 5618–5624.
19. Akalay, D.; Dürner, G.; Bats, J. W.; Göbel, M.W. C₂-symmetric bisamidines: Chiral Brønsted bases catalysing the Diels-Alder reaction of anthrones. *Beilstein J. Org. Chem.* **2008**, *4*, doi:10.3762/bjoc.4.28.
20. Miyata, K.; Kutsuna, H.; Kawakami, S.; Kitamura, M. A Chiral Bidentate sp²-N Ligand, Naph-diPIM: Application to CpRu-Catalyzed Asymmetric Dehydrative C-, N-, and O-Allylation. *Angew. Chem. Int. Ed. Engl.* **2011**, *50*, 4649–4653.
21. Catalytic enantioselective aldol additions to ketones, see: Adachi, S.; Harada, T. Catalytic Enantioselective Aldol additions to ketones. *Eur. J. Org. Chem.* **2009**, 3661–3671.
22. Evans, D.A.; Kozlowski, M.C.; Burgey, C.S.; MacMillan, D.W.C. C₂-symmetric copper(II) complexes as chiral Lewis acids. Catalytic enantioselective aldol additions of enolsilanes to pyruvate esters. *J. Am. Chem. Soc.* **1997**, *119*, 7893–7894.
23. Evans, D.A.; MacMillan, D.W.C.; Campos, K.R. C₂-Symmetric tin(II) complexes as chiral Lewis acids. Catalytic enantioselective *anti* aldol additions of enolsilanes to glyoxylate and pyruvate esters. *J. Am. Chem. Soc.* **1997**, *119*, 10859–10860.

24. Evans, D.A.; Burgey, C.S.; Kozlowski, M.C.; Tregay, S.W. C₂-Symmetric copper(II) complexes as chiral Lewis acids. Scope and mechanism of the catalytic enantioselective aldol additions of enolsilanes to pyruvate esters. *J. Am. Chem. Soc.* **1999**, *121*, 686–699.
25. Langner, M.; Bolm, C. C₁-Symmetric sulfoximines as ligands in copper-catalyzed asymmetric Mukaiyama-type aldol reactions. *Angew. Chem. Int. Ed. Engl.* **2004**, *43*, 5984–5987.
26. Sedelmeier, J.; Hammerer, T.; Bolm, C. C₁-Symmetric oxazolinyl sulfoximines as ligands in copper-catalyzed asymmetric Mukaiyama aldol reactions. *Org. Lett.* **2008**, *10*, 917–920.
27. Akullian, L.C.; Snapper, M.L.; Hoveyda, A.H. A chiral Ag-based catalyst for practical, efficient, and highly enantioselective additions of enolsilanes to α -ketoesters. *J. Am. Chem. Soc.* **2006**, *128*, 6532–6533.
28. Engers, J.L.; Pagenkopf, B.L. A general asymmetric aldol reaction of silyl ketene acetals derived from simple esters to aryl α -keto esters. *Eur. J. Org. Chem.* **2009**, 6109–6111.
29. Ohkouchi, M.; Masui, D.; Yamaguchi, M.; Yamagishi, T. Efficient Mukaiyama aldol reaction by silver(I) carboxylate-bis(phosphine) catalysts. *Nippon Kagaku Kaishi* **2002**, 223–229.
30. Synthesis of α -hydroxyester having sequential quarternary carbons by asymmetric Mukaiyama aldol reaction of methyl pyruvate and cyclopropyl silylketenehemithioacetal, see: Siegel, D.S.; Piizzi, G.; Piersanti, G.; Movassaghi, M. Enantioselective total synthesis of (–)-Acylfulvene and (–)-Irofulven. *J. Org. Chem.* **2009**, *74*, 9292–9304.
31. Frisch, M.J.; Trucks, G.W.; Schlegel, H.B.; Scuseria, G.E.; Robb, M.A.; Cheeseman, J.R.; Montgomery, J.A., Jr.; Vreven, T.; Kudin, K.N.; Burant, J.C.; *et al.* *Gaussian 03*, Revision E.01; Gaussian, Inc.: Wallingford, CT, USA, 2004.
32. Vogl, E.M.; Gröger, H.; Shibasaki, M. Towards perfect asymmetric catalysis: Additives and cocatalysts. *Angew. Chem. Int. Ed. Engl.* **1999**, *38*, 1570–1577.
33. Evans, D.A.; Rovis, T.; Kozlowski, M.C.; Tedrow, J.S. C₂-Symmetric Cu(II) complexes as chiral Lewis acids. Catalytic enantioselective Michael addition of silylketene acetals to alkylidene malonates. *J. Am. Chem. Soc.* **1999**, *121*, 1994–1995.
34. Evans, D.A.; Willis, M.C.; Johnston, J.N. Catalytic enantioselective Michael additions to unsaturated ester derivatives using chiral copper(II) Lewis acid complexes. *Org. Lett.* **1999**, *1*, 865–868.
35. Shuklov, I.A.; Dubrovina, N.V.; Börner, A. Fluorinated alcohols as solvents, cosolvents and additives in homogeneous catalysis. *Synthesis* **2007**, 2925–2943.
36. Wang, J.; Li, H.; Duan, W.; Zu, L.; Wang, W. Organocatalytic asymmetric Michael addition of 2,4-pentandione to nitroolefins. *Org. Lett.* **2005**, *21*, 4713–4716.
37. Wolfe, J.P.; Wagaw, S.; Buchwald, S.L. An Improved Catalyst System for Aromatic Carbon-Nitrogen Bond Formation: The Possible Involvement of Bis(Phosphine) Palladium Complexes as Key Intermediates. *J. Am. Chem. Soc.* **1996**, *118*, 7215–7216.
38. Doucet, H.; Parrain, J.-L.; Santelli, M. Ruthenium(II) and palladium(II) complexes mediated addition of ketene silyl ketal to aldehydes and ketones: Remarkable influence of the nature of the ligand. *Synlett* **2000**, 871–873.