

Supporting Information


Synthesis of Regiospecifically Fluorinated 1,3-Dienamides


Mohammad Chowdhury, Samir K. Mandal, Shaibal Banerjee and Barbara Zajc *


Department of Chemistry, The City College and The City University of New York, 160 Convent Avenue, NY 10031, USA


TABLE OF CONTENTS

Information	Page
¹ H NMR of 2-(benzo[<i>d</i>]thiazol-2-ylsulfonyl)-2-fluoro- <i>N</i> -methyacetamide (1)	S2
¹⁹ F NMR of 2-(benzo[<i>d</i>]thiazol-2-ylsulfonyl)-2-fluoro- <i>N</i> -methyacetamide (1)	S3
¹ H NMR of 2-(2,2-dimethoxyethylthio)benzo[<i>d</i>]thiazole (3)	S4
¹³ C NMR of 2-(2,2-dimethoxyethylthio)benzo[<i>d</i>]thiazole (3)	S5
¹ H NMR of 2-(benzo[<i>d</i>]thiazol-2-ylthio)acetaldehyde (2)	S6
¹ H NMR of (<i>Z</i>)-4-(benzo[<i>d</i>]thiazol-2-ylthio)-2-fluoro- <i>N</i> -methybut-2-enamide (4)	S7
¹³ C NMR of (<i>Z</i>)-4-(benzo[<i>d</i>]thiazol-2-ylthio)-2-fluoro- <i>N</i> -methybut-2-enamide (4)	S8
¹⁹ F NMR of (<i>Z</i>)-4-(benzo[<i>d</i>]thiazol-2-ylthio)-2-fluoro- <i>N</i> -methybut-2-enamide (4)	S9
¹ H NMR of (<i>Z</i>)-4-(benzo[<i>d</i>]thiazol-2-ylsulfonyl)-2-fluoro- <i>N</i> -methoxy- <i>N</i> -methylbut-2-enamide (5)	S10
¹³ C NMR of (<i>Z</i>)-4-(benzo[<i>d</i>]thiazol-2-ylsulfonyl)-2-fluoro- <i>N</i> -methoxy- <i>N</i> -methylbut-2-enamide (5)	S11
¹⁹ F NMR of (<i>Z</i>)-4-(benzo[<i>d</i>]thiazol-2-ylsulfonyl)-2-fluoro- <i>N</i> -methoxy- <i>N</i> -methylbut-2-enamide (5)	S12
¹ H NMR of (<i>Z,4E/Z</i>)- 6a	S13
¹⁹ F NMR of (<i>Z,4E/Z</i>)- 6a	S14
¹ H NMR of (<i>Z,4E/Z</i>)- 6b	S15
¹⁹ F NMR of (<i>Z,4E/Z</i>)- 6b	S16
¹ H NMR of (<i>Z,4E/Z</i>)- 6c	S17
¹⁹ F NMR of (<i>Z,4E/Z</i>)- 6c	S18
¹ H NMR of (<i>Z,4E/Z</i>)- 6d	S19
¹⁹ F NMR of (<i>Z,4E/Z</i>)- 6d	S20
¹ H NMR of (<i>Z,4E/Z</i>)- 6e	S21
¹⁹ F NMR of (<i>Z,4E/Z</i>)- 6e	S22
¹ H NMR of (<i>Z,4E</i>)- 6a	S23
¹³ C NMR of (<i>Z,4E</i>)- 6a	S24
¹⁹ F NMR of (<i>Z,4E</i>)- 6a	S25
¹ H NMR of (<i>Z,4E</i>)- 6b	S26
¹³ C NMR of (<i>Z,4E</i>)- 6b	S27
¹⁹ F NMR of (<i>Z,4E</i>)- 6b	S28
¹ H NMR of (<i>Z,4E</i>)- 6c	S29
¹³ C NMR of (<i>Z,4E</i>)- 6c	S30
¹⁹ F NMR of (<i>Z,4E</i>)- 6c	S31
¹ H NMR of (<i>Z,4E</i>)- 6d	S32
¹³ C NMR of (<i>Z,4E</i>)- 6d	S33
¹⁹ F NMR of (<i>Z,4E</i>)- 6d	S34













1231-CM-02-dimethoxy-sulfide-13C
 Pulse Sequence: s2pul
 Solvent: CDCl₃
 Temp. 25.0 C / 298.1 K
 Operator: barbara
 File: 1231-CM-02-dimethoxy-sulfide-13C
 INOVA-500 "riga"


Relax. delay 4.000 sec
 Pulse 52.1 degrees
 Acc. time 1.300 sec
 Width 29996.3 Hz
 44 repetitions
 OBSERVE C13, 125.6674328 MHz
 DECOUPLE H1, 499.7732084 MHz
 Power 42 dB
 on during acquisition
 WALTZ-16 modulated
 DATA PROCESSING
 Line broadening 0.0 Hz
 FT size 131072
 Total time 96 hr, 39 min, 38 sec









































Pulse Sequence : s2pul1
 Solvent: CDCl₃
 Ambient temperature
 File: 1231-cm-01-159-ER
 Mercury-300BBS "vega300"
 Relax. delay 4.000 sec
 Pulse 25.0 degrees
 Acq. time 0.300 sec
 Width 100.0 kHz
 188 Experiments
 OBSERVE P19, 282.3454132 MHz
 DATA PROCESSING
 Line broadening 1.0 Hz
 FT size 65536
 Total time 17 min, 33 sec

Crude mixture of
 (2Z,4E/Z)-**6d**
 282 MHz, CDCl₃

