

Review

Potential of Natural Products in the Inhibition of Adipogenesis through Regulation of PPAR γ Expression and/or Its Transcriptional Activity

Shi Feng, Laura Reuss and Yu Wang *

Citrus Research and Education Center, Food Science and Human Nutrition, University of Florida, 700 Experiment Station Road, Lake Alfred, FL 33850, USA; fsophias@ufl.edu (S.F.); lereuss@ufl.edu (L.R.)

* Correspondence: yu.wang@ufl.edu; Tel.: +1-863-956-8673

Academic Editor: Min-Hsiung Pan

Received: 10 August 2016; Accepted: 19 September 2016; Published: 23 September 2016

Abstract: Obesity is a global health problem characterized as an increase in the mass of adipose tissue. Adipogenesis is one of the key pathways that increases the mass of adipose tissue, by which preadipocytes mature into adipocytes through cell differentiation. Peroxisome proliferator-activated receptor γ (PPAR γ), the chief regulator of adipogenesis, has been acutely investigated as a molecular target for natural products in the development of anti-obesity treatments. In this review, the regulation of PPAR γ expression by natural products through inhibition of CCAAT/enhancer-binding protein β (C/EBP β) and the farnesoid X receptor (FXR), increased expression of GATA-2 and GATA-3 and activation of the Wnt/ β -catenin pathway were analyzed. Furthermore, the regulation of PPAR γ transcriptional activity associated with natural products through the antagonism of PPAR γ and activation of Sirtuin 1 (Sirt1) and AMP-activated protein kinase (AMPK) were discussed. Lastly, regulation of mitogen-activated protein kinase (MAPK) by natural products, which might regulate both PPAR γ expression and PPAR γ transcriptional activity, was summarized. Understanding the role natural products play, as well as the mechanisms behind their regulation of PPAR γ activity is critical for future research into their therapeutic potential for fighting obesity.

Keywords: obesity; adipogenesis; PPAR γ ; natural products; PPAR γ expression; PPAR γ transcriptional activity

1. Introduction

In recent decades, the prevalence of obesity has received serious attention worldwide. This chronic disease poses a severe threat to overall human health, as it increases the risk for a myriad of clinical conditions that include diabetes, hypertension, coronary atherosclerotic heart disease and certain cancers. Obesity can develop from genetic predisposition, an individual's metabolism, diet, lack of physical activity, as well as social status and lifestyle [1]. It is characterized by an increase in the mass of adipose tissue, specifically white adipose tissue. There are two major types of adipose tissues in mammals: white adipose tissue and brown adipose tissue. White adipose tissue plays a crucial role in lipid homeostasis and maintaining energy balance. It also stores energy in the form of triglycerides. The development of white adipose tissue is lifelong [2]. Unlike white adipose tissue, brown adipose tissue is especially abundant in newborns [3], but tends to decrease as humans age [4]. Brown adipose tissue is responsible for thermoregulation and heat production through non-shivering thermogenesis. An increase in the mass of adipose tissue can arise by increasing cell size (hypertrophy), cell number (hyperplasia or adipogenesis) or both. Adipogenesis is the process by which preadipocytes mature into adipocytes through cell differentiation. Adipocyte number does not necessarily promote obesity directly. Instead, the adipocyte number set during childhood and adolescence seems to play

a major role in determining the lipid-storing capacity of adipose tissue and fat mass in adults [5]. Therefore, regulation of adipogenesis may be one critical pathway for controlling or reversing obesity. Furthermore, identifying potential adipogenic molecular targets that can be modulated by external factors, such as food and drug agents from natural products, may largely contribute to the treatment of obesity. Peroxisome proliferator-activated receptor γ (PPAR γ), the chief regulator of adipogenesis, has been acutely investigated as a molecular target for natural products in the development of anti-obesity treatments. Understanding the role natural products play, as well as the mechanisms behind their inhibition of PPAR γ activity is critical for future research into their therapeutic potential for fighting obesity. This review summarizes the current knowledge of anti-adipogenesis mechanisms and the corresponding natural products that exhibit inhibitory effects of PPAR γ activity. Studies of adipogenesis in white adipose tissue will be the focus of this paper.

2. PPAR γ and Its Role in Adipogenesis

The peroxisome proliferator-activated receptors (PPARs) belonging to the thyroid/retinoid nuclear receptor family are transcription factors activated by lipophilic hormones, including steroids, thyroid hormones and vitamin A metabolites [6]. Primarily, PPARs control the expression of gene networks occurring in adipogenesis, lipid metabolism, inflammation and the maintenance of metabolic homeostasis [7]. They are also involved in the pathology of various diseases, such as obesity, type 2 diabetes, dyslipidemia and inflammatory conditions [8]. Among PPARs, the peroxisome proliferator-activated receptor gamma (PPAR γ), predominately expressed in adipose tissue, has been identified as a critical modulator that is not only sufficient, but also necessary for adipogenesis [9]. The importance of PPAR γ for adipocyte development and function was shown both in vitro and in vivo [10,11]. It was observed that ectopic expression of PPAR γ in NIH-3T3 fibroblasts caused a large percentage of these cells to undergo adipogenesis [12]. Furthermore, PPAR γ added to cultured myoblasts resulted in lipid accumulation and expression of adipocyte-specific markers [13]. In addition, mice chimeric for wild-type and PPAR γ null cells exhibited little or no contribution to adipose tissue by null cells [14]. It was also observed that conditional ablation of PPAR γ in mature adipocytes led to cell death and subsequent repopulation by PPAR γ -positive preadipocytes [15]. Moreover, humans with dominant-negative mutations introduced into the ligand-binding domain of PPAR γ displayed abnormal body fat distribution [16,17].

Activation of PPAR γ by certain agents has been demonstrated to stimulate adipogenesis in vitro and in vivo. It is believed that most pro-adipogenic factors function through activation of PPAR γ expression or at least PPAR γ activity to some extent [18]. PPAR γ exists in three major protein isoforms, PPAR γ 1, PPAR γ 2 and PPAR γ 3 [19,20]. While PPAR γ 1 is broadly expressed, PPAR γ 2 is almost adipocyte specific [21]. PPAR γ 3 is expressed in adipose tissue, as well as macrophages [20]. Additionally, PPAR γ 2 possesses an additional stretch of 30 amino acid residues in the ligand-independent domain of the N-terminal, which leads to higher transcriptional activity as compared to PPAR γ 1 and PPAR γ 3 [19,20,22]. PPAR γ is induced during adipocyte differentiation and before transcriptional activation of most adipocyte genes. It cooperates with other transcriptional factors, especially the CCAAT/enhancer binding proteins (C/EBPs) [23], to regulate adipocyte differentiation. C/EBP β and C/EBP δ induce the expression of PPAR γ and C/EBP α at the initial stage of adipogenesis, and subsequently, the concerted action of both PPAR γ and C/EBP α is required during the terminal differentiation of adipocytes. Moreover, it is proposed that PPAR γ is not only essential for adipogenesis, but also indispensable in the maintenance of the differentiated state [18]. Functioning as an essential regulator of adipogenesis, PPAR γ has been a critical target in the design and development efforts of numerous anti-obesity drugs. PPAR γ can be directly regulated by certain natural products whereas additional natural sources can also target and modulate upstream regulators of PPAR γ . This review investigates the regulation of PPAR γ expression by natural products through inhibition of CCAAT/enhancer-binding protein β (C/EBP β) and farnesoid X receptor (FXR), the increased expression of GATA-2 and GATA-3 and the activation of the Wnt/ β -catenin pathway. Furthermore,

the regulations of PPAR γ transcriptional activity associated with natural products through antagonism of PPAR γ and activation of Sirtuin 1 (Sirt1) and AMP-activated protein kinase (AMPK) were discussed. Lastly, the regulation of mitogen-activated protein kinase (MAPK) by natural products, which might regulate both PPAR γ expression and PPAR γ transcriptional activity, was summarized.

3. Natural Product Regulation of PPAR γ

The regulation of PPAR γ can be carried out directly or/and indirectly via the anti-adipogenic activities of various natural products. The expression and activity of PPAR γ make two straightforward targets for regulation. Thus far, a range of natural products has been reported to suppress adipogenesis via downregulation of PPAR γ expression and/or activity (Table 1). Some of these natural products have been closely investigated, and certain active compounds (Figure 1) have been, at least, partly associated with the regulation of PPAR γ upstream modulators or PPAR γ itself. The possible pathways regulating PPAR γ reviewed in this paper are listed in Figure 2. Nevertheless, for many natural products, the mechanisms regulating PPAR γ remain unclear.

Table 1. Natural products that regulate PPAR γ .

Mechanism of Action	Source (Possible Natural Sources)	Active Component
Inhibition of C/EBP β	Commercial standard (legumes) [24] Not specified (animal liver) [25] <i>Rehmannia glutinosa</i> [26]	Genistein Retinoic acid (RA) Not specified (alcohol extract)
Inhibition of FXR	Resin of the guggul tree [27]	Guggulsterone
Increased expression of GATA-2 and GATA-3	<i>Cortidis rhizome</i> [28]	Berberine
Activation of Wnt/ β -catenin pathway	Commercial standard (legumes) [29] Not specified (rhizomes of <i>Curcuma longa</i>) [30] Commercial standard (<i>Lithospermum erythrorhizon</i> Sieb. Et Zucc) [31]	Genistein Curcumin Shikonin
Antagonism of PPAR γ	<i>Rhaponticum uniflorum</i> [32] Commercial standard (black pepper) [33]	7-Chloroarctonone-b Piperine
Activation of Sirtuin 1	Not specified (Japanese knotweed, peanut) [34] Not specified (broccoli, cabbage) [35] Commercial standard (pomegranate seed oil and brown seaweed extract) [36]	Resveratrol Indole-3-carbinol (I3C) Xanthigen
Activation of AMPK	Commercial standard (<i>Curcuma longa</i>) [37] <i>Lysimachia foenum-graecum</i> [38] Not specified (soybean) [39] Commercial standard (<i>Panax ginseng</i>) [40] Commercial standard (<i>Zanthoxylum armatum</i>) [41] Commercial standard (<i>Mirabilis jalapa</i> , apple) [42] Commercial standard (pomegranate seed oil and brown seaweed extract) [36]	Curcumin Not specified (ethanol extract) Genistein Ginsenoside Rh2 Methyl cinnamate Ursolic acid Xanthigen
Regulation of MAPK	Commercial standard (<i>Evodia fructus</i>) [43] Green tea [44] <i>Hibiscus sabdariffa</i> L. [45]	Evodiamine EGCG Not specified (water extract)
Unknown	<i>Lagerstroemia speciosa</i> L. [46] <i>Hibiscus sabdariffa</i> [45] <i>Undaria pinnatifida</i> [47] Commercial standard (beer hops) [48] Commercial standard (red pepper) [49] <i>Eriobotrya japonica</i> leaves [50] Commercial standard (vinegar, buckwheat) [51] Commercial standard (grape and onion) [52] Brown algae [53] <i>Monascus</i> [54] Commercial standard (onion) [55] Dry <i>Undaria pinnatifida</i> powder [47] <i>Lithospermum erythrorhizon</i> [56] Ginseng [57] <i>Caesalpinia sappan</i> L. [58] <i>C. maculata</i> and <i>C. subternata</i> [59] <i>Adenophora triphylla</i> var. <i>japonica</i> extract [60] <i>Allomyrina dichotoma larvae</i> [61] Chickpea [62] Not specified (<i>Garcinia indica</i> , almonds) [63] Rhizome of <i>Coptis chinensis</i> [64]	Not specified (hot water extract) Not specified (dried flower extract) Fucoxanthin and fucoxanthinol Xanthohumol and isoxanthohumol Capsaicin Corosolic Acid o-Courmaric acid and rutin Resveratrol and quercetin Fucoidan Monascin and ankaflavin Quercetin Amarouciaxanthin A Shikonin Ginsenoside Rh1 Brazilein Not specified (hot water extract) Lupenone Not specified (ethanol extract) Isoflavones Garcinol, pterostilbene Berberine, epiberberine, coptisine, palmatine, and magnoflorine




Figure 1. Chemical structures of active compounds regulating PPAR γ derived from natural products.




Figure 2. Possible pathways by which natural products regulate PPAR γ .

3.1. Regulation of PPAR γ Expression

The regulation of PPAR γ expression by natural products through different possible pathways is demonstrated in Figure 3.




Figure 3. Possible pathways by which natural products regulate PPAR γ expression.

3.1.1. Inhibition of CCAAT/Enhancer-Binding Protein β

C/EBPs are a family of transcription factors controlling the differentiation of a variety of cell types. C/EBP α and C/EBP β are not only the most widely-expressed, but also the most well-studied isoforms found in the C/EBP family [65]. C/EBP α and C/EBP β are well known for their adipogenic transcriptional activities that promote adipogenesis. Conversely, some members of the C/EBP family can repress adipocyte differentiation by forming inactive heterodimers with C/EBP α and C/EBP β [24]. C/EBP homologous protein (CHOP) interacts with the C/EBP transcription factors

to form heterodimers that cannot bind DNA, but instead function as a dominant-negative inhibitor of gene transcription [66,67]. Therefore, increasing the CHOP level may possibly inhibit C/EBP β activity. As previously mentioned, C/EBP β is key for inducing the initial expression of PPAR γ during adipogenesis. Consequently, the direct inhibition of C/EBP β activity can subsequently lead to PPAR γ suppression, thus inhibiting adipogenesis.

Genistein, an isoflavone primarily found in legumes, was shown to have anti-adipogenic effects in 3T3-L1 cells by blocking the DNA binding and transcriptional activity of C/EBP β [24]. This, in turn, inhibited the protein expression of differentiation-induced PPAR γ and C/EBP α . The proposed mechanisms for this were a deactivation of C/EBP β through increased levels of CHOP, as well as inhibition of the tyrosine phosphorylation of C/EBP β . Besides anti-adipogenic effects, the pharmacological activities of genistein have been demonstrated in various published investigations and include tyrosine kinase inhibition, chemoprotective activities against cancers and cardiovascular disease and phytoestrogen activities [68]. As the simplest biosynthetic isoflavonoid compound in legumes, genistein (4',5,7-trihydroxyisoflavone) plays the role of the central intermediary in the biosynthesis of more complex isoflavonoids. Among isoflavonoids, isoflavones including genistein are a group of compounds considered as important as the phytoestrogens that play a beneficial role in fighting obesity. Several studies of obese humans and animals indicate that phytoestrogens have significant anti-obesity effects [69–71]. Dietary sources of phytoestrogens include legumes, seeds and whole grains, all of which may contribute greatly to anti-obesity drug development. In addition to genistein, some other natural products have been shown to inhibit C/EBP β activity, as well. However, the mechanism for this suppression has not been elucidated. For instance, retinoic acid (RA) was illustrated to inhibit adipogenesis by blocking C/EBP β -induced expression of the downstream gene, PPAR γ [25]. RA is considered the most active form of vitamin A, existing in the body in various essential life processes [72]. Sources rich in vitamin A include animal liver, red capsicum, sweet potato, carrots and broccoli. More recently, the extract of *Rehmannia glutinosa*, a traditional Chinese medicine, was demonstrated to inhibit adipocyte differentiation through the inhibition of C/EBP β expression [26].

3.1.2. Inhibition of the Farnesoid X Receptor

The farnesoid X receptor, important in both bile acid and cholesterol homeostasis, was first identified in 1999 as a nuclear receptor for bile acids [73]. FXR expression has been reported in liver, intestine, kidney and the adrenal glands [74]. Recently, FXR expression was reported in adipose tissue, thus the role of FXR regulation of adipogenesis in 3T3-L1 cells was investigated [75]. FXR was demonstrated to promote adipocyte differentiation by at least partially inducing PPAR γ 2 and C/EBP α gene expressions in in vivo experiments. It regulated adipogenesis by both PPAR γ -dependent and independent mechanisms. The process of FXR activation is similar to other nuclear hormone receptors. An agonist ligand is required for FXR activation after heterodimerization of the retinoid X receptor (RXR). Then, FXR regulates target gene expression by binding to FXR response elements. Therefore, investigating antagonist ligands for FXR is one approach for fighting obesity through inhibition of FXR activity and, thereby, suppressing PPAR γ 2 expression.

An early study of the plant sterol guggulsterone, extracted from the resin of the guggul tree, identified a highly efficacious antagonist of FXR [27]. It was demonstrated that guggulsterone exhibited cholesterol-lowering activity based on its inhibition of FXR activation [27]. More recently, guggulsterone was used to illustrate that FXR antagonism prevented preadipocyte differentiation [75]. Later, treatment with *cis*-guggulsterone during the maturation period of adipocytes was illustrated to downregulate the adipocyte-specific transcription factors PPAR γ 2, C/EBP α and C/EBP β in 3T3-L1 cells [76]. In addition to studies of the individual use of guggulsterone, the combination of guggulsterone and genistein was shown to have enhanced anti-adipogenic effects in 3T3-L1 adipocytes compared to the effects of individual compounds alone [77]. Similarly, it was also demonstrated that the combination of guggulsterone and xanthohumol more potently exerted anti-obesity effects than

additive effects of the individual compounds [78]. This suggests that reasonable combinations of natural products exhibiting anti-obesity effects are a possible strategy for the development of effective obesity treatments.

3.1.3. Increase Expression of GATA-2 and GATA-3

The GATA family of transcription factors binds specifically to the consensus DNA sequence (A/T)GATA(A/G) and share highly conservative zinc finger DNA binding domains [79]. They play important roles in a variety of developmental processes, including adipogenesis. Among the GATA transcription factors, GATA-2 and GATA-3 are predominantly present in white adipose tissue, specifically the preadipocytes. GATA-2 and GATA-3 are considered potential preadipocyte markers and significantly contribute to the regulation of adipocyte differentiation. It has been demonstrated that the constitutive expression of GATA-2 and GATA-3 inhibits adipocyte differentiation, trapping cells in the preadipocyte stage. This can be attributed to GATAs' ability to decrease PPAR γ 2 expression through direct suppression of the PPAR γ 2 promoter [79], as well as by interacting with the C/EBP family of transcription factors [80]. Both GATA-2 and GATA-3 can form protein complexes with either C/EBP α or C/EBP β , interfering with their adipocytic functions. These findings indicate that GATA-2 and GATA-3 play a crucial role as molecular gatekeepers during the onset of terminal adipocyte differentiation and, therefore, may serve as targets for the therapeutic intervention of obesity.

Berberine is an isoquinoline derivative alkaloid isolated from many traditional Chinese medicinal herbs. Purified from *Coptidis rhizome*, berberine was shown early on to inhibit PPAR γ by affecting the mRNA and protein levels of PPAR γ , as well as PPAR γ transcriptional activity in 3T3-L1 cells [81]. It was suggested that the inhibition of PPAR γ by berberine is, at least partly, dependent on the C/EBP β signal, an upstream modulator of PPAR γ . The inhibition of PPAR γ transcription has been demonstrated to be the result of reducing PPAR γ protein levels while eliminating the possibility of berberine as the antagonist of PPAR γ . More recently, a new pathway related to the inhibitory effects of berberine on adipocyte differentiation was discussed [28]. In addition to decreasing the expression of PPAR γ , berberine showed the capability to increase GATA-2 and GATA-3 mRNA and protein expression along with the inhibition of adipocyte differentiation in 3T3-L1 cells. In this study, berberine was shown to influence GATA-2 and GATA-3 expression during differentiation and induced increased gene expression of these genes in a dose-dependent manner. Similar to GATA mRNA expression, a significant increase of GATA protein expression was observed in cells grown in the presence of berberine. However, the potential mechanism by which berberine may affect the expression of GATA-2 and GATA-3 was not investigated here, and further elucidation behind this observation is needed. The inhibitory effects of berberine on adipogenesis in high-fat diet-induced obese mice were also reported by the same group of researchers. Berberine was shown to reduce weight gain and food intake in high-fat diet-induced obese mice, while no effect was observed in mice fed a normal diet [82]. Both increased mRNA expression of GATA-2 and GATA-3 was demonstrated, as was the decreased expression of PPAR γ and C/EBP α mRNA in the epididymal fat of high-fat diet-induced obese mice. Meanwhile, berberine plus another botanical alkaloid, evodiamine, were investigated for their inhibitory effect and transcriptional impact individually, as well as in combination on human white preadipocyte differentiation [83]. It was shown that berberine and evodiamine increased both the mRNA expression and the protein expression of GATA-2 and GATA-3. Berberine did not affect cell viability, while evodiamine substantially decreased cell viability. However, when used together, the viability inhibition effect of evodiamine was reversed by berberine. Based on current knowledge, berberine has been shown to exhibit anti-obesity potential both in vitro and in vivo.

3.1.4. Activation of the Wnt/ β -Catenin Pathway

Wnts are a family of secreted signaling proteins that regulate cell-to-cell interactions during development and adult tissue homeostasis. Wnt signaling, likely mediated by Wnt10b, has been showed to govern adipogenesis and maintain preadipocytes in an undifferentiated state by inhibiting

adipogenic transcription factors PPAR γ (Figure 4) and C/EBP α [84]. This may be associated with the Wnt target genes, c-myc, cyclin D1 and PPAR δ , that have been reported to inhibit the expression and activity of both PPAR γ and C/EBP α [85–87]. Wnt signaling is initiated by the binding of Wnts to Frizzled (Fz) receptors, followed by heterodimerization between the Fz receptors and low-density lipoprotein receptor-related protein (LRP) co-receptors [88]. In the β -catenin-dependent pathway, Wnt signaling activates Dishevelled, inhibits the activity of the destruction complex and thereby stabilizes β -catenin through phosphorylation [89]. The destruction complex, containing GSK-3, Axin, APC and β -TrCP/Slimb, promotes rapid degradation of β -catenin in the absence of Wnt signals. Once translocated to the nucleus, β -catenin binds to the T-cell factor/lymphoid-enhancing factor (TCF/LEF) family of transcription factors and then regulates the expression of Wnt target genes [89,90].




Figure 4. Possible mechanism of PPAR γ regulation by genistein, curcumin and shikonin via activation of the Wnt/ β -catenin pathway.

Genistein has been demonstrated to regulate adipogenesis through different pathways. In a recent study, repression of the adipogenic differentiation of human adipose tissue-derived mesenchymal stem cells by genistein was associated with Wnt/ β -catenin signaling [29]. In this study, genistein was shown to promote Wnt signaling by interacting with Wnt ligands, Wnt antagonists and Wnt intermediates. The expression of Wnt3, which inhibits adipogenesis via regulation of PPAR γ and C/EBP α expression in 3T3-L1 cells [88], was increased by genistein. Meanwhile, Wnt signaling antagonists, including sFRP1, DKK2, CK1 and Axin2, were inhibited by genistein. Genistein also augmented mRNA and protein levels of β -catenin. Curcumin, a polyphenol found in the rhizomes of *Curcuma longa*, was also observed to have anti-adipogenic effects associated with the activation of Wnt/ β -catenin signaling in 3T3-L1 cells [30]. During differentiation, curcumin restored nuclear translocation of β -catenin and reduced differentiation-stimulated expression of the components in the destruction complex, including CK1 α , GSK-3 β and Axin. Meanwhile, curcumin was observed to increase mRNA levels of c-Myc and cyclin D1, the Wnt target genes. More recently, shikonin isolated from *Lithospermum erythrorhizon* Sieb. Et Zucc was shown to inhibit adipogenesis through modulation

of the Wnt/ β -catenin pathway [31]. In this study, shikonin maintained the nuclear level of β -catenin while increasing the level of its transcriptional product, cyclin D1 during adipogenesis of 3T3-L1 cells. Meanwhile, shikonin-induced reductions of PPAR γ and C/EBP α were significantly recovered by siRNA-mediated knockdown of β -catenin. Similarly, isoquercitrin and isorhamnetin, active flavonoids in the extract of *Persicaria hydropiper* (L.) spach (an herbal plant used to add spicy flavor to traditional Chinese dishes), were demonstrated to inhibit the adipocyte differentiation of 3T3-L1 cells via the activation of Wnt/ β -catenin signaling [91].

3.2. Regulation of PPAR γ Transcriptional Activity

3.2.1. Antagonist of PPAR γ

PPAR γ functions as a heterodimer with the retinoid X receptor (RXR) and binds to specific DNA sequences to regulate the transcription of target genes [7]. The activity of PPAR γ is regulated by binding agonist ligands, including steroid and thyroid hormones, vitamins, lipid metabolites and xenobiotics [92]. Unliganded (agonist ligand) PPAR γ /RXR heterodimers repress the transcription of target genes by interacting with corepressor molecules [93], while PPAR γ binding with the agonist ligand experiences conformational changes facilitating the dissociation of corepressor molecules to enable recruitment of coactivators to the liganded receptor. Therefore, investigation of PPAR γ antagonists that may directly inhibit PPAR γ activity by interrupting its functional pathway can be considered a logical approach for research in the fight against obesity.

In recent years, researchers have been attempting to screen natural products for novel PPAR γ antagonists with inhibitory effects on adipogenesis. 7-chloroarctinone-b (CAB), a new thiophene-acetylene type of derivative isolated from *Rhaponticum uniflorum*'s roots, was identified as a specific PPAR γ antagonist [32]. The inhibitory effects of CAB on PPAR γ /RXR α heterodimerization and PPAR γ co-activation recruitment were studied. CAB showed high binding affinity for the PPAR γ -ligand binding domain (LBD), thereby considerably antagonizing the PPAR γ agonist rosiglitazone-stimulated PPAR γ -LBD/RXR α -LBD dimerization. Meanwhile, it also considerably antagonized rosiglitazone-stimulated PPAR γ coactivator recruitment. In the 3T3-L1 cell differentiation assay, it was demonstrated that CAB effectively antagonized both hormone and rosiglitazone-induced adipocyte differentiation in cell culture. Thiophene-acetylene represents a unique class of natural products presenting a wide variety of biological activities, including antitumor, antiviral, anti-HIV and antifungal to insecticidal activities [94]. The discovery of PPAR γ 's antagonistic activity of CAB may imply the potential of thiophene-acetylene as an anti-obesity treatment. Similarly, piperine, a major alkaloid-amine component of black pepper, was shown to inhibit adipogenesis by antagonizing PPAR γ activity in 3T3-L1 cells [33]. Piperine significantly repressed the rosiglitazone-induced PPAR γ transcriptional activity while also markedly decreasing mRNA expression of PPAR γ . Furthermore, piperine was demonstrated to disrupt the rosiglitazone-dependent interaction between PPAR γ and coactivator CREB-binding protein (CBP). Before being discovered as a potential antagonist of PPAR γ , piperine had previously been known for its painkilling, antioxidant, antitumor and anti-inflammatory activities [33].

3.2.2. Activation of Sirtuin 1

Sirt1 is an NAD⁺-dependent nuclear deacetylase in mammals. It has been identified as a key mediator of pathways located downstream of calorie restriction and considered a therapeutic target in the treatment of age-related diseases [95]. Sirt1 is involved in processes that include chromatin remodeling, transcriptional silencing, chromosomal stability, cell cycle progression, apoptosis, autophagy, metabolism, growth suppression, inflammation and stress responses [96]. Recently, Sirt1 has emerged as a novel therapeutic target in the fight against obesity. It has been demonstrated that Sirt1 is a negative modulator of adipogenesis in the 3T3-L1 model [97]. In 3T3-L1 adipocytes, overexpression of Sirt1 was shown to attenuate adipogenesis, while the RNA interference of Sirt1

enhanced it. Meanwhile, upregulation of Sirt1 led to lipolysis and the loss of fat in differentiated adipocytes. It is believed that Sirt1 promotes fat mobilization in white adipocytes through PPAR γ repression, interacting with its cofactors nuclear receptor co-repressor (NCoR) and the silencing mediator of retinoid and thyroid hormone receptors (SMRT). Sirt1 and PPAR γ have been shown to bind to the same DNA sequences, suggesting that Sirt1 plays a role as a co-repressor of PPAR γ . This is consistent with a previous study where, in the absence of ligand, PPAR γ recruited NCoR and SMRT, and these co-repressors were capable of downregulating PPAR γ -mediated transcriptional activity [98]. 3T3-L1 cells deficient in NCoR or SMRT were exposed to differentiation media; they then exhibited increased expression of adipocyte-specific genes and increased production of lipid droplets, as compared to the control cells. Therefore, activation of Sirt1 during adipogenesis could be beneficial in the therapeutic intervention of obesity. Besides regulation of PPAR γ , activated Sirt1 further deacetylates the transcriptional coactivator peroxisome proliferator-activated receptor gamma coactivator 1-alpha (PGC-1) at promoter regions to induce the expression of genes involved in mitochondrial biogenesis and fatty acid oxidation [99].

Resveratrol is a member of the natural polyphenols found in plants. It is abundant in Japanese knotweed and found in considerable amounts in peanuts, groundnuts, grapevines and red vines [100]. Resveratrol has gained much attention due to its numerous, beneficial cardio-protective effects [101] and anti-cancer [102], anti-inflammatory and antioxidant actions [103]. Recently, resveratrol has been shown to play a role in multiple activities in adipose tissue, including adipocyte proliferation, adipogenesis, lipolysis and apoptosis [104]. It is the first polyphenolic compound that has been shown to activate Sirt1 [105,106]. Moreover, resveratrol was identified as the most potent polyphenolic compound activator of Sirt1 [34,95]. It has been shown to inhibit proliferation and adipogenic differentiation in a Sirt1-dependent manner in human preadipocytes [107]. Human preadipocytes were used to demonstrate the inhibitory effect of resveratrol on adipogenic differentiation. The fact that the inhibitory activity of resveratrol was abolished by the knockdown of Sirt1 indicated the role of Sirt1 as a mediator of the resveratrol effect. Similarly, in an in vitro study, investigation of the inhibitory effects of resveratrol on adipogenesis was conducted in pig primary preadipocytes [108]. The downregulation of adipogenesis was also associated with an increased expression of Sirt1 mRNA, which, in turn, suppressed PPAR γ . Thus far, the mechanisms that link resveratrol to the activation of Sirt1 are still debated, and recent studies have shown that resveratrol does not directly activate Sirt1 [109,110].

Along with resveratrol, some other polyphenols, such as butein, piceatannol and quercetin, were shown to enhance Sirt1 activity in vitro through the demonstration of the effects on the stimulation of the Sirt1 catalytic rate [34]. There has been a very limited number of studies assessing the ability of other polyphenols, including quercetin and catechins, to induce Sirt1 activity [111,112]. Some conclusions are contradictory, as both the inhibitory effects and stimulation effects have been observed, which may be explained as the powerful effects of polyphenol stability and metabolism on the stimulation of Sirt1. Xanthigen, a source of punicic acid and fucoxanthin derived from pomegranate seed and brown seaweed, was demonstrated to deregulate Sirt1 and activate AMP-activated protein kinase (AMPK) signaling accompanied with downregulation of PPAR γ [36]. More recently, It was reported that indole-3-carbinol (I3C), an efficacious, specific Sirt1 activator in cultured 3T3-L1 cell lines, ameliorated adipogenesis [35]. I3C is the product of the breakdown of glucosinolate glucobrassicin, which is abundant in many vegetables, including broccoli, cabbage, cauliflower and kale. I3C did not inhibit adipocyte differentiation in 3T3-L1 cells where Sirt1 was knocked down. Furthermore, reverse transcriptase polymerase chain reaction analysis showed that I3C treatment reduced mRNA levels of adipogenic genes encoding for C/EBP α , PPAR γ 2, FAS and aP2 in 3T3-L1 cells, but not in Sirt1 knockdown cells. Understanding the role and mechanisms of natural products in Sirt1 regulation is crucial for the discovery and development of pharmacological agents for potential use in the clinical management of obesity.

3.2.3. Activation of AMP-Activated Protein Kinase

AMPK is a serine/threonine protein kinase that works as an essential regulator of cellular metabolism and energy balance. It is a heterotrimeric protein consisting of three subunits, α , β and γ , among which the catalytic subunit α is vital for AMPK activation through its own activation via phosphorylation [113]. AMPK promotes mechanisms that increase energy production and shut down pathways that consume energy. The anti-adipogenic effects of AMPK are expected since adipocyte differentiation can be seen as an energy-consuming process involving new membrane and protein synthesis, two processes that are heavily prohibited by AMPK activation [114,115]. Long-term activation of AMPK can affect the pattern of gene expression in a variety of circumstances. To be more specific, AMPK can directly phosphorylate and regulate proteins involved in gene transcription, such as transcription factors, cofactors and components of the transcriptional core machinery [116]. Activation of AMPK has been shown to attenuate the expression of PPAR γ in 3T3-L1 adipocytes [117]. AMPK was observed as an upstream positive regulator of p38 MAPK [118], which was demonstrated to promote PPAR γ phosphorylation, inhibit its transcriptional activity and thereby block adipocyte differentiation [119,120]. The modulation of AMPK has been viewed as a potential obesity therapy. Berberine was demonstrated to increase the activation (phosphorylation) of AMPK and p38, as well as the deactivation of PPAR γ in adipocytes [121], and they were proposed as the possible pathways by which berberine mediated its metabolic actions. Curcumin was confirmed to regulate adipocyte differentiation by activating AMPK and downregulating PPAR γ transcription [37]. It was observed that AMPK activation led to the inhibition of PPAR γ expression in 3T3-L1 adipocytes, although the second regulator between AMPK and PPAR γ was not elucidated. More recently, the anti-obesity effects of *Lysimachia foenum-graecum* (LFE) were associated with its ability to block adipogenesis in 3T3-L1 adipocytes through the possible link between LFE-induced activation of AMPK and AMPK-mediated suppression of PPAR γ transcriptional activity [38].

Accumulating evidence has shown the important effects of AMPK activation by natural products in the inhibition of adipogenesis. Genistein exerted the inhibition of adipocyte differentiation through the activation of AMPK paralleled with the generation of reactive oxygen species (ROS), an upstream signal for AMPK that rapidly activates AMPK [39]. Anti-obesity effects of ginsenoside Rh2, an active component derived from *Panax ginseng*, are associated with the activation of the AMPK signaling pathway in 3T3-L1 adipocytes, while ROS plays a role in AMPK activation during ginsenoside Rh2 treatment [40]. It was also demonstrated that ursolic acid (UA), a triterpenoid compound identified in many fruits and plants, increased phosphorylation and activation of AMPK by stimulating LKB1 (an upstream kinase of AMPK) activity and consequently inhibited adipogenesis [42].

3.3. Comprehensive Regulation of PPAR γ

Regulation of Mitogen-Activated Protein Kinase

Mitogen-activated protein kinases (MAPKs) are well known for their three subfamilies, ERK, JNK/SAPK and p38. The activation of MAPK occurs through phosphorylation of specific threonine and tyrosine residues on the MAPK molecule by an upstream kinase, MAPK kinase [122]. MAPK cascades are interactive intracellular signal transduction pathways that often regulate cellular control switches [123,124]. These cascades consist of a three-kinase module, MAPK, MEK (MAPK/ERK kinase, MAPK activator) and MEKK (MEK kinase, MEK activator) [122]. In an early study, activation of MAPK was shown to antagonize 3T3-L1 adipocytic differentiation [125]. It is believed that MAPK-mediated phosphorylation of PPAR γ contributes to the reduction of PPAR γ transcriptional activity and, thereby, inhibits adipocyte differentiation [119]. PPAR γ activity is finely modulated; in addition to ligand-mediated activation, its phosphorylation status and the molecular consequences include changes in ligand affinity, DNA binding, recruitment of transcriptional cofactors, proteasome degradation, etc. [120]. By using preadipocyte cultures, evodiamine, a major alkaloidal compound in

the fruit of *Evodia fructus*, was found to increase activation of ERK/MAPK, to reduce the expression of PPAR γ and, thereby, significantly inhibit adipocyte differentiation [43].

Contradictory reports also exist. It was proposed that deactivation of ERK was necessary for inhibition of adipocyte differentiation [126,127]. Green tea polyphenol EGCG was shown to inhibit adipogenesis in 3T3-L1 adipocytes [44]. It was also proposed that the possible mechanisms behind the anti-adipogenic effects of EGCG involve mitogen-activated protein (MAP) kinase, especially the extracellular signal-regulated kinases (ERKs) that are stimulated by growth-related signals. EGCG has been reported to reduce phosphor-ERK1 and phosphor-ERK2 levels in 3T3-L1 preadipocytes [128]. Additionally, ERK1/2 phosphorylation was demonstrated to promote C/EBP α and PPAR γ expression in 3T3-L1 cells [127]. Here, inhibition of MAPK seems to repress adipogenesis. A similar mechanism was also observed in other natural products. The water extract of *Hibiscus sabdariffa* L., a medicinal herb, was investigated for its inhibition of adipocyte differentiation in 3T3-L1 preadipocytes [45]. Hibiscus extract blocked the MAPK pathway by inhibiting the phosphorylation of ERK and, consequently, the expression of C/EBP α and PPAR γ . Further investigation into the influences of the MAPK pathway of adipocyte differentiation is definitely needed, and clear elucidation of this mechanism could provide a potential foundation for future developments in anti-obesity drugs.

4. Possible Future Trends and Applications of Natural Products on Anti-Obesity

Obesity is recognized as a worldwide health crisis. It is not just a severe health concern, but also a risk factor for many chronic diseases, including diabetes, hypertension, cardiovascular diseases, as well as certain cancers. The only obesity-treatment drug currently available, Orlistat, has serious side effects that include increased blood pressure, dry mouth, constipation, headache and insomnia [129]. Sibutramine, a previous prescription drug for weight loss, was withdrawn by the U.S. Food and Drug Administration in 2010 due to an increased risk of cardiovascular adverse events. Therefore, in recent years, natural alternatives that exhibit anti-obesity potential have been widely investigated. As discussed above, natural products, which have shown large potential to regulate PPAR γ expression and/or transcriptional activity, are potential therapies to suppress adipogenesis and, thereby, fight against obesity.

Currently, the majority of natural products investigated are phytochemicals obtained from dietary and medicinal plants, both of which are relatively safe sources and easy to access. Primarily obtained from vegetables and fruits, dietary phytochemicals could be employed as anti-obesity agents due to evidence indicating that they may reduce adipose tissue mass through inhibiting adipogenesis, stimulating lipolysis and inducing apoptosis of existing adipocytes. Among dietary phytochemicals, polyphenols, terpenoids, phytosterols, as well as alkaloids are common groups with active components showing anti-obesity effects [130,131]. The application of medicinal plants in treating metabolic diseases has a long history, especially in Asia. Plants have always been a source of medication, although initially, their active molecules may not be clearly identified. With an increasing understanding of obesity and its key regulator, PPAR γ , investigations into the functional mechanism of PPAR γ along with the effective regulators of PPAR γ have been largely conducted. The identification of both the possible targets that regulate PPAR γ , as well as the active components that modulate PPAR γ are two key steps towards fighting obesity via the inhibition of adipogenesis, and the continuation of these elucidation efforts is imperative.

The current understanding indicates that specific natural products can target more than one known pathway that regulates PPAR γ expression and/or transcriptional activity. For example, genistein has been shown to suppress PPAR γ expression by both the inhibition of C/EBP β , as well as the activation of the Wnt/ β -catenin pathway, whereas it also controls PPAR γ transcriptional activity through activation of AMPK during adipogenesis. This suggests that a natural product such as genistein could be a highly effective anti-obesity agent. However, further investigations into the safety, metabolism and human use of genistein as a medication are necessary. Combined treatment is an additional option for obesity therapy. For instance, the combined effect of both berberine and evodiamine on human white

preadipocyte differentiation has been studied with no observable significant additive or synergetic effects of adipogenesis inhibition. Nevertheless, there is no doubt that the appropriate combination of effective natural products would help the design of novel strategies in preventing the observed current epidemic levels of obesity.

It is not surprising that all known pathways for PPAR γ regulation are not limited to this review. However, the study of these numerous pathways remains in the initial stages, and the application of natural products on these pathways has yet to be investigated. For example, the deficiency of cellular retinol-binding protein type 1 (CRBP1), expressed in preadipocytes prior to the expression of PPAR γ , is tightly associated with enhanced PPAR γ expression and activity [132]. Overexpression of Sirt2 inhibits differentiation, accompanied by decreased expression of PPAR γ [133]. It is proposed that reducing the level of Sirt2 results in an increased level of forkhead box protein O1 (FOXO1) acetylation/phosphorylation, thereby reducing the ability of FOXO1 to interact with the PPAR γ promoter and repress PPAR γ transcription. The high-affinity and PPAR γ -specific antagonist, cyclic phosphatidic acid (cPA), stabilizes the PPAR γ -SMRT corepressor complex and inhibits PPAR γ -mediated gene transcription [134]. It has also been observed that the signal transducer and activator of transcription 5 (STAT5) stimulates the expression of PPAR γ [135]. While the understanding and discovery of new pathways for PPAR γ regulation are proceeding, additional therapeutic targets for obesity treatment are presented, and the further application of possible active natural products could be conducted.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Ordovas, J.M.; Shen, J. Gene-environment interactions and susceptibility to metabolic syndrome and other chronic diseases. *J. Periodontol.* **2008**, *79*, 1508–1513. [[CrossRef](#)] [[PubMed](#)]
2. Flier, J.S. The adipocyte: Storage depot or node on the energy information superhighway? *Cell* **1995**, *80*, 15–18. [[CrossRef](#)]
3. Gesta, S.; Tseng, Y.-H.; Kahn, C.R. Developmental origin of fat: Tracking obesity to its source. *Cell* **2007**, *131*, 242–256. [[CrossRef](#)] [[PubMed](#)]
4. Graja, A.; Schulz, T.J. Mechanisms of aging-related impairment of brown adipocyte development and function. *Gerontology* **2014**, *61*, 211–217. [[CrossRef](#)] [[PubMed](#)]
5. Spalding, K.L.; Arner, E.; Westermark, P.O.; Bernard, S.; Buchholz, B.A.; Bergmann, O.; Blomqvist, L.; Hoffstedt, J.; Näslund, E.; Britton, T.; et al. Dynamics of fat cell turnover in humans. *Nature* **2008**, *453*, 783–787. [[CrossRef](#)] [[PubMed](#)]
6. Mangelsdorf, D.J.; Thummel, C.; Beato, M.; Herrlich, P.; Schütz, G.; Umesono, K.; Blumberg, B.; Kastner, P.; Mark, M.; Chambon, P. The nuclear receptor superfamily: The second decade. *Cell* **1995**, *83*, 835–839. [[CrossRef](#)]
7. Wang, L.; Waltenberger, B.; Pferschy-Wenzig, E.-M.; Blunder, M.; Liu, X.; Malainer, C.; Blazevic, T.; Schwaiger, S.; Rollinger, J.M.; Heiss, E.H.; et al. Natural product agonists of peroxisome proliferator-activated receptor gamma (PPAR γ): A review. *Biochem. Pharmacol.* **2014**, *92*, 73–89. [[CrossRef](#)] [[PubMed](#)]
8. Petersen, R.K.; Christensen, K.B.; Assimopoulou, A.N.; Fretté, X.; Papageorgiou, V.P.; Kristiansen, K.; Kouskoumvekaki, I. Pharmacophore-driven identification of PPAR γ agonists from natural sources. *J. Comput. Aided Mol. Des.* **2011**, *25*, 107–116. [[CrossRef](#)] [[PubMed](#)]
9. Tontonoz, P.; Spiegelman, B.M. Fat and beyond: The diverse biology of PPAR γ . *Annu. Rev. Biochem.* **2008**, *77*, 289–312. [[CrossRef](#)] [[PubMed](#)]
10. Tang, Q.-Q.; Otto, T.C.; Lane, M.D. Mitotic clonal expansion: A synchronous process required for adipogenesis. *Proc. Natl. Acad. Sci. USA* **2003**, *100*, 44–49. [[CrossRef](#)] [[PubMed](#)]
11. Gray, S.; Dalla Nora, E.; Vidal-Puig, A. Mouse models of PPAR- γ deficiency: Dissecting PPAR- γ 's role in metabolic homeostasis. *Biochem. Soc. Trans.* **2005**, *33*, 1053–1058. [[CrossRef](#)] [[PubMed](#)]
12. Tontonoz, P.; Hu, E.; Spiegelman, B.M. Stimulation of adipogenesis in fibroblasts by PPAR γ 2, a lipid-activated transcription factor. *Cell* **1994**, *79*, 1147–1156. [[CrossRef](#)]
13. Hu, E.; Tontonoz, P.; Spiegelman, B.M. Transdifferentiation of myoblasts by the adipogenic transcription factors PPAR gamma and C/EBP alpha. *Proc. Natl. Acad. Sci. USA* **1995**, *92*, 9856–9860. [[CrossRef](#)] [[PubMed](#)]

14. Rosen, E.D.; Sarraf, P.; Troy, A.E.; Bradwin, G.; Moore, K.; Milstone, D.S.; Spiegelman, B.M.; Mortensen, R.M. PPAR γ is required for the differentiation of adipose tissue in vivo and in vitro. *Mol. Cell* **1999**, *4*, 611–617. [[CrossRef](#)]
15. Imai, T.; Takakuwa, R.; Marchand, S.; Dentz, E.; Bornert, J.-M.; Messaddeq, N.; Wendling, O.; Mark, M.; Desvergne, B.; Wahli, W.; et al. Peroxisome proliferator-activated receptor γ is required in mature white and brown adipocytes for their survival in the mouse. *Proc. Natl. Acad. Sci. USA* **2004**, *101*, 4543–4547. [[CrossRef](#)] [[PubMed](#)]
16. Tsai, Y.-S.; Maeda, N. PPAR γ : A critical determinant of body fat distribution in humans and mice. *Trends Cardiovasc. Med.* **2005**, *15*, 81–85. [[CrossRef](#)] [[PubMed](#)]
17. Agostini, M.; Schoenmakers, E.; Mitchell, C.; Szatmari, I.; Savage, D.; Smith, A.; Rajanayagam, O.; Semple, R.; Luan, J.A.; Bath, L.; et al. Non-DNA binding, dominant-negative, human PPAR γ mutations cause lipodystrophic insulin resistance. *Cell Metab.* **2006**, *4*, 303–311. [[CrossRef](#)] [[PubMed](#)]
18. Rosen, E.D.; MacDougald, O.A. Adipocyte differentiation from the inside out. *Nat. Rev. Mol. Cell Biol.* **2006**, *7*, 885–896. [[CrossRef](#)] [[PubMed](#)]
19. Zhu, Y.; Qi, C.; Korenberg, J.R.; Chen, X.-N.; Noya, D.; Rao, M.S.; Reddy, J.K. Structural organization of mouse peroxisome proliferator-activated receptor γ (mPPAR γ) gene: Alternative promoter use and different splicing yield two mPPAR γ isoforms. *Proc. Natl. Acad. Sci. USA* **1995**, *92*, 7921–7925. [[CrossRef](#)] [[PubMed](#)]
20. Willson, T.M.; Lambert, M.H.; Kliewer, S.A. Peroxisome proliferator-activated receptor γ and metabolic disease. *Annu. Rev. Biochem.* **2001**, *70*, 341–367. [[CrossRef](#)] [[PubMed](#)]
21. Vidal-Puig, A.; Jimenez-Liñan, M.; Lowell, B.B.; Hamann, A.; Hu, E.; Spiegelman, B.; Flier, J.S.; Moller, D.E. Regulation of PPAR γ gene expression by nutrition and obesity in rodents. *J. Clin. Invest.* **1996**, *97*, 2553–2561. [[CrossRef](#)] [[PubMed](#)]
22. Werman, A.; Hollenberg, A.; Solanes, G.; Bjørnbæk, C.; Vidal-Puig, A.J.; Flier, J.S. Ligand-independent activation domain in the N terminus of peroxisome proliferator-activated receptor γ (PPAR γ) Differential activity of PPAR γ 1 and-2 isoforms and influence of insulin. *J. Biol. Chem.* **1997**, *272*, 20230–20235. [[CrossRef](#)] [[PubMed](#)]
23. Yeh, W.-C.; Cao, Z.; Classon, M.; McKnight, S.L. Cascade regulation of terminal adipocyte differentiation by three members of the C/EBP family of leucine zipper proteins. *Genes Dev.* **1995**, *9*, 168–181. [[CrossRef](#)] [[PubMed](#)]
24. Harmon, A.W.; Patel, Y.M. Genistein inhibits CCAAT/enhancer-binding protein β (C/EBP β) activity and 3T3-L1 adipogenesis by increasing C/EBP homologous protein expression. *Biochem. J.* **2002**, *367*, 203–208. [[CrossRef](#)] [[PubMed](#)]
25. Schwarz, E.J.; Reginato, M.J.; Shao, D.; Krakow, S.L.; Lazar, M.A. Retinoic acid blocks adipogenesis by inhibiting C/EBP β -mediated transcription. *Mol. Cell Biol.* **1997**, *17*, 1552–1561. [[CrossRef](#)] [[PubMed](#)]
26. Jiang, L.; Zhang, N.-X.; Mo, W.; Wan, R.; Ma, C.-G.; Li, X.; Gu, Y.-L.; Yang, X.-Y.; Tang, Q.-Q.; Song, H.-Y. Rehmannia inhibits adipocyte differentiation and adipogenesis. *Biochem. Biophys. Res. Commun.* **2008**, *371*, 185–190. [[CrossRef](#)] [[PubMed](#)]
27. Urizar, N.L.; Liverman, A.B.; D’Nette, T.D.; Silva, F.V.; Ordentlich, P.; Yan, Y.; Gonzalez, F.J.; Heyman, R.A.; Mangelsdorf, D.J.; Moore, D.D. A natural product that lowers cholesterol as an antagonist ligand for FXR. *Science* **2002**, *296*, 1703–1706. [[CrossRef](#)] [[PubMed](#)]
28. Hu, Y.; Davies, G. Berberine increases expression of GATA-2 and GATA-3 during inhibition of adipocyte differentiation. *Phytomedicine* **2009**, *16*, 864–873. [[CrossRef](#)] [[PubMed](#)]
29. Kim, M.H.; Park, J.S.; Seo, M.S.; Jung, J.W.; Lee, Y.S.; Kang, K.S. Genistein and daidzein repress adipogenic differentiation of human adipose tissue-derived mesenchymal stem cells via Wnt/ β -catenin signalling or lipolysis. *Cell Prolif.* **2010**, *43*, 594–605. [[CrossRef](#)] [[PubMed](#)]
30. Ahn, J.; Lee, H.; Kim, S.; Ha, T. Curcumin-induced suppression of adipogenic differentiation is accompanied by activation of Wnt/ β -catenin signaling. *Am. J. Physiol.-Cell Physiol.* **2010**, *298*, C1510–C1516.
31. Lee, H.; Bae, S.; Kim, K.; Kim, W.; Chung, S.-I.; Yang, Y.; Yoon, Y. Shikonin inhibits adipogenesis by modulation of the WNT/ β -catenin pathway. *Life Sci.* **2011**, *88*, 294–301. [[CrossRef](#)] [[PubMed](#)]
32. Li, Y.-T.; Li, L.; Chen, J.; Hu, T.-C.; Huang, J.; Guo, Y.-W.; Jiang, H.-L.; Shen, X. 7-Chloroarctinone-b as a new selective PPAR γ antagonist potently blocks adipocyte differentiation. *Acta Pharmacol. Sin.* **2009**, *30*, 1351–1358. [[CrossRef](#)] [[PubMed](#)]

33. Park, U.-H.; Jeong, H.-S.; Jo, E.-Y.; Park, T.; Yoon, S.K.; Kim, E.-J.; Jeong, J.-C.; Um, S.-J. Piperine, a component of black pepper, inhibits adipogenesis by antagonizing PPAR γ activity in 3T3-L1 cells. *J. Agric. Food Chem.* **2012**, *60*, 3853–3860. [[CrossRef](#)] [[PubMed](#)]
34. Howitz, K.T.; Bitterman, K.J.; Cohen, H.Y.; Lamming, D.W.; Lavu, S.; Wood, J.G.; Zipkin, R.E.; Chung, P.; Kisielewski, A.; Zhang, L.-L. Small molecule activators of sirtuins extend *Saccharomyces cerevisiae* lifespan. *Nature* **2003**, *425*, 191–196. [[CrossRef](#)] [[PubMed](#)]
35. Choi, Y.; Um, S.; Park, T. Indole-3-carbinol directly targets SIRT1 to inhibit adipocyte differentiation. *Int. J. Obes.* **2013**, *37*, 881–884. [[CrossRef](#)] [[PubMed](#)]
36. Lai, C.-S.; Tsai, M.-L.; Badmaev, V.; Jimenez, M.; Ho, C.-T.; Pan, M.-H. Xanthigen suppresses preadipocyte differentiation and adipogenesis through down-regulation of PPAR γ and C/EBPs and modulation of SIRT-1, AMPK, and FoxO pathways. *J. Agric. Food Chem.* **2012**, *60*, 1094–1101. [[CrossRef](#)] [[PubMed](#)]
37. Lee, Y.K.; Lee, W.S.; Hwang, J.T.; Kwon, D.Y.; Surh, Y.J.; Park, O.J. Curcumin exerts antidifferentiation effect through AMPK α -PPAR- γ in 3T3-L1 adipocytes and antiproliferatory effect through AMPK α -COX-2 in cancer cells. *J. Agric. Food Chem.* **2008**, *57*, 305–310. [[CrossRef](#)] [[PubMed](#)]
38. Seo, J.B.; Choe, S.S.; Jeong, H.W.; Park, S.W.; Shin, H.J.; Choi, S.M.; Park, J.Y.; Choi, E.W.; Kim, J.B.; Seen, D.S. Anti-obesity effects of *Lysimachia foenum-graecum* characterized by decreased adipogenesis and regulated lipid metabolism. *Exp. Mol. Med.* **2011**, *43*, 205–215. [[CrossRef](#)] [[PubMed](#)]
39. Hwang, J.-T.; Park, I.-J.; Shin, J.-I.; Lee, Y.K.; Lee, S.K.; Baik, H.W.; Ha, J.; Park, O.J. Genistein, EGCG, and capsaicin inhibit adipocyte differentiation process via activating AMP-activated protein kinase. *Biochem. Biophys. Res. Commun.* **2005**, *338*, 694–699. [[CrossRef](#)] [[PubMed](#)]
40. Hwang, J.-T.; Kim, S.-H.; Lee, M.-S.; Kim, S.H.; Yang, H.-J.; Kim, M.-J.; Kim, H.-S.; Ha, J.; Kim, M.S.; Kwon, D.Y. Anti-obesity effects of ginsenoside Rh2 are associated with the activation of AMPK signaling pathway in 3T3-L1 adipocyte. *Biochem. Biophys. Res. Commun.* **2007**, *364*, 1002–1008. [[CrossRef](#)] [[PubMed](#)]
41. Chen, Y.-Y.; Lee, M.-H.; Hsu, C.-C.; Wei, C.-L.; Tsai, Y.-C. Methyl cinnamate inhibits adipocyte differentiation via activation of the CaMKK2–AMPK pathway in 3T3-L1 preadipocytes. *J. Agric. Food Chem.* **2012**, *60*, 955–963. [[CrossRef](#)] [[PubMed](#)]
42. He, Y.; Li, Y.; Zhao, T.; Wang, Y.; Sun, C. Ursolic acid inhibits adipogenesis in 3T3-L1 adipocytes through LKB1/AMPK pathway. *PLoS ONE* **2013**, *8*, e70135. [[CrossRef](#)] [[PubMed](#)]
43. Wang, T.; Wang, Y.; Kontani, Y.; Kobayashi, Y.; Sato, Y.; Mori, N.; Yamashita, H. Evodiamine improves diet-induced obesity in a uncoupling protein-1-independent manner: Involvement of antiadipogenic mechanism and extracellularly regulated kinase/mitogen-activated protein kinase signaling. *Endocrinology* **2008**, *149*, 358–366. [[CrossRef](#)] [[PubMed](#)]
44. Lin, J.; Della-Fera, M.A.; Baile, C.A. Green Tea Polyphenol Epigallocatechin Gallate Inhibits Adipogenesis and Induces Apoptosis in 3T3-L1 Adipocytes. *Obes. Res.* **2005**, *13*, 982–990. [[CrossRef](#)] [[PubMed](#)]
45. Kim, J.-K.; So, H.; Youn, M.-J.; Kim, H.-J.; Kim, Y.; Park, C.; Kim, S.-J.; Ha, Y.-A.; Chai, K.-Y.; Kim, S.-M. Hibiscus sabdariffa L. water extract inhibits the adipocyte differentiation through the PI3-K and MAPK pathway. *J. Ethnopharmacol.* **2007**, *114*, 260–267. [[CrossRef](#)] [[PubMed](#)]
46. Huang, T.H.W.; Kota, B.P.; Razmovski, V.; Roufogalis, B.D. Herbal or Natural Medicines as Modulators of Peroxisome Proliferator-Activated Receptors and Related Nuclear Receptors for Therapy of Metabolic Syndrome. *Basic Clin. Pharmacol. Toxicol.* **2005**, *96*, 3–14. [[CrossRef](#)] [[PubMed](#)]
47. Yim, M.-J.; Hosokawa, M.; Mizushima, Y.; Yoshida, H.; Saito, Y.; Miyashita, K. Suppressive effects of amarouciaxanthin A on 3T3-L1 adipocyte differentiation through down-regulation of PPAR γ and C/EBP α mRNA expression. *J. Agric. Food Chem.* **2011**, *59*, 1646–1652. [[CrossRef](#)] [[PubMed](#)]
48. Yang, J.-Y.; Della-Fera, M.A.; Rayalam, S.; Baile, C.A. Effect of xanthohumol and isoxanthohumol on 3T3-L1 cell apoptosis and adipogenesis. *Apoptosis* **2007**, *12*, 1953–1963. [[CrossRef](#)] [[PubMed](#)]
49. Kim, C.-S.; Park, W.-H.; Park, J.-Y.; Kang, J.-H.; Kim, M.-O.; Kawada, T.; Yoo, H.; Han, I.-S.; Yu, R. Capsaicin, a spicy component of hot pepper, induces apoptosis by activation of the peroxisome proliferator-activated receptor γ in HT-29 human colon cancer cells. *J. Med. Food* **2004**, *7*, 267–273. [[CrossRef](#)] [[PubMed](#)]
50. Zong, W.; Zhao, G.-Y. Corosolic acid isolation from the leaves of *Eriobotrya japonica* showing the effects on carbohydrate metabolism and differentiation of 3T3-L1 adipocytes. *Asia Pac. J. Clin. Nutr.* **2007**, *16*, 346–352. [[PubMed](#)]
51. Kubota, H.; Morii, R.; Kojima-Yuasa, A.; Huang, X.; Yano, Y.; Matsui-Yuasa, I. Effect of Zizyphus jujuba extract on the inhibition of adipogenesis in 3T3-L1 preadipocytes. *Am. J. Chin. Med.* **2009**, *37*, 597–608. [[CrossRef](#)] [[PubMed](#)]

52. Yang, J.-Y.; Della-Fera, M.A.; Rayalam, S.; Ambati, S.; Hartzell, D.L.; Park, H.J.; Baile, C.A. Enhanced inhibition of adipogenesis and induction of apoptosis in 3T3-L1 adipocytes with combinations of resveratrol and quercetin. *Life Sci.* **2008**, *82*, 1032–1039. [[CrossRef](#)] [[PubMed](#)]
53. Kim, M.-J.; Chang, U.-J.; Lee, J.-S. Inhibitory effects of fucoidan in 3T3-L1 adipocyte differentiation. *Mar. Biotechnol.* **2009**, *11*, 557–562. [[CrossRef](#)] [[PubMed](#)]
54. Jou, P.-C.; Ho, B.-Y.; Hsu, Y.-W.; Pan, T.-M. The effect of *Monascus* secondary polyketide metabolites, monascin and ankaflavin, on adipogenesis and lipolysis activity in 3T3-L1. *J. Agric. Food Chem.* **2010**, *58*, 12703–12709. [[CrossRef](#)] [[PubMed](#)]
55. Kim, O.Y.; Lee, S.M.; Do, H.; Moon, J.; Lee, K.H.; Cha, Y.J.; Shin, M.J. Influence of Quercetin-rich Onion Peel Extracts on Adipokine Expression in the Visceral Adipose Tissue of Rats. *Phytother. Res.* **2012**, *26*, 432–437. [[CrossRef](#)] [[PubMed](#)]
56. Lee, H.; Kang, R.; Yoon, Y. Shikonin inhibits fat accumulation in 3T3-L1 adipocytes. *Phytother. Res.* **2010**, *24*, 344–351. [[CrossRef](#)] [[PubMed](#)]
57. Siraj, F.M.; Kim, Y.J.; Natarajan, S.; Jung, S.K.; Yang, D.U.; Yang, D.C. Ginseng and obesity: Observations from assorted perspectives. *Food Sci. Biotechnol.* **2014**, *23*, 1007–1016. [[CrossRef](#)]
58. Liang, C.-H.; Chan, L.-P.; Chou, T.-H.; Chiang, F.-Y.; Yen, C.-M.; Chen, P.-J.; Ding, H.-Y.; Lin, R.-J. Brazilein from *Caesalpinia sappan* L. antioxidant inhibits adipocyte differentiation and induces apoptosis through caspase-3 activity and anthelmintic activities against *Hymenolepis nana* and *Anisakis simplex*. *Evid. Based Complement. Altern. Med.* **2013**, *2013*. [[CrossRef](#)]
59. Dudhia, Z.; Louw, J.; Muller, C.; Joubert, E.; de Beer, D.; Kinnear, C.; Pfeiffer, C. *Cyclopia maculata* and *Cyclopia subternata* (honeybush tea) inhibits adipogenesis in 3T3-L1 pre-adipocytes. *Phytomedicine* **2013**, *20*, 401–408. [[CrossRef](#)] [[PubMed](#)]
60. Ahn, E.K.; Oh, J.S. Lupenone Isolated from *Adenophora triphylla* var. japonica Extract Inhibits Adipogenic Differentiation through the Downregulation of PPAR γ in 3T3-L1 Cells. *Phytother. Res.* **2013**, *27*, 761–766. [[CrossRef](#)] [[PubMed](#)]
61. Chung, M.Y.; Yoon, Y.I.; Hwang, J.S.; Goo, T.W.; Yun, E.Y. Anti-obesity effect of *Allomyrina dichotoma* (Arthropoda: INSECTA) larvae ethanol extract on 3T3-L1 adipocyte differentiation. *Entomol. Res.* **2014**, *44*, 9–16. [[CrossRef](#)]
62. Gao, Y.; Yao, Y.; Zhu, Y.; Ren, G. Isoflavones in Chickpeas Inhibit Adipocyte Differentiation and Prevent Insulin Resistance in 3T3-L1 Cells. *J. Agric. Food Chem.* **2015**, *63*, 9696–9703. [[CrossRef](#)] [[PubMed](#)]
63. Hsu, C.-L.; Lin, Y.-J.; Ho, C.-T.; Yen, G.-C. Inhibitory effects of garcinol and pterostilbene on cell proliferation and adipogenesis in 3T3-L1 cells. *Food Funct.* **2012**, *3*, 49–57. [[CrossRef](#)] [[PubMed](#)]
64. Choi, J.S.; Kim, J.-H.; Ali, M.Y.; Min, B.-S.; Kim, G.-D.; Jung, H.A. *Coptis chinensis* alkaloids exert anti-adipogenic activity on 3T3-L1 adipocytes by downregulating C/EBP- α and PPAR- γ . *Fitoterapia* **2014**, *98*, 199–208. [[CrossRef](#)] [[PubMed](#)]
65. Nerlov, C. The C/EBP family of transcription factors: A paradigm for interaction between gene expression and proliferation control. *Trends Cell Biol.* **2007**, *17*, 318–324. [[CrossRef](#)] [[PubMed](#)]
66. Ron, D.; Habener, J.F. CHOP, a novel developmentally regulated nuclear protein that dimerizes with transcription factors C/EBP and LAP and functions as a dominant-negative inhibitor of gene transcription. *Genes Dev.* **1992**, *6*, 439–453. [[CrossRef](#)] [[PubMed](#)]
67. Lekstrom-Himes, J.; Xanthopoulos, K.G. Biological role of the CCAAT/enhancer-binding protein family of transcription factors. *J. Biol. Chem.* **1998**, *273*, 28545–28548. [[CrossRef](#)] [[PubMed](#)]
68. Dixon, R.A.; Ferreira, D. Genistein. *Phytochemistry* **2002**, *60*, 205–211. [[CrossRef](#)]
69. Choi, I.; Kim, Y.; Park, Y.; Seog, H.; Choi, H. Anti-obesity activities of fermented soygerm isoflavones by *Bifidobacterium breve*. *Biofactors* **2007**, *29*, 105–112. [[CrossRef](#)] [[PubMed](#)]
70. Guo, Y.; Wu, G.; Su, X.; Yang, H.; Zhang, J. Antiobesity action of a daidzein derivative on male obese mice induced by a high-fat diet. *Nutr. Res.* **2009**, *29*, 656–663. [[CrossRef](#)] [[PubMed](#)]
71. Bhathena, S.J.; Velasquez, M.T. Beneficial role of dietary phytoestrogens in obesity and diabetes. *Am. J. Clin. Nutr.* **2002**, *76*, 1191–1201. [[PubMed](#)]
72. Gudas, L.J.; Sporn, M.B.; Roberts, A.B. Cellular biology and biochemistry of the retinoids. *Retin. Biol. Chem. Med.* **1994**, *1994*, 443–520.
73. Makishima, M.; Okamoto, A.Y.; Repa, J.J.; Tu, H.; Learned, R.M.; Luk, A.; Hull, M.V.; Lustig, K.D.; Mangelsdorf, D.J.; Shan, B. Identification of a nuclear receptor for bile acids. *Science* **1999**, *284*, 1362–1365. [[CrossRef](#)] [[PubMed](#)]

74. Zhang, Y.; Kast-Woelbern, H.R.; Edwards, P.A. Natural structural variants of the nuclear receptor farnesoid X receptor affect transcriptional activation. *J. Biol. Chem.* **2003**, *278*, 104–110. [[CrossRef](#)] [[PubMed](#)]
75. Rizzo, G.; Disante, M.; Mencarelli, A.; Renga, B.; Gioiello, A.; Pellicciari, R.; Fiorucci, S. The farnesoid X receptor promotes adipocyte differentiation and regulates adipose cell function in vivo. *Mol. Pharmacol.* **2006**, *70*, 1164–1173. [[CrossRef](#)] [[PubMed](#)]
76. Yang, J.Y.; Della-Fera, M.A.; Baile, C.A. Guggulsterone Inhibits Adipocyte Differentiation and Induces Apoptosis in 3T3-L1 Cells. *Obesity* **2008**, *16*, 16–22. [[CrossRef](#)] [[PubMed](#)]
77. Yang, J.-Y.; Della-Fera, M.A.; Rayalam, S.; Ambati, S.; Baile, C.A. Enhanced pro-apoptotic and anti-adipogenic effects of genistein plus guggulsterone in 3T3-L1 adipocytes. *Biofactors* **2007**, *30*, 159–169. [[CrossRef](#)] [[PubMed](#)]
78. Rayalam, S.; Yang, J.-Y.; Della-Fera, M.A.; Park, H.J.; Ambati, S.; Baile, C.A. Anti-obesity effects of xanthohumol plus guggulsterone in 3T3-L1 adipocytes. *J. Med. Food* **2009**, *12*, 846–853. [[CrossRef](#)] [[PubMed](#)]
79. Tong, Q.; Dalgin, G.; Xu, H.; Ting, C.-N.; Leiden, J.M.; Hotamisligil, G.S. Function of GATA transcription factors in preadipocyte-adipocyte transition. *Science* **2000**, *290*, 134–138. [[CrossRef](#)] [[PubMed](#)]
80. Tong, Q.; Tsai, J.; Tan, G.; Dalgin, G.; Hotamisligil, G.S. Interaction between GATA and the C/EBP family of transcription factors is critical in GATA-mediated suppression of adipocyte differentiation. *Mol. Cell. Biol.* **2005**, *25*, 706–715. [[CrossRef](#)] [[PubMed](#)]
81. Huang, C.; Zhang, Y.; Gong, Z.; Sheng, X.; Li, Z.; Zhang, W.; Qin, Y. Berberine inhibits 3T3-L1 adipocyte differentiation through the PPAR γ pathway. *Biochem. Biophys. Res. Commun.* **2006**, *348*, 571–578. [[CrossRef](#)] [[PubMed](#)]
82. Hu, Y.; Davies, G.E. Berberine inhibits adipogenesis in high-fat diet-induced obesity mice. *Fitoterapia* **2010**, *81*, 358–366. [[PubMed](#)]
83. Hu, Y.; Fahmy, H.; Zjawiony, J.K.; Davies, G.E. Inhibitory effect and transcriptional impact of berberine and evodiamine on human white preadipocyte differentiation. *Fitoterapia* **2010**, *81*, 259–268. [[PubMed](#)]
84. Ross, S.E.; Hemati, N.; Longo, K.A.; Bennett, C.N.; Lucas, P.C.; Erickson, R.L.; MacDougald, O.A. Inhibition of adipogenesis by Wnt signaling. *Science* **2000**, *289*, 950–953. [[CrossRef](#)] [[PubMed](#)]
85. Freytag, S.O.; Geddes, T.J. Reciprocal Regulation of Adipogenesis by Myc and C/EBP (α). *Science* **1992**, *256*, 379–382. [[CrossRef](#)] [[PubMed](#)]
86. Fu, M.; Rao, M.; Bouras, T.; Wang, C.; Wu, K.; Zhang, X.; Li, Z.; Yao, T.-P.; Pestell, R.G. Cyclin D1 inhibits peroxisome proliferator-activated receptor γ -mediated adipogenesis through histone deacetylase recruitment. *J. Biol. Chem.* **2005**, *280*, 16934–16941. [[CrossRef](#)] [[PubMed](#)]
87. Ricote, M.; Li, A.C.; Willson, T.M.; Kelly, C.J.; Glass, C.K. The peroxisome proliferator-activated receptor- γ is a negative regulator of macrophage activation. *Nature* **1998**, *391*, 79–82. [[CrossRef](#)] [[PubMed](#)]
88. Kennell, J.A.; MacDougald, O.A. Wnt signaling inhibits adipogenesis through β -catenin-dependent and-independent mechanisms. *J. Biol. Chem.* **2005**, *280*, 24004–24010. [[CrossRef](#)] [[PubMed](#)]
89. Miller, J.R.; Hocking, A.M.; Brown, J.D.; Moon, R.T. Mechanism and function of signal transduction by the Wnt/ β -catenin and Wnt/Ca²⁺ pathways. *Oncogene* **1999**, *18*, 7860–7872. [[CrossRef](#)] [[PubMed](#)]
90. Cadigan, K.M.; Nusse, R. Wnt signaling: A common theme in animal development. *Genes Dev.* **1997**, *11*, 3286–3305. [[CrossRef](#)] [[PubMed](#)]
91. Lee, S.H.; Kim, B.; Oh, M.J.; Yoon, J.; Kim, H.Y.; Lee, K.J.; Lee, J.D.; Choi, K.Y. *Persicaria hydropiper* (L.) Spach and its Flavonoid Components, Isoquercitrin and Isorhamnetin, Activate the Wnt/ β -catenin Pathway and Inhibit Adipocyte Differentiation of 3T3-L1 Cells. *Phytother. Res.* **2011**, *25*, 1629–1635. [[CrossRef](#)] [[PubMed](#)]
92. Chawla, A.; Repa, J.J.; Evans, R.M.; Mangelsdorf, D.J. Nuclear receptors and lipid physiology: Opening the X-files. *Science* **2001**, *294*, 1866–1870. [[CrossRef](#)] [[PubMed](#)]
93. Yu, S.; Reddy, J.K. Transcription coactivators for peroxisome proliferator-activated receptors. *Biochim. Biophys. Acta* **2007**, *1771*, 936–951. [[CrossRef](#)] [[PubMed](#)]
94. Liu, H.L.; Guo, Y.W. Three new thiophene acetylenes from *Rhaphonticum uniflorum* (L.) DC. *Helv. Chim. Acta* **2008**, *91*, 130–135. [[CrossRef](#)]
95. Milne, J.C.; Denu, J.M. The Sirtuin family: Therapeutic targets to treat diseases of aging. *Curr. Opin. Chem. Biol.* **2008**, *12*, 11–17. [[CrossRef](#)] [[PubMed](#)]
96. Saunders, L.; Verdin, E. Sirtuins: Critical regulators at the crossroads between cancer and aging. *Oncogene* **2007**, *26*, 5489–5504. [[CrossRef](#)] [[PubMed](#)]

97. Picard, F.; Kurtev, M.; Chung, N.; Topark-Ngarm, A.; Senawong, T.; de Oliveira, R.M.; Leid, M.; McBurney, M.W.; Guarente, L. Sirt1 promotes fat mobilization in white adipocytes by repressing PPAR- γ . *Nature* **2004**, *429*, 771–776. [[CrossRef](#)] [[PubMed](#)]
98. Yu, C.; Markan, K.; Temple, K.A.; Deplewski, D.; Brady, M.J.; Cohen, R.N. The nuclear receptor corepressors NCoR and SMRT decrease peroxisome proliferator-activated receptor γ transcriptional activity and repress 3T3-L1 adipogenesis. *J. Biol. Chem.* **2005**, *280*, 13600–13605. [[CrossRef](#)] [[PubMed](#)]
99. Gerhart-Hines, Z.; Rodgers, J.T.; Bare, O.; Lerin, C.; Kim, S.H.; Mostoslavsky, R.; Alt, F.W.; Wu, Z.; Puigserver, P. Metabolic control of muscle mitochondrial function and fatty acid oxidation through SIRT1/PGC-1 α . *EMBO J.* **2007**, *26*, 1913–1923. [[CrossRef](#)] [[PubMed](#)]
100. Burns, J.; Yokota, T.; Ashihara, H.; Lean, M.E.; Crozier, A. Plant foods and herbal sources of resveratrol. *J. Agric. Food Chem.* **2002**, *50*, 3337–3340. [[CrossRef](#)] [[PubMed](#)]
101. Das, S.; Alagappan, V.K.; Bagchi, D.; Sharma, H.S.; Maulik, N.; Das, D.K. Coordinated induction of iNOS–VEGF–KDR–eNOS after resveratrol consumption: A potential mechanism for resveratrol preconditioning of the heart. *Vasc. Pharmacol.* **2005**, *42*, 281–289. [[CrossRef](#)] [[PubMed](#)]
102. El-Mowafy, A.M.; Alkhalaf, M. Resveratrol activates adenylyl-cyclase in human breast cancer cells: A novel, estrogen receptor-independent cytostatic mechanism. *Carcinogenesis* **2003**, *24*, 869–873. [[CrossRef](#)] [[PubMed](#)]
103. De la Lastra, C.A.; Villegas, I. Resveratrol as an antioxidant and pro-oxidant agent: Mechanisms and clinical implications. *Biochem. Soc. Trans.* **2007**, *35*, 1156–1160. [[CrossRef](#)] [[PubMed](#)]
104. Baile, C.A.; Yang, J.Y.; Rayalam, S.; Hartzell, D.L.; Lai, C.Y.; Andersen, C.; Della-Fera, M.A. Effect of resveratrol on fat mobilization. *Ann. N. Y. Acad. Sci.* **2011**, *1215*, 40–47. [[CrossRef](#)] [[PubMed](#)]
105. Kaeberlein, M.; McDonagh, T.; Heltweg, B.; Hixon, J.; Westman, E.A.; Caldwell, S.D.; Napper, A.; Curtis, R.; DiStefano, P.S.; Fields, S. Substrate-specific activation of sirtuins by resveratrol. *J. Biol. Chem.* **2005**, *280*, 17038–17045. [[CrossRef](#)] [[PubMed](#)]
106. Högnadóttir, Á.; Rouseff, R.L. Identification of aroma active compounds in orange essence oil using gas chromatography–olfactometry and gas chromatography–mass spectrometry. *J. Chromatogr. A* **2003**, *998*, 201–211. [[CrossRef](#)]
107. Fischer-Posovszky, P.; Kukulius, V.; Tews, D.; Unterkircher, T.; Debatin, K.-M.; Fulda, S.; Wabitsch, M. Resveratrol regulates human adipocyte number and function in a Sirt1-dependent manner. *Am. J. Clin. Nutr.* **2010**, *92*, 5–15. [[CrossRef](#)] [[PubMed](#)]
108. Pang, W.J.; Sun, S.D.; Bai, L.; Yang, Y.J.; Yang, G.S. Effects of resveratrol on pig primary preadipocytes proliferation, differentiation and transcription expression of Sirt1 gene. *Sheng Wu Gong Cheng Xue Bao* **2006**, *22*, 850–855.
109. Pacholec, M.; Bleasdale, J.E.; Chrunchy, B.; Cunningham, D.; Flynn, D.; Garofalo, R.S.; Griffith, D.; Griffior, M.; Loulakis, P.; Pabst, B.; et al. SRT1720, SRT2183, SRT1460, and resveratrol are not direct activators of SIRT1. *J. Biol. Chem.* **2010**, *285*, 8340–8351. [[CrossRef](#)] [[PubMed](#)]
110. Beher, D.; Wu, J.; Cumine, S.; Kim, K.W.; Lu, S.C.; Atangan, L.; Wang, M. Resveratrol is not a direct activator of SIRT1 enzyme activity. *Chem. Biol. Drug Des.* **2009**, *74*, 619–624. [[CrossRef](#)] [[PubMed](#)]
111. De Boer, V.C.; de Goffau, M.C.; Arts, I.C.; Hollman, P.C.; Keijer, J. SIRT1 stimulation by polyphenols is affected by their stability and metabolism. *Mech. Ageing Dev.* **2006**, *127*, 618–627. [[CrossRef](#)] [[PubMed](#)]
112. Choi, K.-C.; Jung, M.G.; Lee, Y.-H.; Yoon, J.C.; Kwon, S.H.; Kang, H.-B.; Kim, M.-J.; Cha, J.-H.; Kim, Y.J.; Jun, W.J.; et al. Epigallocatechin-3-gallate, a histone acetyltransferase inhibitor, inhibits EBV-induced B lymphocyte transformation via suppression of RelA acetylation. *Cancer Res.* **2009**, *69*, 583–592. [[CrossRef](#)] [[PubMed](#)]
113. Hwang, J.-T.; Kwon, D.Y.; Yoon, S.H. AMP-activated protein kinase: A potential target for the diseases prevention by natural occurring polyphenols. *New Biotechnol.* **2009**, *26*, 17–22. [[CrossRef](#)] [[PubMed](#)]
114. Horman, S.; Browne, G.J.; Krause, U.; Patel, J.V.; Vertommen, D.; Bertrand, L.; Lavoigne, A.; Hue, L.; Proud, C.G.; Rider, M.H. Activation of AMP-activated protein kinase leads to the phosphorylation of elongation factor 2 and an inhibition of protein synthesis. *Curr. Biol.* **2002**, *12*, 1419–1423. [[CrossRef](#)]
115. Daval, M.; Fougelle, F.; Ferré, P. Functions of AMP-activated protein kinase in adipose tissue. *J. Physiol.* **2006**, *574*, 55–62. [[CrossRef](#)] [[PubMed](#)]
116. Leff, T. AMP-activated protein kinase regulates gene expression by direct phosphorylation of nuclear proteins. *Biochem. Soc. Trans.* **2003**, *31*, 224–227. [[CrossRef](#)] [[PubMed](#)]

117. Habinowski, S.A.; Witters, L.A. The effects of AICAR on adipocyte differentiation of 3T3-L1 cells. *Biochem. Biophys. Res. Commun.* **2001**, *286*, 852–856. [[CrossRef](#)] [[PubMed](#)]
118. Xi, X.; Han, J.; Zhang, J.-Z. Stimulation of glucose transport by AMP-activated protein kinase via activation of p38 mitogen-activated protein kinase. *J. Biol. Chem.* **2001**, *276*, 41029–41034. [[CrossRef](#)] [[PubMed](#)]
119. Hu, E.; Kim, J.B.; Sarraf, P.; Spiegelman, B.M. Inhibition of adipogenesis through MAP kinase-mediated phosphorylation of PPAR gamma. *Science* **1996**, *274*, 2100. [[CrossRef](#)] [[PubMed](#)]
120. Diradourian, C.; Girard, J.; Pégrier, J.-P. Phosphorylation of PPARs: From molecular characterization to physiological relevance. *Biochimie* **2005**, *87*, 33–38. [[CrossRef](#)] [[PubMed](#)]
121. Lee, Y.S.; Kim, W.S.; Kim, K.H.; Yoon, M.J.; Cho, H.J.; Shen, Y.; Ye, J.-M.; Lee, C.H.; Oh, W.K.; Kim, C.T.; et al. Berberine, a natural plant product, activates AMP-activated protein kinase with beneficial metabolic effects in diabetic and insulin-resistant states. *Diabetes* **2006**, *55*, 2256–2264. [[CrossRef](#)] [[PubMed](#)]
122. Cobb, M.H.; Goldsmith, E.J. How MAP kinases are regulated. *J. Biol. Chem.* **1995**, *270*, 14843–14846. [[CrossRef](#)] [[PubMed](#)]
123. Robinson, M.J.; Cobb, M.H. Mitogen-activated protein kinase pathways. *Curr. Opin. Cell Biol.* **1997**, *9*, 180–186. [[CrossRef](#)]
124. Arbabi, S.; Maier, R.V. Mitogen-activated protein kinases. *Crit. Care Med.* **2002**, *30*, S74–S79. [[CrossRef](#)] [[PubMed](#)]
125. De Mora, J.F.; Porrás, A.; Ahn, N.; Santos, E. Mitogen-activated protein kinase activation is not necessary for, but antagonizes, 3T3-L1 adipocytic differentiation. *Mol. Cell. Biol.* **1997**, *17*, 6068–6075. [[CrossRef](#)]
126. Bost, F.; Aouadi, M.; Caron, L.; Binétruy, B. The role of MAPKs in adipocyte differentiation and obesity. *Biochimie* **2005**, *87*, 51–56. [[CrossRef](#)] [[PubMed](#)]
127. Prusty, D.; Park, B.-H.; Davis, K.E.; Farmer, S.R. Activation of MEK/ERK signaling promotes adipogenesis by enhancing peroxisome proliferator-activated receptor γ (PPAR γ) and C/EBP α gene expression during the differentiation of 3T3-L1 preadipocytes. *J. Biol. Chem.* **2002**, *277*, 46226–46232. [[CrossRef](#)] [[PubMed](#)]
128. Hung, P.-F.; Wu, B.-T.; Chen, H.-C.; Chen, Y.-H.; Chen, C.-L.; Wu, M.-H.; Liu, H.-C.; Lee, M.-J.; Kao, Y.-H. Antimitogenic effect of green tea (–)-epigallocatechin gallate on 3T3-L1 preadipocytes depends on the ERK and Cdk2 pathways. *Am. J. Physiol. Cell Physiol.* **2005**, *288*, C1094–C1108. [[CrossRef](#)] [[PubMed](#)]
129. Yun, J.W. Possible anti-obesity therapeutics from nature—A review. *Phytochemistry* **2010**, *71*, 1625–1641. [[CrossRef](#)] [[PubMed](#)]
130. González-Castejón, M.; Rodríguez-Casado, A. Dietary phytochemicals and their potential effects on obesity: A review. *Pharmacol. Res.* **2011**, *64*, 438–455. [[CrossRef](#)] [[PubMed](#)]
131. Andersen, C.; Rayalam, S.; Della-Fera, M.A.; Baile, C.A. Phytochemicals and adipogenesis. *Biofactors* **2010**, *36*, 415–422. [[CrossRef](#)] [[PubMed](#)]
132. Zizola, C.; Frey, S.; Jitngarmkusol, S.; Kadereit, B.; Yan, N.; Vogel, S. Cellular retinol-binding protein type I (CRBP-I) regulates adipogenesis. *Mol. Cell. Biol.* **2010**, *30*, 3412–3420. [[CrossRef](#)] [[PubMed](#)]
133. Jing, E.; Gesta, S.; Kahn, C.R. SIRT2 regulates adipocyte differentiation through FoxO1 acetylation/deacetylation. *Cell Metab.* **2007**, *6*, 105–114. [[CrossRef](#)] [[PubMed](#)]
134. Tsukahara, T.; Tsukahara, R.; Fujiwara, Y.; Yue, J.; Cheng, Y.; Guo, H.; Bolen, A.; Zhang, C.; Balazs, L.; Re, F.; et al. Phospholipase D2-dependent inhibition of the nuclear hormone receptor PPAR γ by cyclic phosphatidic acid. *Mol. Cell* **2010**, *39*, 421–432. [[CrossRef](#)] [[PubMed](#)]
135. Nanbu-Wakao, R.; Morikawa, Y.; Matsumura, I.; Masuho, Y.; Muramatsu, M.-A.; Senba, E.; Wakao, H. Stimulation of 3T3-L1 adipogenesis by signal transducer and activator of transcription 5. *Mol. Endocrinol.* **2002**, *16*, 1565–1576. [[CrossRef](#)] [[PubMed](#)]

