

Supplementary Materials:

Phytotoxic and Antifungal Metabolites from *Curvularia crepinii* QTYC-1 Isolated from the Gut of *Pantala flavescens*

Caiping Yin, Liping Jin, Feifei Sun, Xiao Xu, Mingwei Shao, Ying-Lao Zhang

Figure S1. ^1H -NMR spectrum (600 MHz) of the compound **2** in Acetone- d_6 .

Figure S2. ^{13}C -NMR spectrum (150 MHz) of compound **2** in Acetone- d_6 .

Figure S3. DEPT 135 spectrum of compound **2** in Acetone- d_6 .

Figure S4. ^1H - ^1H COSY spectrum of compound **2** in Acetone- d_6 .

Figure S5. HMQC spectrum of compound **2** in Acetone- d_6 .

Figure S6. HMBC spectrum of compound **2** in Acetone- d_6 .




Figure S1. ^1H -NMR spectrum (600 MHz) of the compound **2** in Acetone- d_6 .




Figure S2. ^{13}C -NMR spectrum (150 MHz) of compound **2** in Acetone- d_6 .




Figure S3. DEPT 135 spectrum of compound **2** in Acetone- d_6 .




Figure S4. ^1H - ^1H COSY spectrum of compound **2** in Acetone- d_6 .




Figure S5. HMQC spectrum of compound **2** in Acetone- d_6 .




Figure S6. HMBC spectrum of compound **2** in Acetone- d_6 .