Supporting information

Nanomaterials based on Fe₃O₄ and phthalocyanines derived from cashew nut shell liquid

Viviane G. P. Ribeiro ¹, João P. F. Mota ¹, Antônio E. Costa Júnior ¹, Nayane M. A. Lima ¹, Pierre B. A. Fechine ², Juliano C. Denardin ³, Luigi Carbone ⁴, Ermelinda Bloise ⁵, Giuseppe Mele ^{5*} and Selma E. Mazzetto ¹

- ¹ Laboratory of Products and Process Technology (LPT), Organic and Inorganic Chemistry Department, Federal University of Ceará (UFC), Campus do Pici, 60440-900, Fortaleza-CE, Brazil; vivianegpribeiro@live.com (V.R.); jpfmpro@gmail.com (J.P.); eufraziojr@yahoo.com.br (A.E); nayaneal@yahoo.com.br (N. L.); selma@ufc.br (S.M.)
- ² Group of Chemistry of Advanced Materials (GQMat) Department of Analytical Chemistry and Physical-Chemistry, Federal University of Ceará – UFC, Campus do Pici, CP 12100, CEP 60451-970 Fortaleza, CE, Brazil; fechine@ufc.br (P.F.)
- ³ Department of Physics, Universidad de Santiago de Chile and CEDENNA, USACH, Av. Ecuador, 3493 Santiago, Chile; juliano.denardin@usach.cl (J.D.)
- ⁴ CNR NANOTEC-Istituto di Nanotecnologia, c/o Campus Ecotekne, Università del Salento, Via Monteroni, 73100 Lecce, Italy; *luigi.carbone@*nanotec.*cnr*.it (L.C.)
- ⁵ Department of Engineering for Innovation, University of Salento, Via Arnesano, 73100, Lecce, Italy; ermelinda.bloise@unisalento.it (E.B.); giuseppe.mele@unisalento.it (G.M.)
- * Correspondence: giuseppe.mele@unisalento.it



Figure S2. ¹H NMR (300 MHz, CDCl₃) of the (a) Pc, (b) CoPc, (c) CuPc, (d) NiPc and (e) ZnPc.



Figure S3. Mass Spectra (MALDI-TOF-MS) of the (a) Pc, (b) CuPc, (c) NiPc, (d) ZnPc and (e) CoPc.







Figure 1. ¹³C NMR (300 MHz, CDCl₃) of the (a) Pc, (b) CoPc, (c) CuPc, (d) NiPc and (e) ZnPc.