## Supplementary Materials: Identification of Antiinflammatory Compounds from Hawaiian Noni (*Morinda citrifolia* L.) Fruit Juice

Dahae Lee<sup>1,+</sup>, Jae Sik Yu<sup>2,+</sup>, Peng Huang<sup>3</sup>, Mallique Qader<sup>3</sup>, Arulmani Manavalan<sup>3</sup>, Xiaohua Wu<sup>3</sup>, Jin-Chul Kim<sup>4</sup>, Changhyun Pang<sup>5</sup>, Shugeng Cao<sup>3,\*</sup>, Ki Sung Kang<sup>1,\*</sup> and Ki Hyun Kim<sup>2,\*</sup>

- <sup>1</sup> College of Korean Medicine, Gachon University, Seongnam 13120, Republic of Korea; pjsldh@naver.com (D.L.)
- <sup>2</sup> School of Pharmacy, Sungkyunkwan University, Suwon 16419, Republic of Korea; jsyu@bu.edu (J.S.Y.)
- <sup>3</sup> Daniel K. Inouye College of Pharmacy, University of Hawaii at Hilo, Hilo, Hawaii 96720, United States; great7701@126.com (P.H.); mqader@hawaii.edu (M.Q.); arulmanim@gmail.com (A.M.); xiaohua3@hawaii.edu (X.W.)
- <sup>4</sup> KIST Gangneung Institute of Natural products, Natural Product Informatics Research Center, Gangneung, 25451, Republic of Korea; jckim@kist.re.kr (J.C.K.)
- <sup>5</sup> School of Chemical Engineering, Sungkyunkwan University, Suwon 16419, Republic of Korea; chpang@skku.edu (C.P.)

<sup>+</sup>These authors contributed equally to this work

**Figure S1.** <sup>1</sup>H-NMR spectrum of compound **1** (in MeOH-*d*<sub>4</sub>)

Figure S2. <sup>1</sup>H-NMR spectrum of compound 2 (in D<sub>2</sub>O-*d*<sub>2</sub>)

Figure S3. HSQC spectrum of compound 2 (in D<sub>2</sub>O-d<sub>2</sub>)

Figure S4. <sup>1</sup>H-NMR spectrum of compound 3 (in MeOH-d<sub>4</sub>)

Figure S5. <sup>1</sup>H-NMR spectrum of compound 4 (in MeOH-*d*<sub>4</sub>)

Figure S6. 1H-NMR spectrum of compound 5 (in MeOH-d4)



Figure S2. <sup>1</sup>H-NMR spectrum of compound 2 (in H<sub>2</sub>O-d<sub>2</sub>)



**Figure S4**. <sup>1</sup>H-NMR spectrum of compound **3** (in MeOH-*d*<sub>4</sub>)



Figure S6. <sup>1</sup>H-NMR spectrum of compound 5 (in MeOH-*d*<sub>4</sub>)