

Supplementary Material for

Development of Water-In-Oil Emulsions as Delivery Vehicles and Testing with a Natural Antimicrobial Extract

Giovana Colucci^{1,2}, Arantzazu Santamaria-Echart¹, Samara C. Silva^{1,3}, Isabel P.
M. Fernandes^{1,*}, Caroline C. Sipoli² and Maria F. Barreiro^{1,*}

- ¹ Mountain Research Center (CIMO), Polytechnic Institute of Bragança, Santa Apolónia Campus, 5300-253 Bragança, Portugal
- ² Department of Chemical Engineering, Federal University of Technology – Paraná, Apucarana, Brazil
- ³ Laboratory of Separation and Reaction Engineering – Laboratory of Catalysis and Materials (LSRE/LCM) Department of Chemical Engineering, Faculty of Engineering University of Porto, Rua Dr. Roberto Frias, S/N, 4200-465 Porto, Portugal

* Corresponding author: e-mail: ipmf@ipb.pt

* Corresponding author: e-mail: barreiro@ipb.pt




Figure S1. Optical microscopy of the produced 40/60 W/O base emulsions. After 21 HPH cycles: (a) S80/T80 54/46; (b) S80/T80 80/20; (c) S85/T80 80/20. After 24 HPH cycles: (d) S80/T80 54/46; (e) S80/T80 80/20; (f) S85/T80 80/20. Bar = 10 μm , 200 \times magnification.




Figure S2. Optical microscopy of the produced 40/60 W/O emulsions added with cinnamon extract. Primary emulsions: (a) 1.25%; (b) 2.5%; (c) 3.75%; (d) 5%. After 12 HPH cycles: (e) 1.25%; (f) 2.5%; (g) 3.75%; (h) 5%. Bar = 10 μm , 200 \times magnification.