

β -Cyclodextrin-Polyacrylamide Hydrogel for Removal of Organic Micropollutants from Water

Xia Song ^{1,†}, Nana Nyarko Mensah ^{1,2,†}, Yuting Wen ¹, Jingling Zhu ¹, Zhongxing Zhang ¹, Wui Siew Tan ², Xinwei Chen ² and Jun Li ^{1,*}

¹ Department of Biomedical Engineering, Faculty of Engineering, National University of Singapore, 7 Engineering Drive 1, Singapore 117574, Singapore; a0045788@u.nus.edu (X.S.); nn.mensah@u.nus.edu (N.N.M.); bieweny@nus.edu.sg (Y.W.); erizhuj@nus.edu.sg (J.Z.); biezhozh@nus.edu.sg (Z.Z.)

² Institute of Materials Research and Engineering, A*STAR (Agency for Science, Technology and Research), 2 Fusionopolis Way, Singapore 138634, Singapore; wuisiew@gmail.com (W.S.T.); CHEN_Xinwei@nrf.gov.sg (X.C.)

* Correspondence: jun-li@nus.edu.sg or jun-li@u.nus.edu; Tel. +65-65-167-273

† These authors contribute equally.




Figure S1. The 600 MHz ^1H NMR spectra of (A) pure β -CD and (B) β -CD-MA measured in DMSO- d_6 at room temperature.




Figure S2. FTIR spectra of (A) β -CD and (B) β -CD-MA.




Figure S3. XPS survey spectra of (A) PAAm gel and (B) β -CD-PAAm gel.