

*Supplementary Material*

# Flavonoid Glycosides in *Brassica* Species Respond to UV-B Dependence on Exposure Time and Adaptation Time

Susanne Neugart <sup>1,\*</sup> and Christiane Bumke-Vogt <sup>2</sup>

<sup>1</sup> Division Quality and Sensory of Plant Products, Georg-August-Universität Göttingen, Carl-Sprengel-Weg 1, D-37075 Goettingen, Germany

<sup>2</sup> Leibniz-Institute of Vegetable and Ornamental Crops, Theodor-Echtermeyer-Weg 1, 14979 Grossbeeren, Germany; bumke@igzев.de

\* Correspondence: susanne.neugart@uni-goettingen.de; Tel: +49-0551-39-27958

Table S1: Kaempferol diglycosides of different *Brassica* species in µg/g DW. Asterix indicate differences due to the UV-B treatment at the certain time point significant at \*  $p \leq 0.05$ , \*\*  $p \leq 0.01$ , \*\*\*  $p \leq 0.005$

K: kaempferol; soph: sophorose; glc: glucose; cou: coumaroyl; caf: caffeoyl; fer: feruloyl; hfer: hydroxyferuloyl; sin: sinapoyl; h: hour; cond: condition

<i>Brassica</i>		K-3-soph	K-3-cou-soph	K-3-caf-soph	K-3-fer-soph	K-3-hfer-soph	K-3-sin-soph
<i>rapa</i>	Day 1	h	n.d.	n.d.		**	**
		cond					*
		h × cond					
	Day 4	h	n.d.	n.d.		*	
		cond					*
		h × cond					
<i>nigra</i>	Day 7	h	n.d.	n.d.		***	*
		cond				***	
		h × cond					
	Day 1	h	n.d.	n.d.	***	n.d.	***
		cond			**		**

		h × cond		***		
<i>oleracea</i>	Day 4	h	n.d.	n.d.	***	n.d.
		cond				*
		h × cond				
	Day 7	h	n.d.	n.d.		n.d.
		cond			*	
		h × cond				***
	Day 1	h	n.d.	n.d.		n.d.
		cond				
		h × cond				
<i>junccea</i>	Day 4	h	n.d.	n.d.	**	n.d.
		cond				*
		h × cond				
	Day 7	h	n.d.	n.d.	***	n.d.
		cond				**
		h × cond				
	Day 1	h	n.d.	n.d.	***	n.d.
		cond			*	
		h × cond				*
<i>napus</i>	Day 4	h	n.d.	n.d.		n.d.
		cond			*	
		h × cond				
	Day 7	h	n.d.	n.d.		n.d.
		cond				
		h × cond				***
	Day 1	h	n.d.	***		n.d.
		cond				
		h × cond				
	Day 4	h	n.d.			n.d.
		cond				

		h × cond				
	Day 7	h	n.d.	*	*	***
		cond				
		h × cond				
<i>carinata</i>	Day 1	h	n.d.	n.d.	n.d.	*
		cond				***
		h × cond				
	Day 4	h	n.d.	n.d.	n.d.	*
		cond				*
		h × cond				
	Day 7	h	n.d.	n.d.	n.d.	
		cond				***
		h × cond				

Table S2: Quercetin diglycosides of different *Brassica* species in µg/g DW. Asterix indicate differences due to the UV-B treatment at the certain time point significant at \*  $p \leq 0.05$ , \*\*  $p \leq 0.01$ , \*\*\*  $p \leq 0.005$

Q: quercetin; soph: sophorose; glc: glucose; cou: coumaroyl; caf: caffeoyl; fer: feruloyl; hfer: hydroxyferuloyl; sin: sinapoyl; h: hour; cond: condition

<i>Brassica</i>		Q-3-soph	Q-3-cou-soph	Q-3-caf-soph	Q-3-fer-soph	Q-3-hfer-soph	Q-3-sin-soph
<i>rapa</i>	Day 1	h cond h × cond	n.d.	n.d.	n.d.	n.d.	n.d.
	Day 4	h cond h × cond	n.d.	n.d.	n.d.	n.d.	n.d.
	Day 7	h cond h × cond	n.d.	n.d.	n.d.	n.d.	n.d.
<i>nigra</i>	Day 1	h cond h × cond	n.d.	n.d.	n.d.	n.d.	n.d.
	Day 4	h cond h × cond	n.d.	n.d.	n.d.	n.d.	n.d.
	Day 7	h cond h × cond	n.d.	n.d.	n.d.	n.d.	n.d.
<i>oleracea</i>	Day 1	h cond h × cond	n.d.	n.d.	n.d.	n.d.	n.d.
	Day 4	h cond h × cond	n.d.	n.d.	n.d.	n.d.	*
	Day 7	h cond h × cond	n.d.	n.d.	n.d.	n.d.	n.d.

						*
<i>juncea</i>	Day 1	cond				
		h × cond				
		h	n.d.	n.d.	n.d.	n.d.
	Day 4	cond				
		h × cond				
		h	n.d.	n.d.	n.d.	n.d.
	Day 7	cond				
		h × cond				
		h	n.d.	n.d.	n.d.	n.d.
<i>napus</i>	Day 1	cond				
		h × cond				
		h	n.d.	n.d.	n.d.	n.d.
	Day 4	cond				
		h × cond				
		h	n.d.	n.d.	n.d.	n.d.
	Day 7	cond				
		h × cond				
		h	n.d.	n.d.	n.d.	n.d.
<i>carinata</i>	Day 1	cond				
		h × cond				
		h	n.d.	n.d.	n.d.	n.d.
	Day 4	cond				
		h × cond				
		h	n.d.	n.d.	n.d.	n.d.
	Day 7	cond				
		h × cond				
		h	n.d.	n.d.	n.d.	n.d.

Table S3: Kaempferol triglycosides of different *Brassica* species in µg/g DW. Asterix indicate differences due to the UV-B treatment at the certain time point significant at \*  $p \leq 0.05$ , \*\*  $p \leq 0.01$ , \*\*\*  $p \leq 0.005$

K: kaempferol; soph: sophoroside; glc: glucose; cou: coumaroyl; caf: caffeoyl; fer: feruloyl; hfer: hydroxyferuloyl; sin: sinapoyl; h: hour; cond: condition

		<i>Brassica</i>	K-3-soph-7-glc	K-3-cou-7-glc	K-3-caf-7-glc	K-3-fer-7-glc	K-3-hfer-7-glc	K-3-sin-7-glc	K-3-disin-7-glc	K-3-sin,fer-7-glc
<i>rapa</i>	Day 1	h cond  h × cond								
	Day 4	h cond  h × cond		**			*		**	
	Day 7	h cond  h × cond			*					
<i>nigra</i>	Day 1	h cond  h × cond				*	*		n.d.	n.d.
	Day 4	h cond  h × cond		***					n.d.	n.d.
	Day 7	h cond  h × cond			*				n.d.	n.d.
<i>oleracea</i>	Day 1	h cond  h × cond						***	***	
					*					

<i>junccea</i>	Day 4	h	***	***	***	***	***	**	***
		cond	**	*	***	*	***		
		h × cond		**	*	***	***		
	Day 7	h							
		cond		***	**	***	*	***	***
		h × cond	***						
	Day 1	h				*			
		cond		*		*			
		h × cond							
<i>napus</i>	Day 4	h		*			***		
		cond							
		h × cond							
	Day 7	h					***		
		cond	*	*			*		
		h × cond							
	Day 1	h			***	***	n.d.	n.d.	
		cond							
		h × cond							
<i>carinata</i>	Day 4	h				***	n.d.	n.d.	
		cond		*					
		h × cond			*				
	Day 7	h							
		cond							
		h × cond					n.d.	n.d.	
	Day 1	h		*					
		cond							

	h × cond		*			
Day 4	h	*	***		n.d.	n.d.
	cond	*	***			
	h × cond	*	*			
Day 7	h	*	***		n.d.	n.d.
	cond					
	h × cond					

Table S4: Quercetin triglycosides of different *Brassica* species in µg/g DW. Asterix indicate differences due to the UV-B treatment at the certain time point significant at \*  $p \leq 0.05$ , \*\*  $p \leq 0.01$ , \*\*\*  $p \leq 0.005$

Q: quercetin; soph: 9inapoyl9; glc: glucose; cou: coumaroyl; caf: caffeoyl; fer: feruloyl; hfer: hydroxyferuloyl; sin: 9inapoyl; h: hour; cond: condition

<i>Brassica</i>		Q-3-soph-7-glc	Q-3-cou-soph-7-glc	Q-3-caf-soph-7-glc	Q-3-fer-soph-7-glc	Q-3-hfer-soph-7-glc	Q-3-sin-soph-7-glc
<i>rapa</i>	Day 1	h cond h × cond		n.d.		n.d.	
			*				
	Day 4	h cond h × cond		n.d.		n.d.	**
			*				
	Day 7	h cond h × cond		n.d.		n.d.	
			*				
<i>nigra</i>	Day 1	h cond h × cond		n.d.	n.d.	n.d.	***
							**
							***
	Day 4	h cond h × cond		n.d.	n.d.	n.d.	
			*				
	Day 7	h cond h × cond		n.d.	n.d.	n.d.	
<i>oleracea</i>	Day 1	h cond h × cond		n.d.		n.d.	***
	Day 4	h cond	**	n.d.	***	n.d.	**
					***		

		h × cond	***			
<i>junccea</i>	Day 7	h	n.d.	n.d.	n.d.	
		cond				***
		h × cond				
	Day 1	h	n.d. *	n.d.	n.d.	
		cond				
		h × cond		*		
	Day 4	h	n.d.	n.d.	n.d.	
		cond				
		h × cond				
<i>napus</i>	Day 7	h	n.d.	n.d.	n.d.	
		cond				
		h × cond				
	Day 1	h	n.d.	n.d.	n.d.	n.d.
		cond				
		h × cond				
	Day 4	h	n.d.	n.d.	n.d.	n.d.
		cond				
		h × cond				
<i>carinata</i>	Day 7	h	n.d.	n.d.	n.d.	n.d.
		cond				
		h × cond				
	Day 1	h	n.d.	n.d.	***	
		cond			***	
		h × cond			*	
	Day 4	h	n.d.	n.d.	n.d.	n.d.
		cond	*			
		h × cond	*			
	Day 7	h	n.d.	n.d.	n.d.	n.d.

cond

$h \times cond$

Table S5: Kaempferol tetraglycosides of different *Brassica* species in µg/g DW. Asterix indicate differences due to the UV-B treatment at the certain time point significant at \*  $p \leq 0.05$ , \*\*  $p \leq 0.01$ , \*\*\*  $p \leq 0.005$

K: kaempferol; soph: 12inapoyl12; glc: glucose; cou: coumaroyl; caf: caffeoyl; fer: feruloyl; hfer: hydroxyferuloyl; sin: 12inapoyl; h: hour; cond: condition

<i>Brassica</i>		K-3-soph-7-diglc	K-3-cou-soph-7-diglc	K-3-caf-soph-7-diglc	K-3-fer-soph-7-diglc	K-3-hfer-soph-7-diglc	K-3-sin-soph-7-diglc	K-3-disin-striglc-7-glc
<i>rapa</i>	Day 1	h cond h × cond	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
	Day 4	h cond h × cond	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
	Day 7	h cond h × cond	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
<i>nigra</i>	Day 1	h cond h × cond	n.d.	n.d.	***		n.d.	
						**		
	Day 4	h cond h × cond	n.d.	n.d.	***	***	***	n.d.
	Day 7	h cond h × cond	n.d.	n.d.	*		n.d.	
						*		
<i>oleracea</i>	Day 1	h cond h × cond	n.d.	n.d.	n.d.	**		
	Day 4	h cond h × cond	n.d.	n.d.	n.d.	*	***	*
								**

		cond			*		**
		h × cond					
	Day 7	h	n.d.	n.d.	n.d.	**	*
		cond				***	
		h × cond				***	*
<i>junccea</i>	Day 1	h	n.d.	n.d.	n.d.	n.d.	n.d.
		cond					
		h × cond					
	Day 4	h	n.d.	n.d.	n.d.	n.d.	n.d.
		cond					
		h × cond					
	Day 7	h	n.d.	n.d.	n.d.	n.d.	n.d.
		cond					
		h × cond					
<i>napus</i>	Day 1	h	n.d.	n.d.	n.d.	***	*
		cond				*	
		h × cond				*	
	Day 4	h	n.d.	n.d.	n.d.	n.d.	***
		cond					
		h × cond					
	Day 7	h	n.d.	n.d.		n.d.	
		cond					
		h × cond					
<i>carinata</i>	Day 1	h	n.d.	n.d.	n.d.		
		cond					
		h × cond			*		
	Day 4	h	n.d.	n.d.	n.d.		
		cond					
		h × cond					

Day 7	h	n.d.	n.d.	n.d.	***
	cond			*	**
	h × cond				**

Table S6: Quercetin tetraglycosides of different *Brassica* species in µg/g DW. Asterix indicate differences due to the UV-B treatment at the certain time point significant at \*  $p \leq 0.05$ , \*\*  $p \leq 0.01$ , \*\*\*  $p \leq 0.005$

Q: quercetin; soph: 15inapoyl15; glc: glucose; cou: coumaroyl; caf: caffeoyl; fer: feruloyl; hfer: hydroxyferuloyl; sin: 15inapoyl; h: hour; cond: condition

<i>Brassica</i>		Q-3-soph-7-diglc	Q-3-cou-soph-7-diglc	Q-3-caf-soph-7-diglc	Q-3-fer-soph-7-diglc	Q-3-hfer-soph-7-diglc	Q-3-sin-soph-7-diglc	Q-3-disin-triglc-7-glc
<i>rapa</i>	Day 1	h cond h × cond	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
	Day 4	h cond h × cond	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
	Day 7	h cond h × cond	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
<i>nigra</i>	Day 1	h cond h × cond	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
	Day 4	h cond h × cond	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
	Day 7	h cond h × cond	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
<i>oleracea</i>	Day 1	h cond h × cond	n.d.	n.d.	n.d.	n.d.	n.d.	*
	Day 4	h cond h × cond	n.d.	n.d.	n.d.	n.d.	n.d.	*

		cond					***
		h × cond					
	Day 7	h	n.d.	n.d.	n.d.	n.d.	
		cond					***
		h × cond					
<i>junccea</i>	Day 1	h	n.d.	n.d.	n.d.	n.d.	n.d.
		cond					
		h × cond					
	Day 4	h	n.d.	n.d.	n.d.	n.d.	n.d.
		cond					
		h × cond					
	Day 7	h	n.d.	n.d.	n.d.	n.d.	n.d.
		cond					
		h × cond					
<i>napus</i>	Day 1	h	n.d.	n.d.	n.d.	n.d.	n.d.
		cond					
		h × cond					
	Day 4	h	n.d.	n.d.	n.d.	n.d.	n.d.
		cond					
		h × cond					*
	Day 7	h	n.d.	n.d.	n.d.	n.d.	n.d.
		cond					
		h × cond					*
<i>carinata</i>	Day 1	h	n.d.	n.d.	n.d.	n.d.	n.d.
		cond					
		h × cond					
	Day 4	h	n.d.	n.d.	n.d.	n.d.	n.d.
		cond					
		h × cond					*

Day 7	h	n.d.	n.d.	n.d.	n.d.	n.d.
	cond					***
	h × cond					

Table S7: Quercetin tetraglycosides of different *Brassica* species in µg/g DW. Asterix indicate differences due to the UV-B treatment at the certain time point significant at \*  $p \leq 0.05$ , \*\*  $p \leq 0.01$ , \*\*\*  $p \leq 0.005$

Q: quercetin; soph: sinapoyl; glc: glucose; cou: coumaroyl; caf: caffeoyl; fer: feruloyl; hfer: hydroxyferuloyl; sin: sinapoyl; h: hour; cond: condition

<i>Brassica</i>		Chl	Cou-mal	Caf-mal	Fer-mal	Hfer-mal	Sin-mal	Disin-gent	Sin,fer-gent	Trisin-gent	Disin,fer-gent
<i>rapa</i>	Day 1	h									
		cond									
		h × cond									
	Day 4	h		*				***			*
		cond	*				**				
		h × cond									
	Day 7	h								n.d.	
		cond	*				*				
		h × cond									
<i>nigra</i>	Day 1	h	***	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	
		cond	*								
		h × cond									
	Day 4	h		n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	*
		cond									
		h × cond									
	Day 7	h		n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	*
		cond									*
		h × cond									

<i>oleracea</i>	Day 1	h	n.d.	n.d.	n.d.	n.d.	n.d.	*	*	*	*
		cond									
		h × cond									
	Day 4	h	n.d.	n.d.	n.d.	n.d.	n.d.	***	***	***	
		cond								*	***
		h × cond									***
	Day 7	h	n.d.	n.d.	n.d.	n.d.	n.d.				
		cond								*	
		h × cond									
<i>junccea</i>	Day 1	h		*				*			
		cond									
		h × cond						*			
	Day 4	h									
		cond	***		**		*				
		h × cond									
	Day 7	h	*	***		***					n.d.
		cond									
		h × cond									
<i>napus</i>	Day 1	h	*	*	***			*			
		cond									
		h × cond									
	Day 4	h	*		**	***	***				*
		cond	**	*	*		***				

			h × cond							
	Day	h								
	7									
		cond								
		h × cond								
<i>carinata</i>	Day	h		n.d.						
	1									
		cond								
		h × cond								
	Day	h		n.d.						
	4									
		cond								
		h × cond								
	Day	h		n.d.						
	7		***							
		cond	***							
		h × cond								