

Supplementary Materials

A Pilot Study of Rare Renal Amyloidosis Based on FFPE Proteomics

Shuang Meng ^{1,†}, Wenwen Xia ^{2,†}, Li Xia ¹, Li Zhou ¹, Jing Xu ³, Xiaoxia Pan ^{3,*} and Liyuan Meng ^{1,*}

- ¹ Department of Core Facility of Basic Medical Sciences, Shanghai Jiao Tong University School of Basic Medicine, Shanghai 200025, China; mengshuang92@shsmu.edu.cn (S.M.); lixia@shsmu.edu.cn (L.X.); zhouli_bit@shsmu.edu.cn (L.Z.)
 - ² Department of Pathology, Ninth People's Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai 200011, China; xiawenwenliuyedao@163.com
 - ³ Department of Nephrology, Ruijin Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai 200025, China; xj11564@rjh.com.cn
- * Correspondence: pxx10768@rjh.com.cn (X.P.); mengliyuan@shsmu.edu.cn (L.M.);
Tel.: +86-21-64370045 (X.P.); +86-21-63846590 (L.M.)
- † These authors contributed equally to this work.






Figure S1. Identified protein in control and AL-λ groups.



(a)



(b)
Figure S2. (a) . Go enrichment analysis bubble plot for molecular functions. **(b)** . Go enrichment analysis bubble plot for biological processes.




Figure S3: Light micrograph and Immunofluorescent staining for lambda light chain in kidney biopsy. a. light micrograph of glomerulus from a patient with minor glomerular disease showing no glomerular changes; b. mesangial and arteriolar amyloid deposition in patient with amyloidosis (white arrow); c. no lambda light chain deposition in patient with minor glomerular disease; d. positive mesangial and arteriolar lambda light chain deposition in patient with amyloidosis. (a. b. PAS 400'; c. d. light chain lambda 400')