

Rapid Screening of Lipase Inhibitors from *Ophiopogonis Radix* Using High-Performance Thin Layer Chromatography by Two Step Gradient Elution Combined with Bioautographic Method

Xue Hua ^{1,†}, Hui-Jie Hong ^{1,†}, Dai-Yan Zhang ¹, Qiao Liu ¹, Fong Leong ¹, Qi Yang ¹, Yuan-Jia Hu ^{1,*} and Xiao-Jia Chen ^{1,2,*}

¹ Institute of Chinese Medical Sciences, and State Key Laboratory of Quality Research in Chinese Medicine, University of Macau, Macao SAR, China; mb75819@connect.um.edu.mo (X.H.); yc17534@um.edu.mo (H.-J.H.); yc07528@um.edu.mo (D.-Y.Z.); yb87502@um.edu.mo (Q.L.); yb37501@um.edu.mo (F.L.); mc05849@um.edu.mo (Q.Y.)

² Zhuhai UM Science & Technology Research Institute, Zhuhai 519031, China

* Correspondence: YuanjiaHu@um.edu.mo (Y.-J.H.); XiaojiaChen@um.edu.mo (X.-J.C.); Tel.: +853-88228507 (Y.-J.H.); +853-88224915 (X.-J.C.)

[†] These authors contributed equally to this work.




Figure S1: HPTLC chromatograms (A and B) and HPTLC bioautograms (C) of mixed standards and *Ophiopogonis Radix* samples. Sample codes are the same as in Table 2. MS: mixed standards. 1. ophiopogonin D, 2. ophiopojaponin C, 3. ophiopogonin D', 4. L-borneol-7-*O*-[β -D-apiofuranosyl(1 \rightarrow 6)]- β -D-glucopyranoside, 5. ophiopogonin C', 6. β -sitosterol, and 7. methylphiopogonanone, Orlistat: positive control.