

Supplementary materials


Spectroscopic Studies of Quinobenzothiazine Derivative in Terms of the In Vitro Interaction with Selected Human Plasma Proteins. Part 2

Aleksandra Owczarzy ¹, Wojciech Rogóż ¹, Karolina Kulig ¹, Jadwiga Pożycka ¹, Andrzej Zięba ² and Małgorzata Maciążek-Jurczyk ^{1,*}

¹ Department of Physical Pharmacy, Faculty of Pharmaceutical Sciences in Sosnowiec, Medical University of Silesia, 40-055 Katowice, Poland

² Department of Organic Chemistry, Faculty of Pharmaceutical Sciences in Sosnowiec, Medical University of Silesia, 40-055 Katowice, Poland

* Correspondence: mmaciazek@sum.edu.pl; Tel.: +48-32-364-1582







Figure S1. Emission fluorescence spectra of human serum albumin (HSA) in the presence of Salt2 at Salt2:HSA 0:1 and 7:1 and Salt2:HSA 0:1 and 6:1 molar ratios at a) λ_{ex} 275 nm and b) λ_{ex} 295 nm, respectively.







Figure S2. Emission fluorescence spectra of α_1 -acid glycoprotein (AGP) in the presence of Salt2 at Salt2:HSA 0:1 and 8:1 and Salt2:HSA 0:1 and 7:1 molar ratios at a) λ_{ex} 275 nm and b) λ_{ex} 295 nm, respectively.







Figure S3. Emission fluorescence spectra of human gamma globulin (HGG) in the presence of Salt2 at Salt2:HSA 0:1 and 7:1 and Salt2:HSA 0:1 and 6:1 molar ratios at a) λ_{ex} 275 nm and b) λ_{ex} 295 nm, respectively.






Figure S4. Emission fluorescence spectra of control normal serum (CNS) in the presence of Salt2 at Salt2:HSA 0:1 and 7:1 and Salt2:HSA 0:1 and 6:1 molar ratios at a) λ_{ex} 275 nm and b) λ_{ex} 295 nm, respectively.