

A two-step synthesis of unprotected 3-aminoindoles via post functionalization with nitrostyrene


Nicolai A. Aksenov, Nikolai A. Arutiunov, Igor A. Kurenkov, Vladimir V. Malyuga, Dmitrii A. Aksenov, Daria S. Momotova, Anna M. Zatsepilina, Elizaveta A. Chukanova, Alexander V. Leontiev, and Alexander V. Aksenov


Department of Chemistry, North Caucasus Federal University, 1a Pushkin St., Stavropol 355009, Russian Federation


Supporting Information


NMR Spectral Charts.....	S2
¹ H and ¹³ C NMR spectral charts for indolinones 4ab , 4ad	S2
¹ H and ¹³ C NMR spectral charts for 3-aminoindole 5aa-ai , 19 , <i>N</i> -Ac 5aa and <i>N</i> -Ac 5ai	S7
¹ H and ¹³ C NMR spectral charts for 1-methyl-2-phenyl-1 <i>H</i> -indole J	S34
HRMS spectral chart for 4ab , 4ad , 5aa-ai , 19 , <i>N</i> -Ac 5aa and <i>N</i> -Ac 5ai	S36


^1H and ^{13}C NMR spectral charts for 2-(5-halide-3-oxo-2-phenylindolin-2-yl)-2-phenylacetonitrile **4ab**, **4ad**
















¹H and ¹³C NMR spectral charts for 2-aryl-1H-indol-3-amines **5aa-5ai**, **19**, *N*-Ac **5aa** and *N*-Ac **5ai**















































































^1H and ^{13}C NMR spectral charts for 1-methyl-2-phenyl-1*H*-indole **J**





HRMS spectral chart for **4ab**, **4ad**, **5aa-ai**, **19**, *N*-Ac **5aa** and *N*-Ac **5ai**

