

	Cytoplasmic	Nuclear	Ki-67	C 1	Ci	
	Expression	Expression	Expression	Grade	Stage	Age
Cytoplasmic Expression	Q = 1 p-value= 1	Q = 0.319 * p-value= 0.001 *	Q = 0.199 <i>p</i> -value= 0.644	Q = 0.218 * p-value= 0.024 *	Q = 0.072 <i>p</i> -value= 0.458	Q = 0.044 p-value= 0.654
Nuclear Expression		Q = 1 p-value= 1	Q = 0.309 * p-value=0.012*	Q = 0.215 * p-value= 0.026 *	Q = 0.064 <i>p</i> -value= 0.516	Q = -0.052 p-value= 0.592
Ki-67 expression			Q = 1 p-value= 1	Q = 0.196 p-value= 0.115	Q = 0.008 <i>p</i> -value= 0.949	Q = 0.117 <i>p</i> -value=0.35
Grade				Q = 1 p-value= 1	Q = 0.066 <i>p</i> -value= 0.501	Q = 0.113 <i>p</i> -value= 0.246
Stage					Q = 1 p -value= 1	Q = -0.120 p-value= 0.219
Age						Q = 1 p -value= 1

Supplemental Materials: Pearson's correlation of OSCC patients in the authors' own database. For the analysis, the Nuclear and Cytoplasmic expression were categorized as high/low on the basis of a threshold of 60% percentage expression. Staging was categorized in low (S1-S2) and high (S3-S4), Grading in high (G3) and low (G1-G2), and Age in old (>= 65 years' old) and non-old (< 65 years' old).



Supplemental Materials: Expression of Survivin in leukoplakia and healthy mucosa samples was mainly nuclear and confined in the basal third of the epithelium.



Supplemental Materials: Gene Expression Omnibus (GEO) analysis of methylation of the island cg25986496 in both tumor and paired adjacent normal samples.