

Table S2

Primers used for PCR analysis

<i>Actb:</i>	F: GCTACAGCTTCACCACCACA	R: GGTCTTTACGGATGTCAACG
<i>Cdc42bpa:</i>	F: GGATGAACTTGTGGTTCCTGA	R: ACCCATGTGTGCTATGTGGTT
<i>Epor:</i>	F: GGACCCTGTGACTATGGATGA	R: CCCCGAACGTGTAATCTGTTGA
<i>Fam132b:</i>	F: ATGCTGTTCGTCAAGCAGAGT	R: CCTTCAGCAGAACCTCAGATG
<i>Hamp:</i>	F: CTGAGCAGCACCACTATCTC	R: TGGCTCTAGGCTATGTTTTGC
<i>Id1:</i>	F: CGAGGTGGTACTTGGTCTGTC	R: CTGCAGGTCCCTGATGTAGTC
<i>Mcoln1:</i>	F: ATGAGCCCATGTGACAAGTTC	R: AGGTGTCATCAGACCCATCAG
<i>Mfn2:</i>	F: ACTGACCTGGACCACCAAAG	R: TGGCTCTGCTCTGAAGTGAA
<i>Tfr2:</i>	F: CCCTATCTGGTCCTGATCACC	R: CAAGGAACCGGAGAAACATGG
<i>Tfr1:</i>	F: AAAGTGGCTGCAGATGAAGAA	R: ATAGCCCAGGTAGCCACTCAT