



Article

Seventy-Two-Hour LRRK2 Kinase Activity Inhibition Increases Lysosomal GBA Expression in H4, a Human Neuroglioma Cell Line

Clara Ruz ^{1,2}, José Luis Alcantud ², Francisco Vives ^{1,2}, Francisco Arrebola ^{2,3}, John Hardy ⁴, Patrick A. Lewis ^{4,5}, Claudia Manzoni ⁶ and Raquel Duran ^{1,2*}

¹ Department of Physiology, Faculty of Medicine, Universidad de Granada, 18016, Granada, Spain; clararuz@ugr.es (C.R.); fvives@ugr.es (F.V.)

² Institute of Neurosciences “Federico Olóriz”, Centro de Investigación Biomédica (CIBM), Universidad de Granada, 18016 Granada, Spain; clararuz@ugr.es (CR); jlalcantud@ugr.es (JLA); fav@ugr.es (FA)

³ Department of Histology, Faculty of Medicine, Universidad de Granada, 18016 Granada, Spain; fav@ugr.es

⁴ Department of Neurodegenerative Disease, UCL Queen Square Institute of Neurology, WC1N 3BG London, UK; j.hardy@ucl.ac.uk (J.H.); plewis@rvc.ac.uk (P.A.L.)

⁵ Department of Comparative Biomedical Science, Royal Veterinary College, Royal College Street, NW1 0TU London, UK

⁶ Department of Pharmacology, UCL School of Pharmacy, WC1N 1AX London, UK; c.manzoni@ucl.ac.uk

* Correspondence: rduran@ugr.es

Supplementary Materials

Citation: Ruz, C.; Alcantud, J.L.; Vives, F.; Arrebola, F.; Hardy, J.; Lewis, P.A.; Manzoni, C.; Duran, R. Seventy-Two Hour LRRK2 Kinase Activity Inhibition Increases Lysosomal GBA Expression in H4, a Human Neuroglioma Cell Line. *Int. J. Mol. Sci.* **2022**, *23*, 6935. <https://doi.org/10.3390/ijms23136935>

Academic Editor: Alessandro Cannava

Received: 23 May 2022

Accepted: 20 June 2022

Published: 22 June 2022

Publisher’s Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).




Figure S1. GCase enzymatic activity of cell lysates treated with DMSO (as control), LRRK2in1, MLi2 and CBE after 24 h (black bars) and 72 h (grey bars) incubations.




Figure S2. Representative images (10X) from inverted phase-contrast microscope of H4 human neuroglioma cells after incubation with MLi-2 (600nM) or DMSO (0.15% v/v) as control for 24 and 72h.