

Table S1. Spearman rank correlation coefficients, determining the association between baseline biomarker levels and disease activity. Data is presented as ρ (p-value).

Biomarker	HBI ^a	SES-CD ^b	Active disease at baseline
<i>C1M</i>	0.12 (0.545)	0.30 (0.205)	-0.09 (0.638)
<i>C3M</i>	0.08 (0.690)	0.38 (0.097)	-0.02 (0.914)
<i>PRO-C3</i>	-0.14 (0.493)	-0.27 (0.255)	-0.34 (0.081)
<i>C3M/PRO-C3</i>	0.13 (0.508)	0.40 (0.080)	0.34 (0.081)
<i>C4M</i>	0.11 (0.570)	0.30 (0.198)	-0.06 (0.745)
<i>C4G</i>	0.35 (0.066)	-0.13 (0.586)	0.12 (0.538)
<i>PRO-C4</i>	0.01 (0.974)	0.21 (0.366)	0.05 (0.800)
<i>C4M/C4G</i>	-0.24 (0.224)	0.21 (0.373)	-0.18 (0.363)
<i>PRO-C4/C4M</i>	-0.12 (0.537)	-0.24 (0.308)	0.26 (0.174)
<i>PRO-C4/C4G</i>	-0.32 (0.103)	0.20 (0.405)	-0.14 (0.491)
<i>C6Ma3</i>	-0.26 (0.185)	0.05 (0.836)	-0.28 (0.151)
<i>CPa9-HNE</i>	-0.12 (0.544)	0.09 (0.701)	-0.05 (0.800)
<i>CPa9-HNE/C4G</i>	-0.30 (0.120)	0.07 (0.777)	-0.11 (0.587)

^aHBI scores available for 28 patients out of 32 (88%).

^bSES-CD scores available for 20 patients out of 32 (63%).