

Supporting Information

Gallium (III) complexes based on aminobisphenolate ligands: extremely high active ROP-initiators from well-known and easily accessible compounds

Badma N. Mankaev,^{1,2} Leyla. F. Hasanova,¹ Andrei V. Churakov,³ Mikhail P. Egorov,² and Sergey S. Karlov^{1,2,*}

¹*Chemistry Department, Moscow State University, 119991 Moscow, Russia;*

²*N.D. Zelinsky Institute of Organic Chemistry, Russian Academy of Science, 119991 Moscow, Russia;*

³*Institute of General and Inorganic Chemistry, Russian Academy of Science, Moscow, 119991, Russia.*

**Correspondence: s.s.karlov@chemistry.msu.ru*

Table of Contents

Figure S1 ¹ H NMR spectrum of PCL-200 (Table 1, entry 6) in CDCl ₃	S2
Figure S2 ¹ H NMR spectrum of 3 in C ₆ D ₆	S3
Figure S3 ¹³ C NMR spectrum of 3 in C ₆ D ₆	S3
Figure S4 ¹ H NMR spectrum of complex 4 in C ₆ D ₆	S4
Figure S5 ¹³ C NMR spectrum of complex 4 in C ₆ D ₆	S4
Figure S6 ¹ H NMR spectrum of complex 5 in C ₆ D ₆	S5
Figure S7 ¹³ C NMR spectrum of complex 5 in C ₆ D ₆	S5
Figure S8 ¹ H NMR spectrum of complex 6 in C ₆ D ₆	S6
Figure S9 ¹³ C NMR spectrum of complex 6 in C ₆ D ₆	S6




Figure S1 ¹H NMR spectrum of PCL-200 (Table 1, entry 6) in CDCl₃.




Figure S2 ¹H NMR spectrum of **3** in C₆D₆.




Figure S3 ¹³C NMR spectrum of **3** in C₆D₆.




Figure S4 ¹H NMR spectrum of complex 4 in C₆D₆.




Figure S5 ¹³C NMR spectrum of complex 4 in C₆D₆.




Figure S6 ¹H NMR spectrum of complex **5** in C₆D₆.




Figure S7 ¹³C NMR spectrum of complex **5** in C₆D₆.




Figure S8 ¹H NMR spectrum of complex 6 in C₆D₆.




Figure S9 ¹³C NMR spectrum of complex 6 in C₆D₆.