

Dehydrogenation of Betacyanins in Heated Betalain-Rich Extracts of Red Beet (*Beta vulgaris* L.)

Katarzyna Sutor-Świeży¹, Michał Antonik¹, Justyna Proszek¹, Boris Nemzer^{2,3}, Zbigniew Pietrzkowski⁴, Łukasz Popenda⁵, Tomasz Świergosz¹ and Sławomir Wybraniec^{1,*}

- ¹ Department C-1, Faculty of Chemical Engineering and Technology, Cracow University of Technology, ul. Warszawska 24, 31-155 Cracow, Poland; katarzyna.sutor@doktorant.pk.edu.pl (K.S.-Ś.); michal3antonik@gmail.com (M.A.); proszekjustyna@gmail.com (J.P.); tomasz.swiergosz@pk.edu.pl (T.Ś.);
- ² Research and Analytical Center, VDF FutureCeuticals, Inc., 2692 N. State Rt. 1-17, Momence, IL 60954, USA; bnemzer@futureceuticals.com
- ³ Food Science & Human Nutrition, University of Illinois at Urbana-Champaign, 260 Bevier Hall, 905 S. Goodwin Ave., Urbana, IL 61801, USA
- ⁴ Discovery Research Lab, VDF FutureCeuticals, Inc., 23 Peters Canyon Rd., Irvine, CA 92606, USA; zb@futureceuticals.com
- ⁵ NanoBioMedical Centre, Adam Mickiewicz University Poznan, ul. Wszechnicy Piastowskiej 3, 61-614 Poznan, Poland; lukasz.popenda@amu.edu.pl
- * Correspondence: slawomir.wybraniec@pk.edu.pl; Tel.: +48-12-628-3074; Fax: +48-12-628-2036




Figure S1. ¹H NMR spectrum of 15-decarboxy-betanin (D₂O, 298 K).




Figure S2. ¹³C NMR spectrum of 15-decarboxy-betanin (D₂O, 298 K).




Figure S3. ^1H NMR spectrum of 2,15-decarboxy-betanin (CD_3OD , 295 K).




Figure S4. ^{13}C NMR spectrum of 2,15-decarboxy-betanin (CD_3OD , 295 K).