

Table S1. Patients' description diagnosis with moderate-to-severe psoriasis treated with Anti-TNF and Anti-IL12/23.

Variable	Baseline (N=198)		
	N	%	Mean ± standard deviation
Sex			
Female	104	52.53	-
Male	94	47.47	-
Age at baseline	173	-	52.81 ± 14.17
<52 years	72	41.62	-
>52 years	101	58.38	-
BMI at baseline	196	-	27.09 (23.99-31.23)
Normal weight	68	35.42	-
Overweight	64	33.33	-
Obesity Type I	35	18.23	-
Obesity Type II	15	7.81	-
Obesity Type III	10	5.21	-
COMORBIDITIES			
Psoriatic Arthritis	89	44.95	-
Hypertension	65	32.83	-
Dyslipidemia	78	39.39	-
Other comorbidities	129	65.15	-
Age diagnosis PS	198	-	28 (18-42)
Family history PS	115	58.38	-
Type of PS			
Plaque	87	43.94	-
Pustular	4	2.02	-
Inverse	2	1.01	-
Guttate	6	3.03	-
Combination 2 types of PS	73	36.87	-
Combination 3 types of PS	25	12.63	-
Combination 4 types of PS	1	0.51	-
Location of lesions PS			
Trunk and lower and upper limbs	178	89.9	-
Scalp and face	127	64.14	-
Nails	82	41.41	-
Palmoplantar	28	14.14	-
Flexures	55	27.78	-
Genital	23	11.79	-
PS: Psoriasis.			
Qualitative variables are shown as numbers (percentage, %). Lilliefors (Kolmogorov-Smirnov) normality test. Quantitative variables with a normal distribution are shown as mean ± standard deviation. Quantitative variables with a non-normal distribution are shown as p ₅₀ (p ₂₅ - p ₇₅).			