

2-(Bicyclo[4.2.0]octa-1,3,5-trien-3-yl)-adamantan-2-ol

Konstantin S. Levchenko ^{1,*}, Konstantin A. Chudov ¹, Dmitri Yu. Demin ²,
Konstantin A. Lyssenko ^{3,4}, and Pavel S. Shmelin ²

¹ RTU MIREA—Russian Technological University, 78 Vernadsky Avenue, Moscow 119571, Russia; k4udov@gmail.com

² JSC «Technomash», 4 Ivana Franko Str., Moscow 121108, Russia; demindd@mail.ru (D.Y.D.); shmelin@cnititm.ru (P.S.S.)

³ Chemistry Department, M.V. Lomonosov Moscow State University, 1 Leninskie Gory Str., Building 3, Moscow 119991, Russia; kostya@xray.ineos.ac.ru

⁴ G.V. Plekhanov Russian University of Economics, 36 Stremyanny Per., Moscow 117997, Russia

* Correspondence: k.s.levchenko@gmail.com; Tel.: +7-985-188-6215

Supplementary Materials

The ¹H NMR data of **4** in accordance with previously obtained: [1].




Figure S1. ¹H NMR of compound **4**, 4-Bromobenzocyclobutene.

Adamantan-2-one **2** was synthesized as published earlier. The ^1H NMR data of compound **2**, in accordance with previously obtained [2].




Figure S2. ^1H NMR of compound **2**, Adamantan-2-one.




Figure S3. ¹H and ¹³C NMR of compound 5, 2-(bicyclo[4.2.0]octa-1,3,5-trien-3-yl)-adamantan-2-ol.

References

1. Roth M.; Ahles M.; Gawrisch C.; Schwalm Th.; Schmechel R.; Melzer C.; Seggern H.; Rehahn M. Rodlike Tetracene Derivatives. *Chem. Eur. J.* **2017**, *23*, 13445–13454, doi:10.1002/chem.201702382.
2. Geluk, H.W.; Keizer, V.G. Adamantanone. *Org. Synth.* **1973**, *53*, 8, doi:10.15227/orgsyn.053.0008.