

Short Note

# (10*E*,15*Z*)-12-(Dimethylsulfonio)-9,13-dihydroxyoctadeca-10,15-dienoate

Haruka Nishino <sup>1</sup>, Bo-Tao Zhang <sup>1</sup>, Hajime Uchida <sup>2</sup>, Michiya Kamio <sup>1</sup>, Hiroshi Nagai <sup>1,\*</sup> and Masayuki Satake <sup>3,\*</sup>

<sup>1</sup> Department of Ocean Sciences, Tokyo University of Marine Science and Technology, Tokyo 108-8477, Japan; m223051@edu.kaiyodai.ac.jp (H.N.); d202017@edu.kaiyodai.ac.jp (B.-T.Z.); mkamio@kaiyodai.ac.jp (M.K.)

<sup>2</sup> Fisheries Technology Institute, Japan Fisheries Research and Education Agency, Yokohama 236-8648, Japan; uchida\_hajime03@fra.go.jp

<sup>3</sup> Department of Chemistry, The University of Tokyo, Tokyo 113-0033, Japan


\* Correspondence: nagai@kaiyodai.ac.jp (H.N.); satakem@g.ecc.u-tokyo.ac.jp (M.S.)

**Abstract:** A novel oxylipin, okeanoate (**1**), was isolated from the Okinawan cyanobacterium *Okeania hirsuta*. The structure of **1** was elucidated based on spectroscopic data including 1D and 2D NMR, as well as high-resolution mass spectrometry. This is the first oxylipin with a dimethylsulfonium moiety in the middle of the hydrocarbon chain.

**Keywords:** cyanobacteria; oxylipin; nuclear magnetic resonance; marine natural product; dimethylsulfonium

## 1. Introduction

Oxylipins are a group of oxygenated natural products formed from fatty acids [1]. It has been reported that many oxylipins perform physiologically important activities in many organisms [2]. Cyanobacteria are known to produce numerous bioactive substances and are recognized as valuable biochemical resources [3,4]. A few oxylipins have been reported as natural products from cyanobacteria so far [5–10]. Our research group has recently reported two known isolated oxylipins, malyncic acid, 15,16-dihydro malyncic acid, and a new oxylipin, okeanic acid–A [10], from the marine cyanobacterium *Okeania hirsuta*. In this study, further research on the cyanobacterium *O. hirsuta* extracts led to the isolation of a novel oxylipin, (10*E*,15*Z*)-12-(dimethylsulfonio)-9,13-dihydroxyoctadeca-10,15-dienoate (**1**, Figure 1). Only a few oxylipins with a dimethylsulfonium group at the end of the hydrocarbon chain have been reported from nature [11,12]. To the best of our knowledge, there are no reports of oxylipins having a dimethylsulfonium group in the middle of the hydrocarbon chain so far. This is the first finding of an oxylipin with a dimethylsulfonium moiety in the middle of the hydrocarbon chain from nature. The compound (**1**) was designated as okeanoate.



**1:** okeanoate

**Figure 1.** Structure of okeanoate (**1**).

## 2. Results and Discussion

The cyanobacterium *Okeania hirsuta*, identified using 16S rRNA sequence analysis, was collected from Kuba Beach, Okinawa, Japan, in 2010. A frozen sample of *O. hirsuta* was



**Citation:** Nishino, H.; Zhang, B.-T.; Uchida, H.; Kamio, M.; Nagai, H.; Satake, M. (10*E*,15*Z*)-12-(Dimethylsulfonio)-9,13-dihydroxyoctadeca-10,15-dienoate. *Molbank* **2024**, *2024*, M1784. <https://doi.org/10.3390/M1784>

Academic Editor: Oleg A. Rakitin

Received: 19 January 2024

Revised: 26 February 2024

Accepted: 28 February 2024

Published: 3 March 2024



**Copyright:** © 2024 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

extracted with methanol (MeOH). The compound (**1**) was purified using reversed-phase column chromatography. From 9.7 kg of algal cells, 1.1 mg of **1** was isolated.

Compound (**1**) was isolated as a colorless amorphous material with no UV absorption above 210 nm. The molecular formula of **1** was  $C_{20}H_{36}O_4S$  ( $m/z$  373.2408  $[M + H]^+$ , calcd. for  $C_{20}H_{37}O_4S$   $m/z$  373.2407) via high-resolution electron spray ionization mass spectrometry (HR-ESI-MS), suggesting the presence of a sulfur atom in the molecule.  $^1H$ ,  $^{13}C$  NMR, and HSQC spectra revealed the presence of a triplet methyl group ( $\delta_H$  0.97) and two singlet methyls ( $\delta_H$  2.79 and 2.80), nine aliphatic methylenes ( $\delta_C$  39.0, 27.6, 30.5, 30.4, 30.6, 26.3, 38.1, 34.4, 21.9), three methines ( $\delta_H$  4.23, 4.12, 4.16), four olefin signals ( $\delta_C$  149.9, 115.7, 123.7, 136.3), and a carbonyl ( $\delta_C$  180.1) (Table 1). Considered from MS and NMR analysis, the unsaturation of **1** was three, which accounted for a carbonyl and two double bonds. Thus, **1** was a C18:2 fatty acid derivative that contained a sulfonium group, two singlet methyls bonding a hetero atom, two hydroxy groups, and a carboxylate in the molecule. Similar to malyngic acid [5] and okeanic acid-A [10], **1** was deduced to be an  $\omega$ -3 fatty acid derivative by  $^1H$ - $^1H$  COSY correlation, from H<sub>3</sub>-18 ( $\delta_H$  0.97) to H-15 ( $\delta_H$  5.31). Another proton connectivity around the hydroxy groups and the double bonds H-15-H<sub>2</sub>-14-H-13-H<sub>2</sub>-12-H-11-H-10-H-9-H<sub>2</sub>-8 was also determined based on the detailed analyses of  $^1H$ - $^1H$  COSY and TOCSY. The rest of the six methylenes were assigned into a linear alkyl chain from H<sub>2</sub>-2 to H<sub>2</sub>-7. An HMBC correlation from H<sub>2</sub>-2 ( $\delta_H$  2.14) to C-1 ( $\delta_C$  180.1) was observed. Therefore, the skeletal structure of **1** was elucidated. The geometry of  $\Delta^{10}$  was deduced to be *E* from  $^3J_{H-10,H-11}$  (15.5 Hz), and the  $^{13}C$  chemical shift of C-17 at  $\delta_C$  21.7 suggested that the configuration of C-15=C-16 was *Z* [10]. Two singlet methyls observed at  $\delta_H$  2.80 and 2.79 and  $\delta_C$  23.1 and 23.9, respectively, indicated a dimethylsulfonium group in the molecule [11,12]. The  $^{13}C$  chemical shift ( $\delta_C$  64.7) at C-12 and the HMBC correlation from the singlet methyls to C-12 indicated that the dimethylsulfonium group resided on C-12 (Figure 2). The presence of two hydroxy groups was deduced from the molecular formula obtained from HR-MS results and the  $^1H$  and  $^{13}C$  NMR chemical shifts of CH-9 ( $\delta_H$  4.23,  $\delta_C$  72.4) and CH-13 ( $\delta_H$  4.16,  $\delta_C$  72.0). The position of the hydroxy groups was deduced to be at C-9 ( $\delta_C$  72.4) and C-13 ( $\delta_C$  72.0). These MS and NMR analyses allowed us to determine that the structure of okeanoate (**1**) was (10*E*,15*Z*)-12-(dimethylsulfonio)-9,13-dihydroxyoctadeca-10,15-dienoate. Okeanoate (**1**) was unstable in methanol at 5 °C under dark condition. Within a week, more than half of the isolated **1** changed to other unknown compounds under these conditions. Therefore, biological activity tests could not be performed for **1**.

**Table 1.**  $^1H$  (600 MHz) and  $^{13}C$  (150 MHz) NMR data of compound **1** in methanol-*d*<sub>4</sub>.

Position	$\delta_C$ (ppm)	$\delta_H$ (ppm), Multiplicity (J in Hz)
1	180.1	–
2	39.0	2.14 t (7.4, 2H)
3	27.6	1.57–1.62 m (2H)
4	30.5	1.32–1.37 m (2H)
5	30.4	1.32–1.37 m (2H)
6	30.6	1.32–1.37 m (2H)
7	26.3	1.32–1.37 m (2H)
8	38.1	1.57–1.62 m (2H)
9	72.4	4.23 ddd (6.3, 6.1, 5.7, 1H)

Table 1. Cont.

Position	$\delta_C$ (ppm)	$\delta_H$ (ppm), Multiplicity (J in Hz)
10	149.9	6.13 dd (15.5, 5.5, 1H)
11	115.7	5.87 dd (15.5, 10.4, 1H)
12	64.7	4.12 dd (10.5, 2.6, 1H)
13	72.0	4.16 ddd (14.2, 7.0, 2.6, 1H)
14	34.3	2.34 dt (4.3, 6.6, 1H), 2.25 dt (14.7, 7.4, 1H)
15	123.7	5.31 dd (8.4, 7.5, 1H)
16	136.3	5.55 dd (8.6, 7.6, 1H)
17	21.9	2.04–2.08 m (2H)
18	14.5	0.97 t (7.5, 3H)
19	23.9	2.79 s (3H)
20	23.1	2.80 s (3H)




Figure 2. Key COSY and TOCSY (bold line) and HMBC (arrow, H to C) correlations for okeanoate (1).

### 3. Material and Methods

#### 3.1. Collection of Cyanobacteria

The cyanobacterium *Okeania hirsuta* was collected at Kuba Beach, Nakagusuku, in Okinawa Prefecture, Japan, in 2010. *O. hirsuta* was the dominant cyanobacterial species in all samples. The identification of *O. hirsuta* was performed based on 16S rRNA sequence analysis [13]. A voucher specimen (20100713-a) was deposited at the Tokyo University of Marine Science and Technology.

#### 3.2. Isolation and Structural Characterization of 1

The frozen *O. hirsuta* sample (wet weight: 9.7 kg) was extracted with MeOH thrice. After the elimination of MeOH under reduced pressure, the extracts were partitioned between ethyl acetate and H<sub>2</sub>O. The H<sub>2</sub>O layer was extracted with 1-butanol (BuOH). After the evaporation of 1-BuOH, the extract was fractionated by MPLC, using ODS gel (Cosmosil 75C18-OPN, Nacalai Tesque Inc., Kyoto, Japan) with stepwise elution with 50%, 70%, 90%, and 100% MeOH [size of the column, 20 × 300 mm; each eluant volume, 200 mL]. The 70% MeOH eluate was subjected to an ODS column (Cosmosil 5C18-AR-II, 10 × 250 mm, Nacalai Tesque Inc., Kyoto, Japan) with 70% MeOH for 10 min, then a gradient elution from 70% to 100% MeOH for 20 min [flow rate, 3.0 mL/min; detection at 210 nm]. The fraction eluted from 5 to 8 min was combined. Final purification was conducted via HPLC using the reversed-phase column (Cosmosil 5C18-AR-II, 4.6 × 250 mm, Nacalai Tesque Inc., Kyoto, Japan) with gradient elution from 45% to 50% MeOH for 0–5 min, then 50% MeOH for 5–15 min [flow rate, 1.0 mL/min; detection at 210 nm]. Okeanoate (1, 1.1 mg) was isolated at a retention time of 7.0 min.

HR ESI-MS was measured with an micrOTOF-QII (Bruker, MA, USA) Quadrupole time-of-flight mass spectrometer (QTOFMS) with high-performance liquid chromatography of the UltiMate 3000 Series (Thermo Fisher Scientific, Billerica, MA, USA). NMR spectra were measured in methanol- $d_4$  at 300 K using a Bruker AVANCE III 600 spectrometer (Bruker Biospin AG, Fällanden, Switzerland).

#### 4. Conclusions

Okeanoate (**1**), a novel oxylipin, was isolated from the cyanobacterium *Okeania hirsuta* collected in Okinawa, Japan. In this study, the stereochemistry of the two double bonds could be determined, but only the planar structure could be determined for the other three chiral carbon atoms. To the best of our knowledge, okeanoate (**1**) is the first oxylipin with a dimethylsulfonium moiety in the middle of the hydrocarbon chain from nature. The biosynthetic pathway to this chemically unique compound is of interest.

**Supplementary Materials:** Figure S1.  $^1\text{H-NMR}$  (600 MHz, methanol- $d_4$ ) spectrum of okeanoate (**1**). Figure S2.  $^{13}\text{C-NMR}$  (150 MHz, methanol- $d_4$ ) spectrum of okeanoate (**1**). Figure S3. HSQC spectrum (600 MHz, methanol- $d_4$ ) of okeanoate (**1**). Figure S4. HMBC spectrum (600 MHz, methanol- $d_4$ ) of okeanoate (**1**). Figure S5.  $^1\text{H-}^1\text{H COSY}$  spectrum (600 MHz, methanol- $d_4$ ) of okeanoate (**1**). Figure S6.  $^1\text{H-}^1\text{H TOCSY}$  spectrum (600 MHz, methanol- $d_4$ ) of okeanoate (**1**). Figure S7. HR-ESI-MS spectrum of okeanoate (**1**). Figure S8. The spectral area between 4.00 and 4.30 ppm of  $^1\text{H-NMR}$  (600 MHz, methanol- $d_4$ ) spectrum of okeanoate (**1**). Figure S9. The spectral area between 5.20 and 6.25 ppm of  $^1\text{H-NMR}$  (600 MHz, methanol- $d_4$ ) spectrum of okeanoate (**1**).

**Author Contributions:** Conceptualization, H.N. (Hiroshi Nagai) and M.S.; isolation, H.N. (Haruka Nishino); structure analysis, H.N. (Haruka Nishino), B.-T.Z., H.N. (Hiroshi Nagai), and M.S.; analysis, H.U.; writing—original draft preparation, M.S., H.N. (Hiroshi Nagai), H.N. (Haruka Nishino); writing—review and editing, M.K. All authors have read and agreed to the published version of the manuscript.

**Funding:** This work was funded by Japan Society for the Promotion of Science: 22K05817 (Hiroshi Nagai); Japan Society for the Promotion of Science: 23H01962 (M.S.); Japan Science and Technology Agency: JPMJSP2147 (B.Z.).

**Data Availability Statement:** The spectroscopic data presented in this study are available as Supplementary Materials.

**Conflicts of Interest:** The authors declare no conflicts of interest.

#### References

1. Wasternack, C.; Feussner, I. The oxylipin pathways: Biochemistry and function. *Annu. Rev. Plant Biol.* **2018**, *69*, 363–386. [[CrossRef](#)] [[PubMed](#)]
2. Howe, G.A.; Schillmiller, A.L. Oxylipin metabolism in response to stress. *Curr. Opin. Plant Biol.* **2002**, *5*, 230–236. [[CrossRef](#)] [[PubMed](#)]
3. Nunnery, J.K.; Mevers, E.; Gerwick, W.H. Biologically active secondary metabolites from marine cyanobacteria. *Curr. Opin. Biotechnol.* **2010**, *21*, 787–793. [[CrossRef](#)] [[PubMed](#)]
4. Shah, S.A.A.; Akhter, N.; Auckloo, B.N.; Khan, I.; Lu, Y.; Wang, K.; Wu, B.; Guo, Y.W. Structural diversity, biological properties and applications of natural products from cyanobacteria. A review. *Mar. Drugs* **2017**, *15*, 354. [[CrossRef](#)] [[PubMed](#)]
5. Cardellina, J.H.; Moore, R.E. Malyngic acid, a new fatty acid from *Lyngbya majuscula*. *Tetrahedron* **1980**, *36*, 993–996. [[CrossRef](#)]
6. Herz, W.; Kulanthaivel, P. Trihydroxy- $\text{C}_{18}$ -acids and a labdane from *Rudbeckia fulgida*. *Phytochemistry* **1985**, *24*, 89–91. [[CrossRef](#)]
7. Lang, I.; Feussner, I. Oxylipin formation in *Nostoc punctiforme* (PCC73102). *Phytochemistry* **2007**, *68*, 1120–1127. [[CrossRef](#)] [[PubMed](#)]
8. Oku, N.; Matsumoto, M.; Yonejima, K.; Tansei, K.; Igarashi, Y. Sacrolide A, a new antimicrobial and cytotoxic oxylipin macrolide from the edible cyanobacterium *Aphanothece sacrum*. *Beilstein J. Org. Chem.* **2014**, *10*, 1808–1816. [[CrossRef](#)] [[PubMed](#)]
9. Oku, N.; Hana, S.; Matsumoto, M.; Yonejima, K.; Tansei, K.; Isogai, Y.; Igarashi, Y. Two new sacrolide-class oxylipins from the edible cyanobacterium *Aphanothece sacrum*. *J. Antibiot.* **2017**, *70*, 708–709. [[CrossRef](#)] [[PubMed](#)]
10. Nishino, H.; Kanda, N.; Zhang, B.T.; Kamio, M.; Uchida, H.; Sugahara, K.; Nagai, H.; Satake, M. Okeanic acid-A, a trihydroxy fatty acid from the Okinawan cyanobacterium *Okeania hirsuta*. *Nat. Prod. Res.* **2024**. [[CrossRef](#)] [[PubMed](#)]
11. Sciuto, S.; Piattelli, M.; Chillemi, R. (–)-(S)-Dimethylsulfonio-2-methoxybutylate from the red algae *Rytidiphloea tinctoria*. *Phytochemistry* **1982**, *21*, 227–228. [[CrossRef](#)]

12. Nakamura, H.; Fujimaki, K.; Sampei, O.; Murai, A. Gonyol: Methionine-induced sulfonium accumulation in a dinoflagellate *Gonyaulax polyedra*. *Tetrahedron Lett.* **1993**, *34*, 8481–8484. [[CrossRef](#)]
13. Kanda, N.; Zhang, B.T.; Shinjo, A.; Kamiya, M.; Nagai, H.; Uchida, H.; Araki, Y.; Nishikawa, T.; Satake, M. 7-Epi-30-Methylosclatotoxin D from an Okinawan cyanobacterium *Okeania hirsuta*. *Nat. Prod. Commun.* **2023**, *18*, 1–5. [[CrossRef](#)]

**Disclaimer/Publisher’s Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.