

Supplementary Materials: Estimation of the Botanical Composition of Clover-Grass Leys from RGB Images Using Data Simulation and Fully Convolutional Neural Networks

Søren Skovsen ^{1,*}, Mads Dyrmann ¹, Anders Krogh Mortensen ², Kim Arild Steen ³, Ole Green ³, Jørgen Eriksen ⁴, René Gislum ², Rasmus Nyholm Jørgensen ¹ and Henrik Karstoft ¹

¹ Department of Engineering, Aarhus University, Finlandsgade 22, 8200 Aarhus N, Denmark; madsdyrmann@eng.au.dk (M.D.); rnj@eng.au.dk (R.N.J.); hka@eng.au.dk (H.K.)

² Department of Agroecology, Aarhus University, Forsøgsvej 1, 4200 Slagelse, Denmark; anmo@agro.au.dk (A.K.M.); rg@agro.au.dk (R.G.)

³ Agro Intelligence ApS, Agro Food Park 13, 8200 Aarhus N, Denmark; kas@agointelli.com (K.A.S.); olg@agointelli.com (O.G.)

⁴ Department of Agroecology, Aarhus University, Blichers Allé 20, 8830 Tjele, Denmark; jorgen.eriksen@agro.au.dk

* Correspondence: ssk@eng.au.dk

Received: 31 October 2017; Accepted: 12 December 2017; Published: 17 December 2017




Figure 1. Example of the image analysis on a real image. Each element in the image is analyzed and classified as either grass (blue), clover (red), weeds (yellow) or unidentified (black overlay). **(a)** Example input image. **(b)** Automatically analyzed image.