

Table S4. DKK4 deregulation reported in different tumors and its therapeutic potential.

Tumor	Reported function	Selected evidence for therapeutic potential	References
Ovarian cancer	Oncogene	Prognostic factor	1,2
		<i>In vitro</i> migration/invasion	1
Pancreatic cancer	Oncogene	Diagnostic factor	3–5
		Diagnostic factor	6,7
		Prognostic factor	7
Renal cell carcinoma	Oncogene	<i>In vitro</i> proliferation	6,7
		<i>In vitro</i> migration/invasion	6,7
		<i>In vivo</i> tumor growth	6,7
Hepatocellular carcinoma	Tumor suppressor	Diagnostic factor	8
		Prognostic factor	9
		<i>In vitro</i> proliferation	8–12
		<i>In vitro</i> migration/invasion	8,9
		<i>In vivo</i> tumor growth	8–12
Colorectal cancer	Oncogene	Diagnostic factor	13
		Prognostic factor	14
		<i>In vitro</i> proliferation	15
		<i>In vitro</i> migration/invasion	15,16
		<i>In vivo</i> tumor growth	17
Gastrointestinal stromal tumor	Tumor suppressor	<i>In vitro</i> proliferation	18
		Diagnostic factor	20
		Prognostic factor	19
Lung cancer	Oncogene	<i>In vitro</i> proliferation	21
		<i>In vitro</i> migration/invasion	21
		Prognostic factor	22

REFERENCES SUPPLEMENTARY TABLE 4

1. Wang, S., Wei, H. & Zhang, S. Dickkopf-4 is frequently overexpressed in epithelial ovarian carcinoma and promotes tumor invasion. *BMC Cancer* **17**, 455 (2017).
2. Wang, Y., Lei, L., Chi, Y. G., Liu, L. B. & Yang, B. P. A comprehensive understanding of ovarian carcinoma survival prognosis by novel biomarkers. *Eur. Rev. Med. Pharmacol. Sci.* (2019) doi:10.26355/eurrev_201910_19136.
3. Ouyang, Y., Pan, J., Tai, Q., Ju, J. & Wang, H. Transcriptomic changes associated with DKK4 overexpression in pancreatic cancer cells detected by RNA-Seq. *Tumor Biol.* **37**, 10827–10838 (2016).
4. Park, M. *et al.* Distinct Protein Expression Profiles of Solid-Pseudopapillary Neoplasms of the Pancreas. *J. Proteome Res.* (2015) doi:10.1021/acs.jproteome.5b00423.
5. Zhu, Y. *et al.* Proteomic analysis of solid pseudopapillary tumor of the pancreas reveals dysfunction of the endoplasmic reticulum protein processing pathway. *Mol. Cell. Proteomics* (2014) doi:10.1074/mcp.M114.038786.
6. Hirata, H. *et al.* DICKKOPF-4 activates the noncanonical c-Jun-NH₂ kinase signaling pathway while inhibiting the Wnt-canonical pathway in human renal cell carcinoma. *Cancer* **117**, 1649–1660 (2011).
7. Zhai, W. *et al.* High expression of the secreted protein dickkopf homolog 4: Roles in invasion and metastasis of renal cell carcinoma and its association with von Hippel-Lindau gene. *Int. J. Mol. Med.*

- 33**, 1319–1326 (2014).
- 8. Fatima, S. *et al.* Dickkopf 4 (DKK4) acts on Wnt/β-catenin pathway by influencing β-catenin in hepatocellular carcinoma. *Oncogene* **31**, 4233–4244 (2012).
 - 9. Liao, C. H. *et al.* Dickkopf 4 positively regulated by the thyroid hormone receptor suppresses cell invasion in human hepatoma cells. *Hepatology* **55**, 910–920 (2012).
 - 10. Chi, H. C. *et al.* Thyroid hormone receptor inhibits hepatoma cell migration through transcriptional activation of Dickkopf 4. *Biochem. Biophys. Res. Commun.* **439**, 60–65 (2013).
 - 11. Chouhan, S. *et al.* Glucose induced activation of canonical Wnt signaling pathway in hepatocellular carcinoma is regulated by DKK4. *Sci. Rep.* **6**, 27558 (2016).
 - 12. Suresh, S. *et al.* SRC-2-mediated coactivation of anti-tumorigenic target genes suppresses MYC-induced liver cancer. *PLoS Genet.* (2017) doi:10.1371/journal.pgen.1006650.
 - 13. Matsui, A. *et al.* DICKKOPF-4 and -2 genes are upregulated in human colorectal cancer. *Cancer Sci.* **100**, 1923–1930 (2009).
 - 14. Tsukui, Y. *et al.* Dickkopf-4 gene expression is associated with differentiation and lymph node metastasis in colorectal cancer. *JGH Open* (2019) doi:10.1002/jgh3.12177.
 - 15. He, S., Shen, J., Hu, N., Xu, X. & Li, J. DKK4 enhances resistance to chemotherapeutics 5-Fu and YN968D1 in colorectal cancer cells. *Oncol. Lett.* **13**, 587–592 (2017).
 - 16. Pendás-Franco, N. *et al.* DICKKOPF-4 is induced by TCF/β-catenin and upregulated in human colon cancer, promotes tumour cell invasion and angiogenesis and is repressed by 1α,25-dihydroxyvitamin D3. *Oncogene* **27**, 4467–4477 (2008).
 - 17. Chen, H. J. *et al.* Comprehensive models of human primary and metastatic colorectal tumors in immunodeficient and immunocompetent mice by chemokine targeting. *Nat. Biotechnol.* (2015) doi:10.1038/nbt.3239.
 - 18. Baehs, S. *et al.* Dickkopf-4 is frequently down-regulated and inhibits growth of colorectal cancer cells. *Cancer Lett.* **276**, 152–159 (2009).
 - 19. Wang, M. *et al.* Aberrant accumulation of Dickkopf 4 promotes tumor progression via forming the immune suppressive microenvironment in gastrointestinal stromal tumor. *Cancer Med.* (2019) doi:10.1002/cam4.2437.
 - 20. Zeng, S. *et al.* Wnt/β-catenin signaling contributes to tumor malignancy and is targetable in gastrointestinal stromal tumor. *Mol. Cancer Ther.* (2017) doi:10.1158/1535-7163.MCT-17-0139.
 - 21. Yang, X., Liu, Y., Li, W., Li, A. & Sun, Q. DKK4-knockdown enhances chemosensitivity of A549/DTX cells to docetaxel. *Acta Biochim. Biophys. Sin. (Shanghai)*. (2017) doi:10.1093/abbs/gmx086.
 - 22. Bahl, C., Singh, N., Behera, D. & Sharma, S. Genetic Variants in the Wingless Antagonist Genes (sFRP, DKK, and Axin2) Predict the Overall Survival and Prognosis of North Indian Lung Cancer Patients Treated with Platinum-Based Doublet Chemotherapy. *Cancer Biother. Radiopharm.* (2018) doi:10.1089/cbr.2018.2491.