


Supplementary Table S1. The primers used for quantitative RT–PCR

Gene symbols	Sequence	
<i>MYC</i>	Forward	5'- TCCAGCTTGTACCTGCAGGATCTGA-3'
	Reverse	5'- CCTCCAGCAGAAGGTGATCCAGACT-3'
<i>FASLG</i>	Forward	5'- TTGGAGAAGCAAATAGGCCACC-3'
	Reverse	5'- AAGATTGAACACTGCCCCCA-3'
<i>PEG3</i>	Forward	5'- CTCACAACACAATCCAGGAC-3'
	Reverse	5'- TAGACCTCGACTGGTGCTTG-3'
<i>FADD</i>	Forward	5'- GCCATGGACCCGTTCTGG-3'
	Reverse	5'- CTCGATGCTGTCGATCTTG-3'
<i>MCL1</i>	Forward	5'- CTTCGGAACTGGACATCAA-3'
	Reverse	5'- GAAGAACTCCACAAACCCATC-3'
<i>IL12A</i>	Forward	5'- CACAAAGGAGGCGAGGTTCT-3'
	Reverse	5'- GGGTCAGGTTTGATGATGTCC-3'
<i>CXCL10</i>	Forward	5'- AGCAGTTAGCAAGGAAAGGTCT-3'
	Reverse	5'- GGAGGATGGCAGTGGAAGTC-3'
<i>HGF</i>	Forward	5'- CAGCTTTTTGCCTTCGAGCTA-3'
	Reverse	5'- TCTCGTAGGTCCTTGCACTT-3'
<i>AKT1</i>	Forward	5'- AAGACGGGAGCAGGCGG-3'
	Reverse	5'- GGTCTTGATGTACTCCCCTCG-3'
<i>ACTB</i>	Forward	5'- GCTGTGCTATCCCTGTACGC-3'
	Reverse	5'- TGCCTCAGGGCAGCGGAACC-3'



Supplementary Figure S1