

Supplementary Table S1. Crude prevalence and incidence of diagnosed knee osteoarthritis in Beijing, China, 2008–2017.

	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017
Prevalence (%; 95% CI)										
Overall	1.384 (1.377,1.392)	1.767 (1.759,1.775)	2.649 (2.640,2.658)	3.477 (3.468,3.486)	3.883 (3.874,3.892)	3.853 (3.844,3.861)	4.625 (4.616,4.635)	4.963 (4.954,4.972)	4.497 (4.449,4.506)	4.639 (4.631,4.648)
Men	0.796 (0.788,0.803)	1.034 (1.026,1.042)	1.605 (1.596,1.614)	2.139 (2.129,2.148)	2.405 (2.396,2.415)	2.440 (2.431,2.450)	3.066 (3.055,3.076)	3.364 (3.354,3.375)	3.071 (3.061,3.081)	3.213 (3.204,3.223)
Women	2.137 (2.124,2.150)	2.702 (2.688,2.716)	3.966 (3.951,3.982)	5.143 (5.126,5.159)	5.702 (5.686,5.719)	5.589 (5.573,5.637)	6.543 (6.526,6.560)	6.930 (6.914,6.947)	6.258 (6.243,6.274)	6.407 (6.392,6.422)
Incidence (per 1000 person-years; 95% CI)										
Overall	12.62 (12.55,12.69)	15.75 (15.67,15.82)	16.34 (16.27,16.41)	21.48 (21.40,21.56)	21.50 (21.43,21.57)	18.45 (18.38,18.51)	22.69 (22.62,22.76)	18.83 (18.76,19.21)	15.37 (15.32,15.43)	14.41 (14.37,14.46)
Men	7.31 (7.24,7.38)	9.63 (9.55,9.70)	10.43 (10.35,10.50)	14.00 (13.92,14.08)	14.38 (14.30,14.45)	13.06 (12.98,13.13)	17.00 (16.92,17.08)	14.49 (14.41,14.56)	11.88 (11.81,11.95)	11.29 (11.23,11.36)
Women	19.41 (19.28,19.53)	23.65 (23.51,23.79)	23.99 (23.86,24.11)	31.13 (31.00,31.27)	30.72 (30.59,30.84)	25.49 (25.37,25.61)	30.19 (30.07,30.31)	24.60 (24.49,24.71)	20.07 (19.97,20.17)	18.64 (18.55,18.73)