

[illegible]

No	1.00 (Reference)				1.00 (Reference)		1.00 (Reference)	
Yes	0.89 (0.63-1.25)	0.494			1.20 (0.60-2.11)	0.719	0.56 (0.22-1.41)	0.217
Sleepiness during day								
No	1.00 (Reference)		1.00 (Reference)		1.00 (Reference)		1.00 (Reference)	
Yes	2.28 (1.63-3.18)	<0.001	1.60 (1.06-2.42)	0.026	1.64 (0.92-2.93)	0.097	2.36 (0.98-5.70)	0.056
Confusional state								
No	1.00 (Reference)				1.00 (Reference)		1.00 (Reference)	
Yes	2.55 (1.51-4.33)	0.005			1.75 (0.77-3.99)	0.180	3.86 (0.67-22.4)	0.131
Involuntary tremors								
No	1.00 (Reference)				1.00 (Reference)		1.00 (Reference)	
Yes	2.97 (1.64-5.36)	<0.001			1.99 (0.80-4.84)	0.139	21.4 (2.57-178.0)	<0.001
Tingling in the limbs/face								
No	1.00 (Reference)				1.00 (Reference)		1.00 (Reference)	
Yes	1.15 (0.74-1.18)	0.5280			1.17 (0.57-2.43)	0.671	1.92 (0.73-5.62)	0.185
Number of overall symptoms²								
1	1.00 (Reference)				1.00 (Reference)		1.00 (Reference)	
2	2.82 (1.58-5.04)	<0.001			1.52 (0.53-4.34)	0.435	2.00 (0.39-10.2)	0.403
3	7.35 (4.06-13.3)	<0.001			3.30 (1.26-8.64)	0.015	7.00 (1.45-33.7)	0.015
≥4	15.0 (8.83-25.5)	<0.001			5.44 (1.89-15.6)	0.002	15.2 (3.68-63.1)	<0.001

OR =Odds ratio represent the probability of calling the GP over probability of not calling the GP

¹ In the table are reported only the statistically significant variables implemented in the multivariate model

² Not evaluated in multivariate analyses due to multicollinearity issues