

Supplementary Information

Increased Fibroblast Metabolic Activity of Collagen Scaffolds via the Addition of Propolis Nanoparticles

Jeimmy González-Masís ¹, Jorge M. Cubero-Sesin ¹, Yendry R. Corrales-Ureña ²,
 Sara González-Camacho ³, Nohelia Mora-Ugalde ⁴, Mónica Baizán-Rojas ⁴, Randall Loaiza ⁴,
 José Roberto Vega-Baudrit ^{2,5} and Rodolfo J. Gonzalez-Paz ^{2,*}

- ¹ Centro de Investigación y Extensión en Materiales, Escuela de Ciencia e Ingeniería de los Materiales, Instituto Tecnológico de Costa Rica, 159-7050 Cartago, Costa Rica; jeimygonz@gmail.com (J.G.-M.); jorgecubero@gmail.com (J.M.C.-S.)
 - ² Adolphe Merkle Institute, University of Fribourg, Chemin des Verdiers 4, 1700 Fribourg, Switzerland; yendry386@hotmail.com (Y.R.C.-U.); jvegab@gmail.com (J.R.V.-B.)
 - ³ Biological Assay Laboratory (LEBi), Universidad de Costa Rica, 11501-2060 San José, Costa Rica; sara.gonzalez@ucr.ac.cr
 - ⁴ National Center for Biotechnological Innovations (CENIBiot), National Center of High Technology (CeNAT-CONARE), Pavas, 1174-1200 San José, Costa Rica; none.0512@gmail.com (N.M.-U.); mbaizan@cenat.ac.cr (M.B.-R.); rloaiza@cenat.ac.cr (R.L.)
 - ⁵ National Laboratory of Nanotechnology (LANOTEC), National Center of High Technology (LANOTEC-CeNAT-CONARE), Pavas, 1174-1200 San José, Costa Rica
 - ⁶ Laboratorio de Polímeros (POLIUNA), Universidad Nacional, 86-3000 Heredia, Costa Rica
- * Correspondence: osarsip@gmail.com




Figure S1. AFM (A) amplitude, (B) phase image of the collagen control sample. (C) Cross- section of the line profile shown in A. The collagen D-pattern gap is similar in all the microfibers measured and all the microfibers show a homogeneous phase.



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).