

Article

N-[4-(*N,N,N*-Trimethylammonium)Benzyl]Chitosan Chloride as a Gene Carrier: The Influence of Polyplex Composition and Cell Type

Sergei V. Raik ¹, Tatiana V. Mashel ², Albert R. Muslimov ^{3,4,5}, Olga S. Epifanovskaya ³, Mikhail A. Trofimov ⁴, Daria N. Poshina ¹, Kirill V. Lepik ³ and Yury A. Skorik ^{1,*}

¹ Institute of Macromolecular Compounds of the Russian Academy of Sciences, Bolshoi pr. VO 31, 199004 Saint Petersburg, Russia; raiksv@gmail.com (S.V.R.); poschin@yandex.ru (D.N.P.)

² Department of Applied Optics, ITMO University, Kronverkskiy pr. 49, 197101 Saint Petersburg, Russia; t.v.mashel@gmail.com

³ R.M. Gorbacheva Research Institute of Pediatric Oncology, Hematology and Transplantation, Pavlov University, Lva Tolstogo 6/8, 197022 Saint Petersburg, Russia; albert.r.muslimov@gmail.com (A.R.M.); epifolga@rambler.ru (O.A.E.); lepikkv@gmail.com (K.V.L.)

⁴ Renewable Energy Laboratory, St. Petersburg Academic University, Khlopina 8/3 lit. A, 194021 Saint Petersburg, Russia; mihail.trofimov@pharminnotech.com

⁵ "QR.bio", Voronezhskaya 5 lit. A, 191119 Saint Petersburg, Russia; albert.r.muslimov@gmail.com

* Correspondence: yury_skorik@mail.ru




Figure S1. ^1H NMR spectrum of TMAB-CS37-26 (400 MHz, 343 K).




Figure S2. ^1H NMR spectrum of TMAB-CS37-54 (400 MHz, 343 K).




Figure S3. ${}^1\text{H}$ NMR spectrum of TMAB-CS37-71 (400 MHz, 343 K).




Figure S4. ^1H NMR spectrum of TMAB-CS135-64 (400 MHz, 343 K).




Figure S5. Flow cytometry data for HEK 293T transfection (No. 1-6 from Table 4 in the same order).




Figure S6. Flow cytometry data for K562 transfection (No. 1-6 from Table 4 in the same order).