

Supplementary Materials

Table S1. The most serious insect, disease, and parasitic plant agents for each of 419 tree species native to the conterminous United States, up to five per species. Nonnative invasive agents are indicated with *. Taxon (T) is identified for each agent (I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex), and Guild (G) is identified for insect agents (B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders). The agents are sorted by Severity (Sev.) within each species. The sources of the information are also provided.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Abies amabilis</i>	Pacific silver fir	Silver fir beetle (<i>Pseudohylesinus sericeus</i>)	I/B	5	[1,2]
		Balsam woolly adelgid (<i>Adelges piceae</i>)*	I/S	3	[2,3]
		Heterobasidion root disease P-type (<i>Heterobasidion irregularare</i>)	D	3	[2]
		Laminated root rot (<i>Phellinus sulphurascens</i>)	D	3	[4]
		Indian paint fungus (<i>Echinodontium tinctorium</i>)	D	1	[2]
<i>Abies balsamea</i>	balsam fir	Balsam woolly adelgid (<i>Adelges piceae</i>)*	I/S	8	[1,3,5]
		Spruce budworm (<i>Choristoneura fumiferana</i>)	I/F	8	[1]
		Armillaria root disease (<i>Armillaria</i> spp.)	D	1	[1,6]
		Red heart fungus (<i>Stereum sanguinolentum</i>)	D	1	[1,6,7]
		Rust fungus (<i>Melampsorella caryophyllacearum</i>)	D	1	[1,6]
<i>Abies bracteata</i>	bristlecone fir	Lophodermium needle cast (<i>Lophodermium</i> spp.)	D	1	[6,8]
		Needle cast (<i>Lirula nervisequia</i> var. <i>conspicua</i>)	D	1	[8-10]
		True fir bark beetles (<i>Scolytus dentatus</i>)	I/B	1	[11]
<i>Abies concolor</i>	white fir	Armillaria root disease (<i>Armillaria</i> spp.)	D	5	[12,13]
		Fir engraver (<i>Scolytus ventralis</i>)	I/B	5	[1]
		Douglas-fir tussock moth (<i>Orgyia pseudotsugata</i>)	I/F	3	[1,14,15]
		Heterobasidion root disease P-type (<i>Heterobasidion irregularare</i>)	D	3	[16]
		Western spruce budworm (<i>Choristoneura freemani</i>)	I/F	3	[1]
<i>Abies fraseri</i>	Fraser fir	Balsam woolly adelgid (<i>Adelges piceae</i>)*	I/S	8	[1]
		Phytophthora root rot (<i>Phytophthora cinnamomi</i>)*	D	5	[1,17]
		Armillaria root disease (<i>Armillaria</i> spp.)	D	3	[1]
		Two-spotted spider mite (<i>Tetranychus urticae</i>)	I/S	1	[1]
<i>Abies grandis</i>	grand fir	Armillaria root disease (<i>Armillaria</i> spp.)	D	8	[1,6,18]
		Fir engraver (<i>Scolytus ventralis</i>)	I/B	8	[1,19]

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Abies lasiocarpa</i>	subalpine fir	Laminated root rot (<i>Phellinus sulphurascens</i>)	D	8	[1,20]
		Balsam woolly adelgid (<i>Adelges piceae</i>)*	I/S	3	[2,3]
		Heterobasidion root disease P-type (<i>Heterobasidion irregulare</i>)	D	3	[1]
		Armillaria root disease (<i>Armillaria</i> spp.)	D	8	[1,18,21]
		Balsam woolly adelgid (<i>Adelges piceae</i>)*	I/S	8	[1,22,23]
		Western balsam bark beetle (<i>Dryocoetes confusus</i>)	I/B	8	[1,12,24]
		Heterobasidion root disease P-type (<i>Heterobasidion irregulare</i>)	D	3	[16]
		Western spruce budworm (<i>Choristoneura freemani</i>)	I/F	1	[1,25]
		Western balsam bark beetle (<i>Dryocoetes confusus</i>)	I/B	5	[1,26]
		Armillaria root disease (<i>Armillaria</i> spp.)	D	3	[1,18,21]
<i>Abies lasiocarpa</i> var. <i>arizonica</i>	corkbark fir	Heterobasidion root disease P-type (<i>Heterobasidion irregulare</i>)	D	3	[16]
		Western spruce budworm (<i>Choristoneura freemani</i>)	I/F	3	[1,25]
		Fir engraver (<i>Scolytus ventralis</i>)	I/B	3	[1,26,27]
		Heterobasidion root disease P-type (<i>Heterobasidion irregulare</i>)	D	3	[16]
		Indian paint fungus (<i>Echinodontium tinctorium</i>)	D	1	[1,28]
<i>Abies magnifica</i>	California red fir	Laminated root rot (<i>Phellinus sulphurascens</i>)	D	1	[4]
		Red fir dwarf mistletoe (<i>Arceuthobium abietinum</i> f. sp. <i>magnifica</i>)	P	1	[1]
		Fir engraver (<i>Scolytus ventralis</i>)	I/B	3	[29]
		Armillaria root disease (<i>Armillaria</i> spp.)	D	1	[29]
		Indian paint fungus (<i>Echinodontium tinctorium</i>)	D	1	[28]
<i>Abies magnifica</i> var. <i>shastensis</i>	Shasta red fir	Red fir dwarf mistletoe (<i>Arceuthobium abietinum</i> f. sp. <i>magnifica</i>)	P	1	[30]
		Red ring rot canker (<i>Phenillus cancriformans</i>)	D	1	[29]
		Indian paint fungus (<i>Echinodontium tinctorium</i>)	D	5	[1,31]
		Heterobasidion root disease P-type (<i>Heterobasidion irregulare</i>)	D	3	[16]
		Red ring rot (<i>Phellinus pini</i> var. <i>cancriformans</i>)	D	3	[1]
<i>Abies procera</i>	noble fir				

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Acacia farnesiana</i>	sweet acacia	Schweinitzii root and butt rot (<i>Phaeolus schweinitzii</i>)	D	3	[1]
		Laminated root rot (<i>Phellinus sulphurascens</i>)	D	1	[4]
		Clitocybe root rot (<i>Armillaria tabescens</i>)	D	1	[13,32]
		Cotton root rot (<i>Phymatotrichopsis omnivora</i>)	D	1	[13,32]
		Rust (<i>Ravenelia</i> spp.)	D	1	[32]
<i>Acacia greggii</i>	catclaw acacia	Twig girdler (<i>Oncideres pustulatus</i>)	I/B	1	[32]
		Mistletoe (<i>Phoradendron californicum</i>)	P	1	[33]
		Rust (<i>Ravenelia versatilis</i>)	D	1	[8]
		White rot (<i>Phellinus badius</i>)	D	1	[8]
<i>Acer floridanum</i>	Florida maple	Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	8	[34,35]
		Japanese beetle (<i>Popillia japonica</i>)*	I/F	1	[34,35]
		Oystershell scale (<i>Lepidosaphes ulmi</i>)*	I/S	1	[34,35]
<i>Acer glabrum</i>	Rocky Mountain maple	Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	8	[34,35]
		White spongy rot (<i>Phellinus igniarius</i>)	D	3	[13]
		Japanese beetle (<i>Popillia japonica</i>)*	I/F	1	[34,35]
		Oak bracket (<i>Inonotus dryadeus</i>)	D	1	[36]
		Oystershell scale (<i>Lepidosaphes ulmi</i>)*	I/S	1	[34,35]
		Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	8	[34,35]
<i>Acer grandidentatum</i>	bigtooth maple	Fall cankerworm (<i>Alsophila pometaria</i>)	I/F	3	[37]
		Japanese beetle (<i>Popillia japonica</i>)*	I/F	1	[34,35]
		Linden looper (<i>Erranis tiliae</i>)	I/F	1	[37]
		Oystershell scale (<i>Lepidosaphes ulmi</i>)*	I/S	1	[34,35]
		Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	8	[34,35]
<i>Acer leucoderme</i>	chalk maple	Japanese beetle (<i>Popillia japonica</i>)*	I/F	1	[34,35]
		Oystershell scale (<i>Lepidosaphes ulmi</i>)*	I/S	1	[34,35]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Acer macrophyllum</i>	bigleaf maple	Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	8	[34,35]
		Polyphagous shot hole borer (<i>Euwallacea</i> spp.)*	I/B	5	[38]
		Sudden oak death (<i>Phytophthora ramorum</i>)*	D	1	[7,39]
		Verticillium wilt (<i>Verticillium albo-atrum</i>)	D	1	[1]
<i>Acer negundo</i>	boxelder	Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	8	[34,35]
		Polyphagous shot hole borer (<i>Euwallacea</i> spp.)*	I/B	5	[38,40]
		Eutypella canker (<i>Eutypella parasitica</i>)	D	1	[1,41]
		Heart rot (<i>Inonotus glomeratus</i>)	D	1	[1,36]
		Verticillium wilt (<i>Verticillium albo-atrum</i>)	D	1	[1,42]
<i>Acer nigrum</i>	black maple	Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	8	[34,35]
		Eutypella canker (<i>Eutypella parasitica</i>)	D	1	[1,41]
		Japanese beetle (<i>Popillia japonica</i>)*	I/F	1	[34,35]
		Verticillium wilt (<i>Verticillium albo-atrum</i>)	D	1	[1,42]
<i>Acer pensylvanicum</i>	striped maple	Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	8	[34,35]
		Japanese beetle (<i>Popillia japonica</i>)*	I/F	1	[34,35]
		Leaf spot disease (<i>Cristulariella depraedans</i>)	D	1	[1]
		Pezicula trunk and branch cankers (<i>Pezicula subcarnea</i>)	D	1	[1]
		Verticillium wilt (<i>Verticillium albo-atrum</i>)	D	1	[1]
<i>Acer rubrum</i>	red maple	Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	8	[34,35,43]
		Armillaria root disease (<i>Armillaria</i> spp.)	D	1	[1]
		Bacterial leaf scorch (<i>Xylella fastidiosa</i>)	D	1	[42,44]
		Heart rot (<i>Inonotus glomeratus</i>)	D	1	[1,36,42]
		Verticillium wilt (<i>Verticillium albo-atrum</i>)	D	1	[35,42,45]
<i>Acer saccharinum</i>	silver maple	Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	8	[34,35]
		Armillaria root disease (<i>Armillaria</i> spp.)	D	1	[1]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Acer saccharum</i>	sugar maple	Japanese beetle (<i>Popillia japonica</i>)*	I/F	1	[34,35]
		Nectria canker (<i>Nectria ditissima</i>)	D	1	[1]
		Verticillium wilt (<i>Verticillium albo-atrum</i>)	D	1	[1,35]
		Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	8	[34,35]
		Armillaria root disease (<i>Armillaria</i> spp.)	D	1	[1]
		Bacterial leaf scorch (<i>Xylella fastidiosa</i>)	D	1	[42,44]
		Nectria canker (<i>Nectria ditissima</i>)	D	1	[1,36]
<i>Acer spicatum</i>	mountain maple	Verticillium wilt (<i>Verticillium albo-atrum</i>)	D	1	[1,42]
		Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	8	[34,35]
		Japanese beetle (<i>Popillia japonica</i>)*	I/F	1	[34,46]
		Nectria canker (<i>Nectria</i> sp.)	D	1	[33]
		Powdery mildew (<i>Phyllactinia guttata</i>)	D	1	[33]
		Verticillium wilt (<i>Verticillium dahliae</i>)	D	1	[33,47]
		Ganoderma butt rot (<i>Ganoderma zonatum</i>)	D	1	[36,48]
<i>Acoelorrhaphes wrightii</i>	Everglades palm	Nematode (<i>Rotylenchulus reniformis</i>)	D	1	[49]
		Stigmina leaf spot (<i>Stigmina palmivora</i>)	D	1	[36]
		Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	5	[34,35]
		Bacterial leaf scorch (<i>Xylella fastidiosa</i>)	D	1	[35,36,50]
		Eastern mistletoe (<i>Phoradendron leucarpum</i>)	P	1	[36]
<i>Aesculus californica</i>	California buckeye	Sudden oak death (<i>Phytophthora ramorum</i>)*	D	1	[36,51]
		Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	5	[34,35]
		Bacterial leaf scorch (<i>Xylella fastidiosa</i>)	D	1	[35,36,50]
		Eastern mistletoe (<i>Phoradendron leucarpum</i>)	P	1	[36]
		Sudden oak death (<i>Phytophthora ramorum</i>)*	D	1	[36,51]
		Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	5	[34,35]
		Heart rot (<i>Flammulina velutipes</i>)	D	1	[1]
<i>Aesculus flava</i>	yellow buckeye	Oak bracket (<i>Inonotus dryadeus</i>)	D	1	[36]
		Powdery mildew (<i>Erysiphe flexuosa</i>)	D	1	[1]
		Root-knot nematode (<i>Meloidogyne</i> spp.)	D	1	[52]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Aesculus glabra</i>	Ohio buckeye	Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	5	[34,35]
		Artist's cork (<i>Ganoderma applanatum</i>)	D	1	[1,35]
		Guignardia blotch (<i>Guignardia aesculi</i>)	D	1	[1]
		Powdery mildew (<i>Erysiphe flexuosa</i>)	D	1	[1]
<i>Aesculus glabra</i> var. <i>arguta</i>	Texas buckeye	Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	5	[34]
<i>Aesculus pavia</i>	red buckeye	Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	5	[34,35]
<i>Aesculus sylvatica</i>	painted buckeye	Guignardia blotch (<i>Guignardia aesculi</i>)	D	1	[36]
		Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	5	[34,35]
		Powdery mildew (<i>Erysiphe flexuosa</i>)	D	1	[53]
<i>Alnus incana</i> ssp. <i>tenuifolia</i>	thinleaf alder	Phytophthora disease of alders (<i>Phytophthora</i> sp.)	D	5	[54]
		Alder canker (<i>Valsa melanodiscus</i>)	D	3	[13,55]
		Green alder sawfly (<i>Monsoma pulveratum</i>)*	I/F	3	[56,57]
		Collar rot (<i>Phytophthora siskiyouensis</i>)	D	1	[36]
		Gypsy moth (<i>Lymantria dispar</i> <i>dispar</i>)*	I/F	3	[37]
<i>Alnus maritima</i>	seaside alder		.	.	.
<i>Alnus oblongifolia</i>	Arizona alder		.	.	.
<i>Alnus rhombifolia</i>	white alder		.	.	.
		Phytophthora disease of alders (<i>Phytophthora</i> sp.)	D	5	[54]
		Alder canker (<i>Valsa melanodiscus</i>)	D	3	[13]
		Green alder sawfly (<i>Monsoma pulveratum</i>)*	I/F	3	[56,57]
		Collar rot (<i>Phytophthora siskiyouensis</i>)	D	1	[36]
<i>Alnus rubra</i>	red alder	Alder bark beetle (<i>Alniphagus aspericollis</i>)	I/B	5	[11]
		Phytophthora disease of alders (<i>Phytophthora</i> sp.)	D	5	[54]
		Alder canker (<i>Valsa melanodiscus</i>)	D	3	[58]
		Forest tent caterpillar (<i>Malacosoma disstria</i>)	I/F	3	[1,59]
		Gypsy moth (<i>Lymantria dispar</i> <i>dispar</i>)*	I/F	3	[60]
<i>Amelanchier arborea</i>	common serviceberry	Blister canker (<i>Biscogniauxia marginata</i>)	D	1	[36]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Amelanchier sanguinea</i>	roundleaf serviceberry	Fire blight (<i>Erwinia amylovora</i>)	D	1	[61,62]
		Witches'-broom (<i>Apiosporina collinsii</i>)	D	1	[36]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[60]
		Blister canker (<i>Biscogniauxia marginata</i>)	D	1	[36]
		Fire blight (<i>Erwinia amylovora</i>)	D	1	[61,62]
		Witches'-broom (<i>Apiosporina collinsii</i>)	D	1	[36]
<i>Amyris elemifera</i>	sea torchwood	none	.	.	.
<i>Annona glabra</i>	pond-apple	none	.	.	.
<i>Arbutus arizonica</i>	Arizona madrone	Bark beetle (<i>Liparthrum arizonicum</i>)	I/B	1	[11]
<i>Arbutus menziesii</i>	Pacific madrone	Crown canker and root rot (<i>Phytophthora cactorum</i>)	D	3	[1,63]
		Heterobasidion root disease (<i>Heterobasidion</i> spp.)	D	3	[1,63]
		Madrone canker (<i>Neofusicoccum arbuti</i>)	D	3	[1,63,64]
		Sudden oak death (<i>Phytophthora ramorum</i>)*	D	3	[34,65]
		Arbutus canker (<i>Nattrassia mangiferae</i>)	D	1	[7,66]
		Crown canker and root rot (<i>Phytophthora cactorum</i>)	D	3	[1,63]
<i>Arbutus xalapensis</i>	Texas madrone	Heterobasidion root disease (<i>Heterobasidion</i> spp.)	D	3	[1,63]
		Madrone canker (<i>Neofusicoccum arbuti</i>)	D	3	[1,63,64]
		Arbutus canker (<i>Nattrassia mangiferae</i>)	D	1	[7,66]
		Sudden oak death (<i>Phytophthora ramorum</i>)*	D	1	[34,65]
		Coral spot fungus (<i>Nectria cinnabarina</i>)	D	1	[13,36]
		Japanese beetle (<i>Popillia japonica</i>)*	I/F	1	[67]
<i>Avicennia germinans</i>	black-mangrove	Butt rot fungi (<i>Phellinus swieteniae</i>)	D	3	[68]
<i>Betula alleghaniensis</i>	yellow birch	Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	5	[34,35]
		Bronze birch borer (<i>Agrilus anxius</i>)	I/B	1	[1,69,70]
		Eutypella canker (<i>Eutypella parasitica</i>)	D	1	[1,35,36]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Betula lenta</i>	sweet birch	Nectria canker (<i>Nectria ditissima</i>)	D	1	[1,35]
		Sterile conk trunk rot (<i>Inonotus obliquus</i>)	D	1	[1,36]
		Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	5	[34,35]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	1	[1,35]
		Nectria canker (<i>Nectria ditissima</i>)	D	1	[1,35]
<i>Betula nigra</i>	river birch	White spongy rot (<i>Phellinus igniarius</i>)	D	1	[1,35]
		Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	5	[34,35]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34,35]
		Birch leafminer (<i>Fenusia pusilla</i>)*	I/F	1	[34,35]
		Eastern mistletoe (<i>Phoradendron leucarpum</i>)	P	1	[1]
<i>Betula occidentalis</i>	water birch	Oystershell scale (<i>Lepidosaphes ulmi</i>)*	I/S	1	[34,35]
		Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	5	[34,35]
		Birch rust (<i>Melampsoridium betulinum</i>)	D	1	[36]
		Bronze birch borer (<i>Agrilus anxius</i>)	I/B	1	[35,69]
		Sterile conk trunk rot (<i>Inonotus obliquus</i>)	D	1	[36]
<i>Betula papyrifera</i>	paper birch	Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	5	[34,35]
		Bronze birch borer (<i>Agrilus anxius</i>)	I/B	5	[1,35,36]
		Birch leafminer (<i>Fenusia pusilla</i>)*	I/F	3	[1,34]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[1,35]
		Tinder conk (<i>Fomes fomentarius</i>)	D	1	[71]
<i>Betula populifolia</i>	gray birch	Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	5	[34,35]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34,35]
		Birch leafminer (<i>Fenusia pusilla</i>)*	I/F	1	[35,72]
		Bronze birch borer (<i>Agrilus anxius</i>)	I/B	1	[70,72]
		Sterile conk trunk rot (<i>Inonotus obliquus</i>)	D	1	[36]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Betula uber</i>	Virginia roundleaf birch	Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	1	[1,35]
		Japanese beetle (<i>Popillia japonica</i>)*	I/F	1	[34,73]
		Nectria canker (<i>Nectria ditissima</i>)	D	1	[1,35]
		White spongy rot (<i>Phellinus igniarius</i>)	D	1	[1,35]
<i>Bursera simaruba</i>	gumbo limbo	Rugose spiraling whitefly (<i>Aleurodicus rugioperculatus</i>)*	I/S	1	[74]
		Scale insect (<i>Phalacrocooccus howertoni</i>)	I/S	1	[74]
<i>Callitropsis nootkatensis</i>	Alaska yellow-cedar	Cedar bark beetle (<i>Phloeosinus spp.</i>)	I/B	3	[13]
		Juniper scale (<i>Carulaspis juniperi</i>)*	I/S	1	[34]
		Valsa canker and dieback (<i>Valsa abietis</i>)	D	1	[36]
		Heterobasidion root disease P-type (<i>Heterobasidion irregularare</i>)	D	3	[1,7]
<i>Calocedrus decurrens</i>	incense-cedar	Pocket dry rot (<i>Tyromyces amarus</i>)	D	3	[1,75]
		Broom rust (<i>Gymnosporangium libocedri</i>)	D	1	[1,7]
		Cedar bark beetle (<i>Phloeosinus spp.</i>)	I/B	1	[11]
		Juniper scale (<i>Carulaspis juniperi</i>)*	I/S	1	[1,76]
		Bacterial necrosis (<i>Erwinia cacticida</i>)	D	1	[77,78]
<i>Carnegiea gigantea</i>	saguaro	Cactus longhorn beetle (<i>Moneilema gigas</i>)	I/B	1	[78]
<i>Carpinus caroliniana</i>	musclewood	Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[35,79]
		Hornbeam anthracnose (<i>Apiosporopsis carpinea</i>)	D	1	[36]
		Nectria canker (<i>Nectria ditissima</i>)	D	1	[35]
		White rot (<i>Bjerkandera adusta</i>)	D	1	[80]
		Butt rot (<i>Hericium erinaceus</i>)	D	1	[36]
<i>Carya aquatica</i>	water hickory	Hickory borer (<i>Goes pulcher</i>)	I/B	1	[1]
		Armillaria root disease (<i>Armillaria spp.</i>)	D	1	[36]
		Hickory bark beetle (<i>Scolytus quadrispinosus</i>)	I/B	1	[1]
<i>Carya carolinae-septentrionalis</i>	southern shagbark hickory	Hickory borer (<i>Goes pulcher</i>)	I/B	1	[1]
<i>Carya cordiformis</i>	bitternut hickory				

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Carya floridana</i>	scrub hickory	Nectria canker (<i>Nectria ditissima</i>)	D	1	[1]
		Spiculosia canker (<i>Phellinus spiculosus</i>)	D	1	[1]
		White spongy rot (<i>Phellinus igniarius</i>)	D	1	[1]
<i>Carya glabra</i>	pignut hickory	Wood decay (<i>Phellinus gilvus</i>)	D	1	[81]
		Hickory bark beetle (<i>Scolytus quadrispinosus</i>)	I/B	1	[1]
<i>Carya illinoiensis</i>	pecan	Nectria canker (<i>Nectria ditissima</i>)	D	1	[1]
		Spiculosia canker (<i>Phellinus spiculosus</i>)	D	1	[1]
		Twig girdler (<i>Oncideres cingulata</i>)	I/B	1	[1]
		Bacterial leaf scorch (<i>Xylella fastidiosa</i>)	D	1	[44]
		Crown gall (<i>Agrobacterium tumefaciens</i>)	D	1	[1]
		Hickory bark beetle (<i>Scolytus quadrispinosus</i>)	I/B	1	[1]
		Hickory borer (<i>Goes pulcher</i>)	I/B	1	[1]
<i>Carya laciniosa</i>	shellbark hickory	Spiculosia canker (<i>Phellinus spiculosus</i>)	D	1	[1,82]
		Hickory bark beetle (<i>Scolytus quadrispinosus</i>)	I/B	1	[1]
		Hickory borer (<i>Goes pulcher</i>)	I/B	1	[1]
		Hickory spiral borer (<i>Agrilus arcuatus torquatus</i>)	I/B	1	[1]
<i>Carya myristiciformis</i>	nutmeg hickory	Twig girdler (<i>Oncideres cingulata</i>)	I/B	1	[1]
		Hickory bark beetle (<i>Scolytus quadrispinosus</i>)	I/B	1	[1]
		Twig girdler (<i>Oncideres cingulata</i>)	I/B	1	[1]
<i>Carya ovalis</i>	red hickory	Hickory bark beetle (<i>Scolytus quadrispinosus</i>)	I/B	1	[1]
		Nectria canker (<i>Nectria ditissima</i>)	D	1	[1]
<i>Carya ovata</i>	shagbark hickory	Spiculosia canker (<i>Phellinus spiculosus</i>)	D	1	[1]
		Twig girdler (<i>Oncideres cingulata</i>)	I/B	1	[1]
		Crown gall (<i>Agrobacterium tumefaciens</i>)	D	1	[1]
		Hickory bark beetle (<i>Scolytus quadrispinosus</i>)	I/B	1	[1]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
		Spiculosia canker (<i>Phellinus spiculosus</i>)	D	1	[1]
		Twig girdler (<i>Oncideres cingulata</i>)	I/B	1	[1]
		Witches broom (<i>Phytoplasma</i>)	D	1	[1,36]
<i>Carya pallida</i>	sand hickory	Wood decay (<i>Phellinus gilvus</i>)	D	1	[81]
<i>Carya texana</i>	black hickory	Twig girdler (<i>Oncideres cingulata</i>)	I/B	1	[83]
<i>Carya tomentosa</i>	mockernut hickory	Hickory bark beetle (<i>Scolytus quadrispinosus</i>)	I/B	1	[1]
		Hickory borer (<i>Goes pulcher</i>)	I/B	1	[1]
		Hickory spiral borer (<i>Agrilus arcuatus torquatus</i>)	I/B	1	[1]
		Pecan carpenterworm (<i>Cossula magnifica</i>)	I/B	1	[1]
		Spiculosia canker (<i>Phellinus spiculosus</i>)	D	1	[1]
<i>Castanea dentata</i>	American chestnut	Chestnut blight (<i>Cryphonectria parasitica</i>)*	D	10	[34,84]
		Phytophthora root rot (<i>Phytophthora cinnamomi</i>)*	D	8	[84]
		Twolined chestnut borer (<i>Agrilus bilineatus</i>)	I/B	3	[37]
		Asiatic oak weevil (<i>Cyrtepistomus castaneus</i>)*	I/F	1	[85]
		Oak bracket (<i>Inonotus dryadeus</i>)	D	1	[36]
<i>Castanea pumila</i>	Allegheny chinquapin	Chestnut blight (<i>Cryphonectria parasitica</i>)*	D	10	[86]
		Phytophthora root rot (<i>Phytophthora cinnamomi</i>)*	D	5	[86]
		Twolined chestnut borer (<i>Agrilus bilineatus</i>)	I/B	3	[37]
		Asiatic oak weevil (<i>Cyrtepistomus castaneus</i>)*	I/F	1	[86]
		Oak wilt (<i>Bretziella fagacearum</i>)*	D	1	[34]
<i>Castanea pumila</i> var. <i>ozarkensis</i>	Ozark chinquapin	Chestnut blight (<i>Cryphonectria parasitica</i>)*	D	10	[86]
		Phytophthora root rot (<i>Phytophthora cinnamomi</i>)*	D	5	[86]
		Twolined chestnut borer (<i>Agrilus bilineatus</i>)	I/B	3	[37]
		Asiatic oak weevil (<i>Cyrtepistomus castaneus</i>)*	I/F	1	[85,86]
		Chestnut gall wasp (<i>Dryocosmus kuriphilus</i>)*	I/F	1	[86]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Catalpa bignonioides</i>	southern catalpa	Verticillium wilt (<i>Verticillium dahliae</i>)	D	3	[87]
		Catalpa midge (<i>Cecidomyia catalpae</i>)	I/F	1	[87]
		Catalpa sphinx moth (<i>Ceratomia catalpae</i>)	I/F	1	[87]
		Decay (<i>Trametes versicolor</i>)	D	1	[87]
<i>Catalpa speciosa</i>	northern catalpa	Verticillium wilt (<i>Verticillium dahliae</i>)	D	3	[88]
		Catalpa midge (<i>Cecidomyia catalpae</i>)	I/F	1	[88]
		Catalpa sphinx moth (<i>Ceratomia catalpae</i>)	I/F	1	[88]
		Decay (<i>Trametes versicolor</i>)	D	1	[88]
<i>Celtis laevigata</i>	sugarberry	Basal canker and butt rot (<i>Kretzschmaria deusta</i>)	D	1	[36]
		Eastern mistletoe (<i>Phoradendron leucarpum</i>)	P	1	[1]
		Hackberry butterfly (<i>Asterocampa celtis</i>)	I/F	1	[1]
		Hackberry witches-broom complex (<i>Podosphaera phytophthora/Aceria celtis</i>)	D	1	[36]
<i>Celtis laevigata</i> var. <i>reticulata</i>	netleaf hackberry	Basal canker and butt rot (<i>Kretzschmaria deusta</i>)	D	1	[36]
		Eastern mistletoe (<i>Phoradendron leucarpum</i>)	P	1	[36]
<i>Celtis lindheimeri</i>	Lindheimer hackberry	Basal canker and butt rot (<i>Kretzschmaria deusta</i>)	D	1	[36]
		Eastern mistletoe (<i>Phoradendron leucarpum</i>)	P	1	[36]
<i>Celtis occidentalis</i>	hackberry	Bacterial leaf scorch (<i>Xylella fastidiosa</i>)	D	1	[44]
		Eastern mistletoe (<i>Phoradendron leucarpum</i>)	P	1	[1]
		Hackberry engraver (<i>Scolytus muticus</i>)	I/B	1	[1]
		Hackberry nipplegall maker (<i>Pachypsylla celtidismamma</i>)*	I/F	1	[1]
		Hackberry witches-broom complex (<i>Podosphaera phytophthora/Aceria celtis</i>)	D	1	[1,36]
		Botryosphaeria canker (<i>Botryosphaeria ribis</i>)	D	1	[1]
		Coral spot fungus (<i>Nectria cinnabrina</i>)	D	1	[36]
<i>Cercis canadensis</i>	eastern redbud	Verticillium wilt (<i>Verticillium albo-atrum</i>)	D	1	[1]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Cercocarpus ledifolius</i>	curlleaf mountain-mahogany	Geometrid moth (<i>Stamnodes animata</i>)*	I/F	3	[89]
		Mountain mahogany bark beetle (<i>Chaetophloeus heterodoxus</i>)	I/B	3	[13]
		Tent caterpillar (<i>Malacosoma</i> spp.)	I/F	3	[11,37]
		Mountain-mahogany looper (<i>Iridopsis clivinaria</i>)	I/F	1	[7]
		Wood-boring beetle (<i>Dicerca horni</i>)	I/B	1	[11]
<i>Chamaecyparis lawsoniana</i>	Port-Orford-cedar	Port-Orford-cedar root disease (<i>Phytophthora lateralis</i>)*	D	10	[1,90]
		Phytophthora root rot (<i>Phytophthora cinnamomi</i>)*	D	5	[7,13]
		Juniper scale (<i>Carulaspis juniperi</i>)*	I/S	1	[34]
		Seiridium canker (<i>Seiridium cardinale</i>)	D	1	[36]
<i>Chamaecyparis thyoides</i>	Atlantic whitecedar	Bagworm (<i>Thyridopteryx ephemeraeformis</i>)	I/F	1	[1]
		Heterobasidion root disease (<i>Heterobasidion</i> spp.)	D	1	[34]
		Juniper scale (<i>Carulaspis juniperi</i>)*	I/S	1	[34]
		Red-belted fungus (<i>Fomitopsis pinicola</i>)	D	1	[1]
		Schweinitzii root and butt rot (<i>Phaeolus schweinitzii</i>)	D	1	[1]
<i>Chilopsis linearis</i>	desert-willow	none	.	.	.
<i>Chionanthus virginicus</i>	fringetree	Emerald ash borer (<i>Agrilus planipennis</i>)*	I/B	3	[91,92]
<i>Chrysolepis chrysophylla</i>	golden chinquapin	Canker (<i>Phytophthora cambivora</i>)	D	5	[93]
		Oak wilt (<i>Bretziella fagacearum</i>)*	D	5	[34]
		Ganoderma rot of hardwoods (<i>Ganoderma lucidum</i>)	D	1	[35]
		White spongy rot (<i>Phellinus igniarius</i>)	D	1	[1]
		Cardin's whitefly (<i>Metaleurodicus cardini</i>)*	I/S	1	[94]
<i>Citharexylum fruticosum</i>	Florida fiddlewood	Fiddlewood leafroller (<i>Epicorsia oedipodalis</i>)	I/F	1	[95]
		Verticillium wilt (<i>Verticillium albo-atrum</i>)	D	3	[96]
		Fiddlewood leafroller (<i>Epicorsia oedipodalis</i>)	I/F	1	[95]
<i>Coccoloba diversifolia</i>	tietongue	Ganoderma root and butt rot (<i>Ganoderma australe</i>)	D	1	[36]
		False smut of palms (<i>Graphiooa phoenicis</i>)	D	3	[36]
<i>Coccothrinax argentata</i>	Florida silver palm				

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Colubrina elliptica</i>	soldierwood	Ganoderma butt rot (<i>Ganoderma zonatum</i>) none	D . .	1 . .	[36]
<i>Condalia hookeri</i>	bluewood	none
<i>Conocarpus erectus</i>	button mangrove	Lobate lac scale (<i>Paratachardina pseudolobata</i>)* Root rot (<i>Cylindrocladium scoparium</i>) Sri Lankan weevil (<i>Myllocerus undecimpustulatus undatus</i>)*	I/S D I/F	1 1 1	[97] [36,98] [99]
<i>Cordia boissieri</i>	anacahuita	none
<i>Cordia sebestena</i>	longleaf geigertree	Geiger tortoise beetle (<i>Eurypepla calochroma</i>) Tar spot (<i>Diatractium cordianum</i>)	I/F D	1 1	[100,101] [102]
<i>Cornus drummondii</i>	roughleaf dogwood	Dogwood anthracnose (<i>Discula destructiva</i>)*	D	5	[34]
<i>Cornus florida</i>	flowering dogwood	Dogwood anthracnose (<i>Discula destructiva</i>)* Dogwood borer (<i>Synanthedon scitula</i>) Bacterial leaf scorch (<i>Xylella fastidiosa</i>) Crown canker and root rot (<i>Phytophthora cactorum</i>) Dogwood clubgall midge (<i>Resseliella clavula</i>)	D I/B D D I/F	5 3 1 1 1	[34,36] [1,103] [44] [1,36] [1]
<i>Cornus nuttallii</i>	Pacific dogwood	Dogwood anthracnose (<i>Discula destructiva</i>)* Crown canker and root rot (<i>Phytophthora cactorum</i>) Madrone thrips (<i>Thrips madronii</i>) Nectria canker (<i>Nectria ditissima</i>)	D D I/F D	8 1 1 1	[34,36] [36] [13] [13,35]
<i>Cotinus obovatus</i>	smoketree	Verticillium wilt (<i>Verticillium albo-atrum</i>)	D	1	[13,104]
<i>Crataegus brainerdii</i>	Brainerd's hawthorn	Cherry bark tortrix (<i>Enarmonia formosana</i>)* Fire blight (<i>Erwinia amylovora</i>) Leaf blight and fruit rot (<i>Monilinia johnsonii</i>) Rust (<i>Gymnosporangium</i> spp.) Winter moth (<i>Operophtera brumata</i>)*	I/B D D D I/F	3 1 1 1 1	[34,105] [106] [106] [106] [34]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Crataegus calpodendron</i>	pear hawthorn	Cherry bark tortrix (<i>Enarmonia formosana</i>)*	I/B	3	[34,105]
		Fire blight (<i>Erwinia amylovora</i>)	D	1	[107]
		Leaf blight and fruit rot (<i>Monilinia johnsonii</i>)	D	1	[107]
		Rust (<i>Gymnosporangium</i> spp.)	D	1	[107]
		Winter moth (<i>Operophtera brumata</i>)*	I/F	1	[34]
		Cherry bark tortrix (<i>Enarmonia formosana</i>)*	I/B	3	[34,105]
<i>Crataegus chrysocarpa</i>	fireberry hawthorn	Fire blight (<i>Erwinia amylovora</i>)	D	1	[107]
		Leaf blight and fruit rot (<i>Monilinia johnsonii</i>)	D	1	[107]
		Rust (<i>Gymnosporangium</i> spp.)	D	1	[107]
		Winter moth (<i>Operophtera brumata</i>)*	I/F	1	[34]
		Cherry bark tortrix (<i>Enarmonia formosana</i>)*	I/B	3	[34,105]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34,108]
<i>Crataegus crus-galli</i>	cockspur hawthorn	Hawthorn rust (<i>Gymnosporangium globosum</i>)	D	1	[109]
		Winter moth (<i>Operophtera brumata</i>)*	I/F	1	[34]
		Cherry bark tortrix (<i>Enarmonia formosana</i>)*	I/B	3	[34,105]
		Fire blight (<i>Erwinia amylovora</i>)	D	1	[110]
		Leaf blight and fruit rot (<i>Monilinia johnsonii</i>)	D	1	[110]
		Rust (<i>Gymnosporangium</i> spp.)	D	1	[110]
<i>Crataegus dilatata</i>	broadleaf hawthorn	Winter moth (<i>Operophtera brumata</i>)*	I/F	1	[34]
		Cherry bark tortrix (<i>Enarmonia formosana</i>)*	I/B	3	[34,105]
		Fire blight (<i>Erwinia amylovora</i>)	D	1	[110]
		Leaf blight and fruit rot (<i>Monilinia johnsonii</i>)	D	1	[110]
		Rust (<i>Gymnosporangium</i> spp.)	D	1	[110]
		Winter moth (<i>Operophtera brumata</i>)*	I/F	1	[34]
<i>Crataegus flabellata</i>	fanleaf hawthorn	Cherry bark tortrix (<i>Enarmonia formosana</i>)*	I/B	3	[34,105]
		Fire blight (<i>Erwinia amylovora</i>)	D	1	[111]
		Leaf blight and fruit rot (<i>Monilinia johnsonii</i>)	D	1	[111]
		Rust (<i>Gymnosporangium</i> spp.)	D	1	[111]
		Winter moth (<i>Operophtera brumata</i>)*	I/F	1	[34]
		Cherry bark tortrix (<i>Enarmonia formosana</i>)*	I/B	3	[34,105]
<i>Crataegus mollis</i>	downy hawthorn				

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Crataegus pedicellata</i>	scarlet hawthorn	Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34,108]
		Winter moth (<i>Operophtera brumata</i>)*	I/F	1	[34]
		Cherry bark tortrix (<i>Enarmonia formosana</i>)*	I/B	3	[34,105]
		Fire blight (<i>Erwinia amylovora</i>)	D	1	[107]
		Leaf blight and fruit rot (<i>Monilinia johnsonii</i>)	D	1	[107]
		Rust (<i>Gymnosporangium</i> spp.)	D	1	[107]
		Winter moth (<i>Operophtera brumata</i>)*	I/F	1	[34]
<i>Diospyros texana</i>	Texas persimmon	Verticillium wilt (<i>Verticillium albo-atrum</i>)	D	3	[36]
<i>Diospyros virginiana</i>	common persimmon	Persimmon wilt (<i>Acremonium diospyri</i>)	D	5	[1,36]
		Verticillium wilt (<i>Verticillium albo-atrum</i>)	D	3	[36]
<i>Ebenopsis ebano</i>	blackbead ebony	none	.	.	.
<i>Ehretia anacua</i>	Anacua knockaway	none	.	.	.
<i>Eugenia rhombea</i>	red stopper	none	.	.	.
<i>Exothea paniculata</i>	butterbough	none	.	.	.
<i>Fagus grandifolia</i>	American beech	Beech bark disease (<i>Neonectria</i> spp./ <i>Cryptococcus fagisuga</i>)*	C	8	[1,36,112,113]
		White spongy rot (<i>Phellinus igniarius</i>)	D	3	[1,36]
		Armillaria root disease (<i>Armillaria</i> spp.)	D	1	[1]
		Artist's conk (<i>Ganoderma applanatum</i>)	D	1	[1,36]
		Forest tent caterpillar (<i>Malacosoma disstria</i>)	I/F	1	[1]
		Lobate lac scale (<i>Paratachardina pseudolobata</i>)*	I/S	8	[97]
		Scale insect (<i>Phalacrocooccus howertoni</i>)	I/S	3	[114]
<i>Ficus aurea</i>	Florida strangler fig	Edwards wasp moth (<i>Lymire edwardsii</i>)	I/F	1	[115]
		Ficus whitefly (<i>Singhiella simplex</i>)*	I/S	1	[116]
		Lobate lac scale (<i>Paratachardina pseudolobata</i>)*	I/S	8	[97]
		Scale insect (<i>Phalacrocooccus howertoni</i>)	I/S	3	[114]
<i>Ficus citrifolia</i>	wild banyantree				

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Frangula purshiana</i>	cascara buckthorn	Edwards wasp moth (<i>Lymire edwardsii</i>)	I/F	1	[115]
		Ficus whitefly (<i>Singhiella simplex</i>)*	I/S	1	[116]
		Hispidus canker (<i>Inonotus hispidus</i>)	D	1	[36]
		Sudden oak death (<i>Phytophthora ramorum</i>)*	D	1	[117]
<i>Fraxinus americana</i>	white ash	Emerald ash borer (<i>Agrilus planipennis</i>)*	I/B	10	[34,35,118]
		Ash yellows (<i>Candidatus Phytoplasma fraxini</i>)	D	5	[36,119]
		Oystershell scale (<i>Lepidosaphes ulmi</i>)*	I/S	3	[34,35]
		Forest tent caterpillar (<i>Malacosoma disstria</i>)	I/F	1	[1]
		Verticillium wilt (<i>Verticillium spp.</i>)	D	1	[35,120,121]
<i>Fraxinus berlandieriana</i>	Berlandier ash	Emerald ash borer (<i>Agrilus planipennis</i>)*	I/B	9	[34,35]
		Ash yellows (<i>Candidatus Phytoplasma fraxini</i>)	D	5	[36]
		Oystershell scale (<i>Lepidosaphes ulmi</i>)*	I/S	3	[34,35]
		Forest tent caterpillar (<i>Malacosoma disstria</i>)	I/F	1	[37]
<i>Fraxinus caroliniana</i>	Carolina ash	Emerald ash borer (<i>Agrilus planipennis</i>)*	I/B	10	[34,35]
		Ash yellows (<i>Candidatus Phytoplasma fraxini</i>)	D	5	[13]
		Oystershell scale (<i>Lepidosaphes ulmi</i>)*	I/S	3	[34,35]
		Forest tent caterpillar (<i>Malacosoma disstria</i>)	I/F	1	[37]
<i>Fraxinus cuspidata</i>	fragrant ash	Emerald ash borer (<i>Agrilus planipennis</i>)*	I/B	9	[34,35]
		Ash yellows (<i>Candidatus Phytoplasma fraxini</i>)	D	5	[13]
		Oystershell scale (<i>Lepidosaphes ulmi</i>)*	I/S	3	[34,35]
		Forest tent caterpillar (<i>Malacosoma disstria</i>)	I/F	1	[37]
<i>Fraxinus dipetala</i>	two-petal ash	Emerald ash borer (<i>Agrilus planipennis</i>)*	I/B	8	[34,35,118]
		Ash yellows (<i>Candidatus Phytoplasma fraxini</i>)	D	5	[13]
		Oystershell scale (<i>Lepidosaphes ulmi</i>)*	I/S	3	[34,35]
		Forest tent caterpillar (<i>Malacosoma disstria</i>)	I/F	1	[37]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Fraxinus greggii</i>	Gregg ash	Emerald ash borer (<i>Agrilus planipennis</i>)*	I/B	9	[34,35]
		Ash yellows (<i>Candidatus Phytoplasma fraxini</i>)	D	5	[13]
		Oystershell scale (<i>Lepidosaphes ulmi</i>)*	I/S	3	[34,35]
		Forest tent caterpillar (<i>Malacosoma disstria</i>)	I/F	1	[37]
<i>Fraxinus latifolia</i>	Oregon ash	Emerald ash borer (<i>Agrilus planipennis</i>)*	I/B	8	[34,35]
		Ash yellows (<i>Candidatus Phytoplasma fraxini</i>)	D	5	[36]
		Oystershell scale (<i>Lepidosaphes ulmi</i>)*	I/S	3	[34,35]
		Forest tent caterpillar (<i>Malacosoma disstria</i>)	I/F	1	[37]
		Verticillium wilt (<i>Verticillium spp.</i>)	D	1	[13]
<i>Fraxinus nigra</i>	black ash	Emerald ash borer (<i>Agrilus planipennis</i>)*	I/B	10	[34,35,118]
		Ash yellows (<i>Candidatus Phytoplasma fraxini</i>)	D	5	[119]
		Oystershell scale (<i>Lepidosaphes ulmi</i>)*	I/S	3	[34,35]
		Forest tent caterpillar (<i>Malacosoma disstria</i>)	I/F	1	[37]
		Nectria canker (<i>Nectria ditissima</i>)	D	1	[1]
<i>Fraxinus papillosa</i>	Chihuahua ash	Emerald ash borer (<i>Agrilus planipennis</i>)*	I/B	9	[34,35]
		Ash yellows (<i>Candidatus Phytoplasma fraxini</i>)	D	5	[13]
		Oystershell scale (<i>Lepidosaphes ulmi</i>)*	I/S	3	[34,35]
		Forest tent caterpillar (<i>Malacosoma disstria</i>)	I/F	1	[37]
<i>Fraxinus pennsylvanica</i>	green ash	Emerald ash borer (<i>Agrilus planipennis</i>)*	I/B	10	[34,35,118]
		Ash yellows (<i>Candidatus Phytoplasma fraxini</i>)	D	5	[36]
		Oystershell scale (<i>Lepidosaphes ulmi</i>)*	I/S	3	[34,35]
		Forest tent caterpillar (<i>Malacosoma disstria</i>)	I/F	1	[37]
		Verticillium wilt (<i>Verticillium dahliae</i>)	D	1	[36,122]
<i>Fraxinus profunda</i>	pumpkin ash	Emerald ash borer (<i>Agrilus planipennis</i>)*	I/B	10	[34,35]
		Ash yellows (<i>Candidatus Phytoplasma fraxini</i>)	D	5	[36]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Fraxinus quadrangulata</i>	blue ash	Oystershell scale (<i>Lepidosaphes ulmi</i>)*	I/S	3	[34,35]
		Forest tent caterpillar (<i>Malacosoma disstria</i>)	I/F	1	[37]
		Emerald ash borer (<i>Agrilus planipennis</i>)*	I/B	8	[34,35,118]
		Ash yellows (<i>Candidatus Phytoplasma fraxini</i>)	D	5	[36,119]
		Oystershell scale (<i>Lepidosaphes ulmi</i>)*	I/S	3	[34,35]
		Forest tent caterpillar (<i>Malacosoma disstria</i>)	I/F	1	[37]
<i>Fraxinus texensis</i>	Texas ash	Verticillium wilt (<i>Verticillium</i> spp.)	D	1	[121]
		Emerald ash borer (<i>Agrilus planipennis</i>)*	I/B	9	[34,35]
		Ash yellows (<i>Candidatus Phytoplasma fraxini</i>)	D	5	[36]
		Oystershell scale (<i>Lepidosaphes ulmi</i>)*	I/S	3	[34,35]
		Forest tent caterpillar (<i>Malacosoma disstria</i>)	I/F	1	[37]
		Emerald ash borer (<i>Agrilus planipennis</i>)*	I/B	9	[34,35]
<i>Fraxinus velutina</i>	velvet ash	Ash yellows (<i>Candidatus Phytoplasma fraxini</i>)	D	5	[36,123]
		Oystershell scale (<i>Lepidosaphes ulmi</i>)*	I/S	3	[34,35]
		Forest tent caterpillar (<i>Malacosoma disstria</i>)	I/F	1	[37]
		Emerald ash borer (<i>Agrilus planipennis</i>)*	I/B	9	[34,35]
		Ash yellows (<i>Candidatus Phytoplasma fraxini</i>)	D	5	[36,123]
		Oystershell scale (<i>Lepidosaphes ulmi</i>)*	I/S	3	[34,35]
<i>Gleditsia aquatica</i>	waterlocust	Forest tent caterpillar (<i>Malacosoma disstria</i>)	I/F	1	[37]
		Calico scale (<i>Eulecanium cerasorum</i>)*	I/S	1	[34]
		Mimosa webworm (<i>Homadaula anisocentra</i>)*	I/F	1	[1]
		Thyronectria canker (<i>Pleonectria austroamericana</i>)	D	1	[1,35,36]
		Twig girdler (<i>Oncideres cingulata</i>)	I/B	1	[1]
		White mottled rot (<i>Ganoderma</i> spp.)	D	1	[1,36]
<i>Gleditsia triacanthos</i>	honeylocust	Honeylocust agrilus (<i>Agrilus difficilis</i>)	I/B	3	[1]
		Calico scale (<i>Eulecanium cerasorum</i>)*	I/S	1	[34]
		Thyronectria canker (<i>Pleonectria austroamericana</i>)	D	1	[1,35,36]
		Twig girdler (<i>Oncideres cingulata</i>)	I/B	1	[1]
		White mottled rot (<i>Ganoderma</i> spp.)	D	1	[1,36]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Gordonia lasianthus</i>	loblolly bay	none	.	.	.
<i>Guapira discolor</i>	beeftree	none	.	.	.
<i>Gymnocladus dioicus</i>	Kentucky coffeetree	Verticillium wilt (<i>Verticillium</i> spp.) White mottled rot (<i>Ganoderma</i> spp.)	D D	1 1	[36] [36]
<i>Halesia carolina</i>	Carolina silverbell	none	.	.	.
<i>Halesia diptera</i>	two-wing silverbell	none	.	.	.
<i>Halesia parviflora</i>	little silverbell	none	.	.	.
<i>Helietta parvifolia</i>	barreta	none	.	.	.
<i>Hesperocyparis abramsiana</i>	Santa Cruz cypress	Seiridium canker (<i>Seiridium cardinale</i>)	D	1	[13]
<i>Hesperocyparis arizonica</i>	Arizona cypress	Cercospora blight (<i>Asperisporium sequoiae</i>) Cedar bark beetle (<i>Phloeosinus</i> spp.) Seiridium canker (<i>Seiridium cardinale</i>) Juniper scale (<i>Carulaspis juniperi</i>)* Rust (<i>Gymnosporangium cunninghamianum</i>)	D I/B D I/S D	5 3 3 1 1	[36,124,125] [11] [36] [34] [36]
<i>Hesperocyparis bakeri</i>	Baker cypress	Seiridium canker (<i>Seiridium cardinale</i>) Mistletoe (<i>Phoradendron bolleanum</i>) Rust (<i>Gymnosporangium cunninghamianum</i>)	D P D	3 1 1	[36] [126] [36]
<i>Hesperocyparis forbesii</i>	Tecate cypress	Cercospora blight (<i>Asperisporium sequoiae</i>) Seiridium canker (<i>Seiridium cardinale</i>) Juniper scale (<i>Carulaspis juniperi</i>)* Rust (<i>Gymnosporangium cunninghamianum</i>)	D D I/S D	5 3 1 1	[36,124] [36] [34] [36]
<i>Hesperocyparis goveniana</i>	Gowen cypress	Seiridium canker (<i>Seiridium cardinale</i>) Mistletoe (<i>Phoradendron bolleanum</i>)	D P	3 1	[36] [126]
<i>Hesperocyparis macnabiana</i>	Macnab's cypress	Mistletoe (<i>Phoradendron bolleanum</i>) Seiridium canker (<i>Seiridium cardinale</i>)	P D	1 1	[126] [36]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Hesperocyparis macrocarpa</i>	Monterey cypress	Cercospora blight (<i>Asperisporium sequoiae</i>) Seiridium canker (<i>Seiridium cardinale</i>)	D D	5 5	[36,124] [36]
<i>Hesperocyparis nevadensis</i>	Paiute cypress	Seiridium canker (<i>Seiridium cardinale</i>) Mistletoe (<i>Phoradendron bolleanum</i>)	D P	3 1	[36] [36]
<i>Hesperocyparis sargentii</i>	Sargent's cypress	Mistletoe (<i>Phoradendron bolleanum</i>) Seiridium canker (<i>Seiridium cardinale</i>)	P D	1 1	[126] [7]
<i>Hesperocyparis stephensonii</i>	Cuyamaca cypress	Seiridium canker (<i>Seiridium cardinale</i>) Mistletoe (<i>Phoradendron bolleanum</i>)	D P	3 1	[36] [36]
<i>Hippomane mancinella</i>	manchineel	none	.	.	.
<i>Ilex cassine</i>	dahoon	Florida wax scale (<i>Ceroplastes floridensis</i>)	I/S	3	[127]
<i>Ilex montana</i>	mountain winterberry	none	.	.	.
<i>Ilex myrtifolia</i>	myrtle dahoon	none	.	.	.
<i>Ilex opaca</i>	American holly	Florida wax scale (<i>Ceroplastes floridensis</i>) Black root rot (<i>Thielaviopsis basicola</i>)	I/S D	3 1	[127] [36]
<i>Juglans californica</i>	southern California black walnut	Thousand cankers disease (<i>Geosmithia morbida/Pityophthorus juglandis</i>)	D	3	[128,129]
<i>Juglans cinerea</i>	butternut	Butternut canker (<i>Sirococcus clavigignenti-juglandacearum</i>)* Thousand cankers disease (<i>Geosmithia morbida/Pityophthorus juglandis</i>) Bunch disease (Phytoplasma strains in subgroup of Prunus X-disease group)	D D D	10 5 1	[1,34] [128,129] [1,36]
<i>Juglans hindsii</i>	northern California black walnut	Blackline (Cherry leafroll virus) Thousand cankers disease (<i>Geosmithia morbida/Pityophthorus juglandis</i>)	D D	5 3	[36] [128,129]
<i>Juglans hindsii</i>	northern California black walnut	Sooty canker (<i>Fusicoccum mangiferum</i>)	D	1	[36]
<i>Juglans major</i>	Arizona walnut	Thousand cankers disease (<i>Geosmithia morbida/Pityophthorus juglandis</i>) White rot (<i>Phellinus weiri</i>)	D D	1 1	[128-130] [36]
<i>Juglans microcarpa</i>	Texas walnut	Bunch disease (Phytoplasma strains in subgroup of Prunus X-disease group)	D	3	[36]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Juglans nigra</i>	black walnut	Thousand cankers disease (<i>Geosmithia morbida/Pityophthorus juglandis</i>)	D	3	[128-130]
		Thousand cankers disease (<i>Geosmithia morbida/Pityophthorus juglandis</i>)	D	5	[128,129]
		Nectria canker (<i>Nectria ditissima</i>)	D	3	[1]
		Black stem borer (<i>Xylosandrus germanus</i>)*	I/B	1	[1]
		Flatheaded apple tree borer (<i>Chrysobothris femorata</i>)	I/B	1	[1]
		Walnut caterpillar (<i>Datana integerrima</i>)	I/F	1	[1,131]
<i>Juniperus ashei</i>	Ashe juniper	Bagworm (<i>Thyridopteryx ephemeraeformis</i>)	I/F	1	[35,132,133]
		Juniper scale (<i>Carulaspis juniperi</i>)*	I/S	1	[34]
<i>Juniperus californica</i>	California juniper	Juniper scale (<i>Carulaspis juniperi</i>)*	I/S	1	[34]
<i>Juniperus coahuilensis</i>	redberry juniper	Juniper scale (<i>Carulaspis juniperi</i>)*	I/S	1	[34]
<i>Juniperus deppeana</i>	alligator juniper	Juniper scale (<i>Carulaspis juniperi</i>)*	I/S	1	[34]
<i>Juniperus flaccida</i>	drooping juniper	Bagworm (<i>Thyridopteryx ephemeraeformis</i>)	I/F	1	[35,132]
<i>Juniperus monosperma</i>	oneseed juniper	Juniper mistletoe (<i>Phoradendron juniperinum</i>)	P	1	[7]
		Black-horned juniper borer (<i>Callidium texanum</i>)	I/B	1	[12]
		Cedar bark beetle (<i>Phloeosinus spp.</i>)	I/B	1	[11,134]
		Juniper borer (<i>Atimia huachucae</i>)	I/B	1	[12]
		Juniper mistletoe (<i>Phoradendron juniperinum</i>)	P	1	[36]
		Juniper twig pruner (<i>Styloxus bicolor</i>)	I/B	1	[12]
<i>Juniperus occidentalis</i>	western juniper	Cedar bark beetle (<i>Phloeosinus spp.</i>)	I/B	3	[11]
		Heterobasidion root disease P-type (<i>Heterobasidion irregulare</i>)	D	3	[135]
		Juniper mistletoe (<i>Phoradendron juniperinum</i>)	P	1	[36]
		Juniper pocket rot (<i>Pyrofomes demidoffii</i>)	D	1	[1]
		Juniper scale (<i>Carulaspis juniperi</i>)*	I/S	1	[34]
		Cedar bark beetle (<i>Phloeosinus spp.</i>)	I/B	1	[11]
<i>Juniperus osteosperma</i>	Utah juniper	Juniper mistletoe (<i>Phoradendron juniperinum</i>)	P	1	[36]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Juniperus pinchotii</i>	Pinchot juniper	Juniper pocket rot (<i>Pyrofomes demidoffii</i>)	D	1	[136,137]
		Juniper scale (<i>Carulaspis juniperi</i>)*	I/S	1	[34]
		Rust (<i>Gymnosporangium</i> spp.)	D	1	[1]
<i>Juniperus scopulorum</i>	Rocky Mountain juniper	Juniper scale (<i>Carulaspis juniperi</i>)*	I/S	1	[34]
		Cedar bark beetle (<i>Phloeosinus</i> spp.)	I/B	3	[11]
<i>Juniperus virginiana</i>	eastern redcedar	Cercospora blight of juniper (<i>Pseudocercospora juniperi</i>)	D	3	[36,124]
		Juniper mistletoe (<i>Phoradendron juniperinum</i>)	P	1	[36]
		Juniper scale (<i>Carulaspis juniperi</i>)*	I/S	1	[34]
		Rust (<i>Gymnosporangium</i> spp.)	D	1	[1]
		Cercospora blight of juniper (<i>Pseudocercospora juniperi</i>)	D	3	[36,124]
		Heterobasidion root disease P-type (<i>Heterobasidion irregulare</i>)	D	3	[1]
		Phytophthora root rot (<i>Phytophthora cinnamomi</i>)*	D	3	[34]
<i>Juniperus virginiana</i> var. <i>silicicola</i>	southern redcedar	Eastern juniper bark beetle (<i>Phloeosinus dentatus</i>)	I/B	1	[1]
		Juniper scale (<i>Carulaspis juniperi</i>)*	I/S	1	[1]
		Cercospora blight of juniper (<i>Pseudocercospora juniperi</i>)	D	3	[36,124]
		Heterobasidion root disease P-type (<i>Heterobasidion irregulare</i>)	D	3	[1]
		Phytophthora root rot (<i>Phytophthora cinnamomi</i>)*	D	3	[34]
<i>Laguncularia racemosa</i>	white-mangrove	Eastern juniper bark beetle (<i>Phloeosinus dentatus</i>)	I/B	1	[1]
		Juniper scale (<i>Carulaspis juniperi</i>)*	I/S	1	[1]
<i>Larix laricina</i>	tamarack	none	.	.	.
<i>Larix lyallii</i>	subalpine larix	Larch casebearer (<i>Coleophora laricella</i>)*	I/F	5	[1]
		Larch sawfly (<i>Pristiphora erichsonii</i>)	I/F	5	[1]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Spruce budworm (<i>Choristoneura fumiferana</i>)	I/F	3	[1]
		European larch canker (<i>Lachnellula willkommii</i>)*	D	1	[36]
		Heterobasidion root disease P-type (<i>Heterobasidion irregulare</i>)	D	3	[36]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Larix occidentalis</i>	western larix	Brown trunk rot (<i>Fomitopsis officinalis</i>)	D	1	[1]
		Larch needle blight (<i>Hypodermella laricis</i>)	D	1	[7]
		Larch sawfly (<i>Pristiphora erichsonii</i>)	I/F	1	[34]
		Larch dwarf mistletoe (<i>Arceuthobium laricis</i>)	P	5	[36]
		Armillaria root disease (<i>Armillaria</i> spp.)	D	3	[36,138]
		Larch casebearer (<i>Coleophora laricella</i>)*	I/F	3	[1]
		Larch sawfly (<i>Pristiphora erichsonii</i>)	I/F	3	[1]
<i>Leucaena pulverulenta</i>	great leadtree	none	.	.	.
<i>Liquidambar styraciflua</i>	sweetgum	Botryosphaeria canker (<i>Botryosphaeria ribis</i>)	D	1	[13,36]
		Calico scale (<i>Eulecanium cerasorum</i>)*	I/S	1	[34]
		Forest tent caterpillar (<i>Malacosoma disstria</i>)	I/F	1	[1,35,139,140]
		Ganoderma rot of hardwoods (<i>Ganoderma lucidum</i>)	D	1	[1]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	1	[34]
		Stem canker (<i>Fusarium solani</i>)	D	1	[1]
		Tuliptree scale (<i>Toumeyella liriodendri</i>)	I/S	1	[1]
<i>Liriodendron tulipifera</i>	yellow-poplar	Verticillium wilt (<i>Verticillium</i> spp.)	D	1	[36]
		Yellow poplar weevil (<i>Odontopus calceatus</i>)	I/F	1	[1]
		Sudden oak death (<i>Phytophthora ramorum</i>)*	D	10	[36]
		California oakworm (<i>Phryganidea californica</i>)	I/F	1	[1]
		Canker-rot (<i>Inonotus</i> spp.)	D	1	[1]
<i>Lithocarpus densiflorus</i>	tanoak	none	.	.	.
<i>Lyonia ferruginea</i>	Catalina ironwood	Ganoderma rot of hardwoods (<i>Ganoderma lucidum</i>)	D	1	[36]
		Lac scale (<i>Tachardiella mexicana</i>)	I/S	1	[141]
		Nectriella gall (<i>Nectriella pironii</i>)	D	1	[36]
		none	.	.	.
<i>Lysiloma watsonii</i>	littleleaf lysiloma	none	.	.	.

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Maclura pomifera</i>	Osage-orange	Cotton root rot (<i>Phymatotrichopsis omnivora</i>)	D	1	[1]
		Eastern mistletoe (<i>Phoradendron leucarpum</i>)	P	1	[1]
		Verticillium wilt (<i>Verticillium albo-atrum</i>)	D	1	[1]
<i>Magnolia acuminata</i>	cuckertree	Magnolia scale (<i>Neolecanium cornuparvum</i>)	I/S	1	[1]
		Nectria canker (<i>Nectria ditissima</i>)	D	1	[1]
<i>Magnolia ashei</i>	Ashe's magnolia	none	.	.	.
<i>Magnolia fraseri</i>	mountain magnolia	Magnolia scale (<i>Neolecanium cornuparvum</i>)	I/S	1	[1]
		Nectria canker (<i>Nectria magnoliae</i>)	D	1	[1]
<i>Magnolia grandiflora</i>	Southern magnolia	Magnolia scale (<i>Neolecanium cornuparvum</i>)	I/S	1	[1]
		Tuliptree scale (<i>Toumeyella liriodendri</i>)	I/S	1	[1]
		Verticillium wilt (<i>Verticillium albo-atrum</i>)	D	1	[142]
<i>Magnolia macrophylla</i>	bigleaf magnolia	none	.	.	.
<i>Magnolia pyramidata</i>	pyramid magnolia	none	.	.	.
<i>Magnolia tripetala</i>	umbrella magnolia	none	.	.	.
<i>Magnolia virginiana</i>	sweetbay	none	.	.	.
<i>Malus angustifolia</i>	southern crabapple	Apple scab (<i>Venturia inaequalis</i>)	D	1	[36]
		Fire blight (<i>Erwinia amylovora</i>)	D	1	[143]
<i>Malus coronaria</i>	sweet crabapple	Apple scab (<i>Venturia inaequalis</i>)	D	1	[36]
		Blister canker (<i>Biscogniauxia marginata</i>)	D	1	[36]
<i>Malus fusca</i>	Oregon crab apple	Apple mosaic virus (ApMV)	D	1	[36]
		Apple scab (<i>Venturia inaequalis</i>)	D	1	[36]
<i>Malus ioensis</i>	prairie crabapple	Apple mosaic virus (ApMV)	D	1	[36]
		Apple scab (<i>Venturia inaequalis</i>)	D	1	[36]
<i>Metopium toxiferum</i>	Florida poisontree	none	.	.	.
<i>Morus microphylla</i>	Texas mulberry	Bacterial leaf scorch (<i>Xylella fastidiosa</i>)	D	1	[36]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Morus rubra</i>	red mulberry	Pink hibiscus mealybug (<i>Maconellicoccus hirsutus</i>)*	I/S	1	[34]
		Bacterial leaf scorch (<i>Xylella fastidiosa</i>)	D	1	[36]
<i>Nyssa aquatica</i>	water tupelo	Pink hibiscus mealybug (<i>Maconellicoccus hirsutus</i>)*	I/S	1	[34]
		Forest tent caterpillar (<i>Malacosoma disstria</i>)	I/F	1	[1]
<i>Nyssa biflora</i>	swamp tupelo	Stem canker (<i>Fusarium solani</i>)	D	1	[36]
		Tupelo leafminer (<i>Antispila nysaefoliella</i>)	I/F	1	[1]
<i>Nyssa ogeche</i>	Ogeechee tupelo	Forest tent caterpillar (<i>Malacosoma disstria</i>)	I/F	1	[144]
		Stem canker (<i>Fusarium solani</i>)	D	1	[144]
<i>Nyssa sylvatica</i>	blackgum	Tupelo leafminer (<i>Antispila nysaefoliella</i>)	I/F	1	[1]
		Forest tent caterpillar (<i>Malacosoma disstria</i>)	I/F	1	[1]
<i>Olneya tesota</i>	desert ironwood	Stem canker (<i>Fusarium solani</i>)	D	1	[1]
		Tupelo leafminer (<i>Antispila nysaefoliella</i>)	I/F	1	[1]
<i>Osmanthus americanus</i>	devilwood	Mistletoe (<i>Phoradendron californicum</i>)	P	1	[36]
<i>Ostrya chisosensis</i>	Chisos hophornbeam	none	.	.	.
<i>Ostrya knowltonii</i>	Knowlton's hophornbeam	none	.	.	.
<i>Ostrya virginiana</i>	eastern hophornbeam	Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
<i>Oxydendrum arboreum</i>	sourwood	Black twig borer (<i>Xylosandrus compactus</i>)*	I/B	1	[145]
		Dogwood twig borer (<i>Oberea tripunctata</i>)	I/B	1	[1]
		Nectria canker (<i>Nectria ditissima</i>)	D	1	[36]
		Rhododendron stem borer (<i>Oberea myops</i>)	I/B	1	[146]
		Twig girdler (<i>Oncideres cingulata</i>)	I/B	1	[1]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Parkinsonia aculeata</i>	Jerusalem-thorn	Mistletoe (<i>Phoradendron californicum</i>)	P	1	[36]
		White mottled rot (<i>Ganoderma</i> spp.)	D	1	[36]
<i>Parkinsonia florida</i>	blue paloverde	Mistletoe (<i>Phoradendron californicum</i>)	P	1	[36]
		Palo verde root borers (<i>Derobrachus geminatus</i>)	I/B	1	[147,148]
<i>Parkinsonia texana</i> var. <i>macra</i>	border paloverde	White mottled rot (<i>Ganoderma</i> spp.)	D	1	[36]
		Mistletoe (<i>Phoradendron californicum</i>)	P	1	[36]
		White mottled rot (<i>Ganoderma</i> spp.)	D	1	[36]
<i>Persea borbonia</i>	redbay	Laurel wilt (<i>Raffaelea lauricola</i>)/redbay ambrosia beetle (<i>Xyleborus glabratus</i>)*	C	10	[34,149]
		Black twig borer (<i>Xylosandrus compactus</i>)*	I/B	1	[145]
<i>Picea breweriana</i>	Brewer spruce	Lobate lac scale (<i>Paratachardina pseudolobata</i>)*	I/S	1	[97]
		Spruce beetle (<i>Dendroctonus rufipennis</i>)	I/B	5	[37]
		Heterobasidion root disease (<i>Heterobasidion</i> spp.)	D	3	[1]
		White pine weevil (<i>Pissodes strobi</i>)	I/B	3	[37]
		Dwarf mistletoe (<i>Arceuthobium</i> spp.)	P	1	[1,36]
		Spruce beetle (<i>Dendroctonus rufipennis</i>)	I/B	8	[1,37]
<i>Picea engelmannii</i>	Engelmann spruce	Western spruce budworm (<i>Choristoneura freemani</i>)	I/F	5	[1]
		Armillaria root disease (<i>Armillaria</i> spp.)	D	3	[35,150]
		Red root and butt rot (<i>Onnia tomentosa</i>)	D	3	[12]
		Western spruce dwarf mistletoe (<i>Arceuthobium microcarpum</i>)	P	3	[1]
		Spruce beetle (<i>Dendroctonus rufipennis</i>)	I/B	5	[37]
<i>Picea glauca</i>	white spruce	Western spruce budworm (<i>Choristoneura freemani</i>)	I/F	5	[37]
		Eastern dwarf mistletoe (<i>Arceuthobium pusillum</i>)	P	3	[1]
		White pine weevil (<i>Pissodes strobi</i>)	I/B	3	[37]
		Red root and butt rot (<i>Onnia tomentosa</i>)	D	1	[36]
		Spruce beetle (<i>Dendroctonus rufipennis</i>)	I/B	5	[1]
<i>Picea mariana</i>	black spruce				

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Picea pungens</i>	blue spruce	Spruce budworm (<i>Choristoneura fumiferana</i>)	I/F	5	[37]
		Eastern dwarf mistletoe (<i>Arceuthobium pusillum</i>)	P	3	[1,36]
		Red root and butt rot (<i>Onnia tomentosa</i>)	D	1	[36]
		Western spruce budworm (<i>Choristoneura freemani</i>)	I/F	5	[1,37]
		Red root and butt rot (<i>Onnia tomentosa</i>)	D	3	[36]
		Blue spruce engraver (<i>Ips hunteri</i>)	I/B	1	[37]
<i>Picea rubens</i>	red spruce	Cytospora canker of spruce (<i>Cytospora kunzei</i>)	D	1	[13,36]
		Western spruce dwarf mistletoe (<i>Arceuthobium microcarpum</i>)	P	1	[1]
		Spruce budworm (<i>Choristoneura fumiferana</i>)	I/F	5	[1]
		Eastern dwarf mistletoe (<i>Arceuthobium pusillum</i>)	P	3	[36]
		Spruce beetle (<i>Dendroctonus rufipennis</i>)	I/B	3	[37]
		Red-belted fungus (<i>Fomitopsis pinicola</i>)	D	1	[36]
<i>Picea sitchensis</i>	Sitka spruce	Spruce beetle (<i>Dendroctonus rufipennis</i>)	I/B	5	[1]
		Heterobasidion root disease (<i>Heterobasidion</i> spp.)	D	3	[36]
		Laminated root rot (<i>Phellinus weiri</i>)	D	3	[36]
		White pine weevil (<i>Pissodes strobi</i>)	I/B	3	[1]
		Schweinitzii root and butt rot (<i>Phaeolus schweinitzii</i>)	D	1	[36]
		Mountain pine beetle (<i>Dendroctonus ponderosae</i>)	I/B	8	[1,37]
<i>Pinus albicaulis</i>	whitebark pine	White pine blister rust (<i>Cronartium ribicola</i>)*	D	8	[1,36]
		Armillaria root disease (<i>Armillaria</i> spp.)	D	3	[13,35,36]
		Limber pine dwarf mistletoe (<i>Arceuthobium cyanocarpum</i>)	P	3	[1,36]
		Foliage diseases (<i>Bifusella</i> spp.)	D	1	[151]
		Mountain pine beetle (<i>Dendroctonus ponderosae</i>)	I/B	8	[152]
		White pine blister rust (<i>Cronartium ribicola</i>)*	D	5	[7,36]
<i>Pinus aristata</i>	Rocky Mountain bristlecone pine	Limber pine dwarf mistletoe (<i>Arceuthobium cyanocarpum</i>)	P	3	[36]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Pinus arizonica</i>	Arizona pine	Pine engraver beetles (<i>Ips</i> spp.)	I/B	3	[35,37]
		Twig beetle (<i>Pityophthorus boycei</i>)	I/B	1	[37]
		Armillaria root disease (<i>Armillaria</i> spp.)	D	3	[13,35,36]
		Dwarf mistletoe (<i>Arceuthobium</i> spp.)	P	3	[36]
		Limb rust (<i>Cronartium arizonicum</i>)	D	3	[36]
		Red root and butt rot (<i>Onnia tomentosa</i>)	D	3	[35,36]
<i>Pinus attenuata</i>	Knobcone pine	Gall rust (<i>Cronartium quercuum</i>)	D	1	[36]
		Pine engraver beetles (<i>Ips</i> spp.)	I/B	5	[37,153]
		Dwarf mistletoe (<i>Arceuthobium</i> spp.)	P	3	[36]
		Pitch canker (<i>Fusarium circinatum</i>)*	D	3	[36]
		Elytroderma needle cast (<i>Elytroderma deformans</i>)	D	1	[36]
		Western gall rust (<i>Endocronartium harknessii</i>)	D	1	[36]
<i>Pinus balfouriana</i>	foxtail pine	Mountain pine beetle (<i>Dendroctonus ponderosae</i>)	I/B	8	[7]
		White pine blister rust (<i>Cronartium ribicola</i>)*	D	8	[7,36]
		Limber pine dwarf mistletoe (<i>Arceuthobium cyanocarpum</i>)	P	3	[7,36]
		Jack pine budworm (<i>Choristoneura pinus</i>)	I/F	5	[7,37]
		Lodgepole pine dwarf mistletoe (<i>Arceuthobium americanum</i>)	P	3	[36]
		North American pine engraver (<i>Ips pini</i>)	I/B	3	[35,37]
<i>Pinus banksiana</i>	jack pine	White pine weevil (<i>Pissodes strobi</i>)	I/B	3	[37]
		Comandra blister rust (<i>Cronartium comandrae</i>)	D	1	[36]
		Pinyon dwarf mistletoe (<i>Arceuthobium divaricatum</i>)	P	3	[7,36]
		Pinyon blister rust (<i>Cronartium occidentale</i>)	D	1	[7,36]
		Pinyon needle scale (<i>Matsucoccus acalyptus</i>)	I/S	1	[37]
		Southern pine beetle (<i>Dendroctonus frontalis</i>)	I/B	3	[154]
<i>Pinus clausa</i>	sand pine	Armillaria root disease (<i>Armillaria</i> spp.)	D	1	[1,36]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Pinus contorta</i>	lodgepole pine	Pine engraver beetles (<i>Ips</i> spp.)	I/B	1	[1,37,155]
		Red root and butt rot (<i>Onnia tomentosa</i>)	D	1	[36]
		Virginia pine sawfly (<i>Neodiprion pratti</i>)	I/F	1	[1]
		Mountain pine beetle (<i>Dendroctonus ponderosae</i>)	I/B	8	[1,37]
		Armillaria root disease (<i>Armillaria</i> spp.)	D	3	[1]
		Comandra blister rust (<i>Cronartium comandrae</i>)	D	3	[1,36]
		Lodgepole pine dwarf mistletoe (<i>Arceuthobium americanum</i>)	P	3	[1,36]
<i>Pinus coulteri</i>	Coulter pine	Pine engraver beetles (<i>Ips</i> spp.)	I/B	3	[1,156]
		Western pine beetle (<i>Dendroctonus brevicomis</i>)	I/B	8	[37]
		Pine engraver beetles (<i>Ips</i> spp.)	I/B	5	[37,153]
		Dwarf mistletoe (<i>Arceuthobium</i> spp.)	P	3	[36]
		Elytroderma needle cast (<i>Elytroderma deformans</i>)	D	1	[36]
		Pitch canker (<i>Fusarium circinatum</i>)*	D	1	[34]
		Pinyon dwarf mistletoe (<i>Arceuthobium divaricatum</i>)	P	3	[36]
<i>Pinus discolor</i>	border pinyon	Pinyon blister rust (<i>Cronartium occidentale</i>)	D	1	[36]
		Pinyon needle scale (<i>Matsucoccus acalyptus</i>)	I/S	1	[37]
		Phytophthora root rot (<i>Phytophthora cinnamomi</i>)*	D	5	[1,36]
		Southern pine beetle (<i>Dendroctonus frontalis</i>)	I/B	5	[154]
		Pine engraver beetles (<i>Ips</i> spp.)	I/B	1	[1,37,155]
		Pitch canker (<i>Fusarium circinatum</i>)*	D	1	[36]
		Redheaded pine sawfly (<i>Neodiprion lecontei</i>)	I/F	1	[1]
<i>Pinus edulis</i>	common pinyon	Pinyon ips (<i>Ips confusus</i>)	I/B	8	[12,153]
		Pinyon blister rust (<i>Cronartium occidentale</i>)	D	3	[1,36]
		Pinyon dwarf mistletoto (<i>Arceuthobium divaricatum</i>)	P	3	[36]
		Pinyon needle scale (<i>Matsucoccus acalyptus</i>)	I/S	1	[1,37]
		Pinyon sawfly (<i>Neodiprion edulicolus</i>)	I/F	1	[1,37]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Pinus elliottii</i>	slash pine	Fusiform rust (<i>Cronartium quercuum</i> f. sp. <i>fusiforme</i>)	D	3	[1,36]
		Heterobasidion root disease (<i>Heterobasidion</i> spp.)	D	1	[1,36]
		Pine engraver beetles (<i>Ips</i> spp.)	I/B	1	[1,37,155]
		Pitch canker (<i>Fusarium circinatum</i>)*	D	1	[36]
		Southern pine beetle (<i>Dendroctonus frontalis</i>)	I/B	1	[154]
<i>Pinus engelmannii</i>	Apache pine	Dwarf mistletoe (<i>Arceuthobium</i> spp.)	P	3	[36]
		Limb rust (<i>Cronartium</i> sp.)	D	3	[36]
		Southern pine beetle (<i>Dendroctonus frontalis</i>)	I/B	3	[37]
<i>Pinus flexilis</i>	limber pine	Mountain pine beetle (<i>Dendroctonus ponderosae</i>)	I/B	8	[1,37]
		White pine blister rust (<i>Cronartium ribicola</i>)*	D	8	[1,36]
		Limber pine dwarf mistletoe (<i>Arceuthobium cyanocarpum</i>)	P	3	[1,36]
		Pine engraver beetles (<i>Ips</i> spp.)	I/B	3	[35,37]
		Twig beetles (<i>Pityophthorus</i> spp.)	I/B	1	[37]
		Southern pine beetle (<i>Dendroctonus frontalis</i>)	I/B	3	[154]
<i>Pinus glabra</i>	spruce pine	Ganoderma root and butt rot (<i>Ganoderma meredithae</i>)	D	1	[36]
		Pine engraver beetles (<i>Ips</i> spp.)	I/B	1	[37,155]
		Jeffrey pine beetle (<i>Dendroctonus jeffreyi</i>)	I/B	5	[1,37]
<i>Pinus jeffreyi</i>	Jeffrey pine	Pine engraver beetles (<i>Ips</i> spp.)	I/B	5	[1,37]
		Black-stain root disease (<i>Leptographium wageneri</i>)	D	3	[36]
		Elytroderma needle cast (<i>Elytroderma deformans</i>)	D	3	[1]
		Western dwarf mistletoe (<i>Arceuthobium campylopodum</i>)	P	3	[1,36]
		Mountain pine beetle (<i>Dendroctonus ponderosae</i>)	I/B	8	[1,37]
<i>Pinus lambertiana</i>	sugar pine	White pine blister rust (<i>Cronartium ribicola</i>)*	D	8	[1,36]
		Dwarf mistletoe (<i>Arceuthobium californicum</i>)	P	3	[1]
		Heterobasidion root disease (<i>Heterobasidion</i> spp.)	D	3	[1]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Pinus leiophylla</i>	Chihuahuan pine	White pine needle cast (<i>Lophodermella arcuata</i>)	D	1	[1,36]
		Armillaria root disease (<i>Armillaria</i> spp.)	D	3	[13,35,36]
		Dwarf mistletoe (<i>Arceuthobium gillii</i>)	P	3	[36]
		Red root and butt rot (<i>Onnia tomentosa</i>)	D	3	[35,36]
<i>Pinus longaeva</i>	Great Basin bristlecone pine	Southern pine beetle (<i>Dendroctonus frontalis</i>)	I/B	3	[37]
		White pine blister rust (<i>Cronartium ribicola</i>)*	D	8	[34]
		Western dwarf mistletoe (<i>Arceuthobium campylopodium</i>)	P	3	[7]
		Brown spot needle blight (<i>Mycosphaerella dearnessii</i>)	D	1	[36]
<i>Pinus monophylla</i>	singleleaf pinyon	Mountain pine beetle (<i>Dendroctonus ponderosae</i>)	I/B	1	[7]
		Pinyon ips (<i>Ips confusus</i>)	I/B	8	[1,37,153]
		Pinyon dwarf mistletoe (<i>Arceuthobium divaricatum</i>)	P	3	[1]
		Pinyon blister rust (<i>Cronartium occidentale</i>)	D	1	[1]
<i>Pinus monophylla</i> var. <i>fallax</i>	Arizona pinyon pine	Pinyon needle scale (<i>Matsucoccus acalyptus</i>)	I/S	1	[1]
		Pinyon sawfly (<i>Neodiprion edulicolus</i>)	I/F	1	[1]
		Pinyon ips (<i>Ips confusus</i>)	I/B	8	[1,37,153]
		Pinyon dwarf mistletoe (<i>Arceuthobium divaricatum</i>)	P	3	[1]
<i>Pinus monticola</i>	western white pine	Pinyon blister rust (<i>Cronartium occidentale</i>)	D	1	[1]
		Pinyon needle scale (<i>Matsucoccus acalyptus</i>)	I/S	1	[1]
		Pinyon sawfly (<i>Neodiprion edulicolus</i>)	I/F	1	[1]
		Mountain pine beetle (<i>Dendroctonus ponderosae</i>)	I/B	8	[1,37]
<i>Pinus muricata</i>	bishop pine	White pine blister rust (<i>Cronartium ribicola</i>)*	D	8	[1,36]
		Armillaria root disease (<i>Armillaria</i> spp.)	D	3	[1,36]
		Dwarf mistletoto (<i>Arceuthobium monticola</i>)	P	3	[36]
		Red ring rot (<i>Phellinus pini</i>)	D	1	[1,36]
		Pitch canker (<i>Fusarium circinatum</i>)*	D	5	[34,36]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Pinus palustris</i>	longleaf pine	Coastal dwarf mistletoe (<i>Arceuthobium littorum</i>)	P	3	[36]
		Monterey pine weevil (<i>Pissodes radiatae</i>)*	I/B	1	[37]
		Fusiform rust (<i>Cronartium quercuum</i> f. sp. <i>fusiforme</i>)	D	3	[1,36]
		Heterobasidion root disease (<i>Heterobasidion</i> spp.)	D	1	[1]
		Pine engraver beetles (<i>Ips</i> spp.)	I/B	1	[1,37,155]
		Pitch canker (<i>Fusarium circinatum</i>)*	D	1	[36]
<i>Pinus ponderosa</i>	ponderosa pine	Southern pine beetle (<i>Dendroctonus frontalis</i>)	I/B	1	[1]
		Mountain pine beetle (<i>Dendroctonus ponderosae</i>)	I/B	8	[37]
		Pine engraver beetles (<i>Ips</i> spp.)	I/B	8	[1,37]
		Armillaria root disease (<i>Armillaria</i> spp.)	D	5	[1,36]
		Dwarf mistletoe (<i>Arceuthobium</i> spp.)	P	3	[1,36]
		Comandra blister rust (<i>Cronartium comandrae</i>)	D	1	[36]
<i>Pinus ponderosa</i> var. <i>washoensis</i>	Washoe pine	Mountain pine beetle (<i>Dendroctonus ponderosae</i>)	I/B	5	[37]
		Western dwarf mistletoe (<i>Arceuthobium campylopodum</i>)	P	3	[1,36]
		Armillaria root disease (<i>Armillaria</i> spp.)	D	1	[1,36]
		Comandra blister rust (<i>Cronartium comandrae</i>)	D	1	[35,36]
		Elytroderma needle cast (<i>Elytroderma deformans</i>)	D	1	[1,36]
		Southern pine beetle (<i>Dendroctonus frontalis</i>)	I/B	3	[1]
<i>Pinus pungens</i>	Table Mountain pine	Comandra blister rust (<i>Cronartium comandrae</i>)	D	1	[1]
		European pine sawfly (<i>Neodiprion sertifer</i>)*	I/F	1	[1]
		Pine engraver beetles (<i>Ips</i> spp.)	I/B	1	[37,152]
		Schweinitzii root and butt rot (<i>Phaeolus schweinitzii</i>)	D	1	[1]
		Pinyon dwarf mistletoe (<i>Arceuthobium divaricatum</i>)	P	3	[7,36]
		Pinyon ips (<i>Ips confusus</i>)	I/B	3	[37,153]
<i>Pinus quadrifolia</i>	four-leaf pine	Pinon twig beetle (<i>Pityophthorus punctifrons</i>)	I/B	1	[37]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Pinus radiata</i>	Monterey pine	Pine engraver beetles (<i>Ips</i> spp.)	I/B	5	[1,37]
		Pitch canker (<i>Fusarium circinatum</i>)*	D	5	[34,36]
		Western gall rust (<i>Endocronartium harknessii</i>)	D	3	[1,36]
		Monterey pine weevil (<i>Pissodes radiatae</i>)*	I/B	1	[1]
<i>Pinus remota</i>	papershell pinyon pine	Pinyon dwarf mistletoe (<i>Arceuthobium divaricatum</i>)	P	3	[1,36]
		Pinyon blister rust (<i>Cronartium occidentale</i>)	D	1	[1,36]
		Pinyon needle scale (<i>Matsucoccus acalyptus</i>)	I/S	1	[37]
<i>Pinus resinosa</i>	red pine	Red pine scale (<i>Matsucoccus resinosae</i>)*	I/S	3	[1,37]
		Heterobasidion root disease (<i>Heterobasidion</i> spp.)	D	1	[1,36]
		Sawfly (<i>Neodiprion</i> spp.)	I/F	1	[1,37]
		Scleroderris canker (<i>Gremmeniella abietina</i>)*	D	1	[1,36]
		White pine weevil (<i>Pissodes strobi</i>)	I/B	1	[37]
<i>Pinus rigida</i>	pitch pine	Gall rust (<i>Cronartium quercuum</i>)	D	3	[36]
		Southern pine beetle (<i>Dendroctonus frontalis</i>)	I/B	3	[1,37]
		Fusiform rust (<i>Cronartium quercuum</i> f. sp. <i>fusiforme</i>)	D	1	[1,36]
		Heterobasidion root disease (<i>Heterobasidion</i> spp.)	D	1	[1]
		Sawfly (<i>Neodiprion</i> spp.)	I/F	1	[37]
<i>Pinus sabiniana</i>	gray pine	Bark beetle (<i>Orthotomicus spinifer</i>)	I/B	3	[1]
		Dwarf mistletoe (<i>Arceuthobium occidentale</i>)	P	3	[1,36]
		Pitch canker (<i>Fusarium circinatum</i>)*	D	1	[36,157]
		Western gall rust (<i>Endocronartium harknessii</i>)	D	1	[1]
<i>Pinus serotina</i>	pond pine	Gall rust (<i>Cronartium quercuum</i>)	D	3	[36]
		Southern pine beetle (<i>Dendroctonus frontalis</i>)	I/B	3	[154]
		Fusiform rust (<i>Cronartium quercuum</i> f. sp. <i>fusiforme</i>)	D	1	[1,36]
		Pine engraver beetles (<i>Ips</i> spp.)	I/B	1	[1,37,152]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Pinus strobus</i>	white pine	Red ring rot (<i>Phellinus pini</i>)	D	1	[1]
		White pine blister rust (<i>Cronartium ribicola</i>)*	D	8	[36]
		Pine engraver beetles (<i>Ips</i> spp.)	I/B	5	[35,37]
		Roundheaded pine beetle (<i>Dendroctonus adjunctus</i>)	I/B	5	[37]
		Apache dwarf mistletoe (<i>Arceuthobium apachecum</i>)	P	3	[36]
<i>Pinus taeda</i>	loblolly pine	Armillaria root disease (<i>Armillaria</i> spp.)	D	3	[13,35,36]
		White pine blister rust (<i>Cronartium ribicola</i>)*	D	5	[1,36]
		Canadian pine scale (<i>Matsucoccus macrocicatrices</i>)/ <i>Caliciopsis pinea</i> complex	C	3	[158]
		Heterobasidion root disease (<i>Heterobasidion</i> spp.)	D	1	[1,36]
		Pine shoot beetle (<i>Tomicus piniperda</i>)*	I/B	1	[37]
<i>Pinus torreyana</i>	Torrey pine	White pine weevil (<i>Pissodes strobi</i>)	I/B	1	[1,37]
		Southern pine beetle (<i>Dendroctonus frontalis</i>)	I/B	5	[1,37]
		Fusiform rust (<i>Cronartium quercuum</i> f. sp. <i>fusiforme</i>)	D	3	[36]
		Pine engraver beetles (<i>Ips</i> spp.)	I/B	3	[1,37,155]
		Heterobasidion root disease (<i>Heterobasidion</i> spp.)	D	1	[1,36]
<i>Pinus virginiana</i>	Virginia pine	Sawfly (<i>Neodiprion</i> spp.)	I/F	1	[35,37]
		California fivespined ips (<i>Ips paraconfusus</i>)	I/B	5	[7]
		Dwarf mistletoe (<i>Arceuthobium occidentale</i>)	P	3	[1,36]
		Pitch canker (<i>Fusarium circinatum</i>)*	D	3	[36,157]
		Bark beetle (<i>Ips spinifer</i>)	I/B	1	[1]
<i>Pinus strobiformis</i>	southwestern white pine	Western gall rust (<i>Endocronartium harknessii</i>)	D	1	[1]
		Pitch canker (<i>Fusarium circinatum</i>)*	D	3	[1,36]
		Southern pine beetle (<i>Dendroctonus frontalis</i>)	I/B	3	[1]
		Gall rust of Virginia pine (<i>Cronartium quercuum</i> f. sp. <i>virginianae</i>)	D	1	[1,36]
		Red ring rot (<i>Phellinus pini</i>)	D	1	[1]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Piscidia piscipula</i>	fishpoison tree	Sawfly (<i>Neodiprion</i> spp.)	I/F	1	[1,35]
<i>Pistacia mexicana</i>	Texas pistache	Psyllid (<i>Euphalerus nidifex</i>)	I/F	1	[159]
<i>Planera aquatica</i>	water-elm	none	.	.	.
<i>Platanus occidentalis</i>	American sycamore	Dutch elm disease (<i>Ophiostoma novo-ulmi</i>)*	D	5	[160,161]
		Sycamore canker (<i>Ceratocystis fimbriata</i> f. <i>platani</i>)	D	5	[1,36]
		Anthracnose (<i>Apiognomonia veneta</i>)	D	1	[1,36]
		Bacterial leaf scorch (<i>Xylella fastidiosa</i>)	D	1	[36]
		Sycamore lace bug (<i>Corythucha ciliata</i>)	I/F	1	[1]
<i>Platanus racemosa</i>	California sycamore	Polyphagous shot hole borer (<i>Euwallacea</i> spp.)*	I/B	8	[38,162]
		Anthracnose (<i>Apiognomonia veneta</i>)	D	1	[36]
		Sycamore borer (<i>Synanthedon resplendens</i>)	I/B	1	[163]
<i>Platanus wrightii</i>	Arizona sycamore	Anthracnose (<i>Apiognomonia veneta</i>)	D	1	[36]
		Eastern mistletoe (<i>Phoradendron leucarpum</i>)	P	1	[164]
		Oak bracket (<i>Inonotus dryadeus</i>)	D	1	[36]
<i>Populus angustifolia</i>	narrowleaf cottonwood	Poplar borer (<i>Saperda calcarata</i>)	I/B	3	[37]
		Cottonwood leaf beetle (<i>Chrysomela scripta</i>)	I/F	1	[165]
		Poplar rust (<i>Melampsora abietis-canadensis</i>)	D	1	[8]
		Poplar-and-willow borer (<i>Cryptorhynchus lapathi</i>)*	I/B	1	[34]
<i>Populus balsamifera</i>	balsam poplar	Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Forest tent caterpillar (<i>Malacosoma disstria</i>)	I/F	1	[12]
		Oystershell scale (<i>Lepidosaphes ulmi</i>)*	I/S	1	[34]
		Poplar borer (<i>Saperda calcarata</i>)	I/B	1	[1,37]
		Poplar rust (<i>Melampsora abietis-canadensis</i>)	D	1	[1,36]
<i>Populus balsamifera</i> ssp. <i>trichocarpa</i>	black cottonwood	Cottonwood leaf beetle (<i>Chrysomela scripta</i>)	I/F	1	[165]
		Cytospora canker (<i>Cytospora chrysosperma</i>)	D	1	[13]
		Oystershell scale (<i>Lepidosaphes ulmi</i>)*	I/S	1	[1]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Populus deltoides</i>	eastern cottonwood	Poplar rust (<i>Melampsora occidentalis</i>)	D	1	[36]
		Poplar-and-willow borer (<i>Cryptorhynchus lapathi</i>)*	I/B	1	[1]
		Artist's conk (<i>Ganoderma applanatum</i>)	D	1	[36]
		Cottonwood borer (<i>Plectrodera scalator</i>)	I/B	1	[1]
		Forest tent caterpillar (<i>Malacosoma disstria</i>)	I/F	1	[12]
		Poplar borer (<i>Saperda calcarata</i>)	I/B	1	[1,37]
<i>Populus deltoides</i> ssp. <i>monilifera</i>	plains cottonwood	Poplar rust (<i>Melampsora abietis-canadensis</i>)	D	1	[1,36]
		Artist's conk (<i>Ganoderma applanatum</i>)	D	1	[36]
		Cottonwood borer (<i>Plectrodera scalator</i>)	I/B	1	[1]
		Forest tent caterpillar (<i>Malacosoma disstria</i>)	I/F	1	[12]
		Poplar borer (<i>Saperda calcarata</i>)	I/B	1	[1,37]
		Poplar rust (<i>Melampsora abietis-canadensis</i>)	D	1	[36]
<i>Populus fremontii</i>	Fremont cottonwood	Canker (<i>Cryptosphaeria pullmanensis</i>)	D	1	[166]
		Cottonwood leaf beetle (<i>Chrysomela scripta</i>)	I/F	1	[165]
		Sooty canker (<i>Fusicoccum mangiferum</i>)	D	1	[36]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Hypoxylon canker of aspen (<i>Entoleuca mammata</i>)	D	3	[13,36]
		Artist's conk (<i>Ganoderma applanatum</i>)	D	1	[1]
<i>Populus grandidentata</i>	bigtooth aspen	Forest tent caterpillar (<i>Malacosoma disstria</i>)	I/F	1	[167]
		Poplar borer (<i>Saperda calcarata</i>)	I/B	1	[1,37]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Cottonwood leaf beetle (<i>Chrysomela scripta</i>)	I/F	1	[165]
		Forest tent caterpillar (<i>Malacosoma disstria</i>)	I/F	1	[35,37]
		Melampsora leaf rust (<i>Melampsora medusae</i>)	D	1	[1]
<i>Populus heterophylla</i>	swamp cottonwood	Poplar borer (<i>Saperda calcarata</i>)	I/B	1	[1,37]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Populus tremuloides</i>	quaking aspen	Bronze poplar borer (<i>Agrilus liragus</i>)	I/B	3	[1,168-170]
		Cytospora canker (<i>Cytospora chrysosperma</i>)	D	3	[36]
		Forest tent caterpillar (<i>Malacosoma disstria</i>)	I/F	3	[1,12]
		Poplar borer (<i>Saperda calcarata</i>)	I/B	3	[37]
		Sooty bark canker (<i>Encoelia pruinosa</i>)	D	3	[1,171]
<i>Prosopis glandulosa</i>	honey mesquite	Borer (<i>Megacyllene antennata</i>)	I/B	1	[37]
		Mistletoe (<i>Phoradendron californicum</i>)	P	1	[36]
		White mottled rot (<i>Ganoderma</i> spp.)	D	1	[36]
<i>Prosopis pubescens</i>	screwbean mesquite	Borer (<i>Megacyllene antennata</i>)	I/B	1	[37]
		Mistletoe (<i>Phoradendron californicum</i>)	P	1	[36]
		White mottled rot (<i>Ganoderma</i> spp.)	D	1	[36]
<i>Prosopis velutina</i>	velvet mesquite	Borer (<i>Megacyllene antennata</i>)	I/B	1	[37]
		Mistletoe (<i>Phoradendron californicum</i>)	P	1	[36]
		White mottled rot (<i>Ganoderma</i> spp.)	D	1	[36]
<i>Prunus alleghaniensis</i>	Allegheny plum	Bacterial spot (<i>Xanthomonas campestris</i> pv. <i>pruni</i>)	D	1	[36]
		Black knot (<i>Apiosporina morbosa</i>)	D	1	[36]
		Cherry leaf spot (<i>Blumeriella jaapii</i>)	D	1	[36]
		Eastern tent caterpillar (<i>Malacosoma americanum</i>)	I/F	1	[35,37]
		Prunus X-disease (<i>Phytoplasma</i>)	D	5	[36]
<i>Prunus americana</i>	American plum	Bacterial spot (<i>Xanthomonas campestris</i> pv. <i>pruni</i>)	D	1	[36]
		Black knot (<i>Apiosporina morbosa</i>)	D	1	[36]
		Cherry leaf spot (<i>Blumeriella jaapii</i>)	D	1	[36]
		Tent caterpillar (<i>Malacosoma</i> spp.)	I/F	1	[35,37]
		Bacterial spot (<i>Xanthomonas campestris</i> pv. <i>pruni</i>)	D	1	[36]
<i>Prunus angustifolia</i>	Chickasaw plum	Black knot (<i>Apiosporina morbosa</i>)	D	1	[36]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Prunus caroliniana</i>	Carolina laurelcherry	Cherry leaf spot (<i>Blumeriella jaapii</i>)	D	1	[36]
		Tent caterpillar (<i>Malacosoma</i> spp.)	I/F	1	[35,37]
		Bacterial spot (<i>Xanthomonas campestris</i> pv. <i>pruni</i>)	D	1	[36]
		Black knot (<i>Apiosporina morbosa</i>)	D	1	[36]
		Cherry leaf spot (<i>Blumeriella jaapii</i>)	D	1	[36]
<i>Prunus emarginata</i>	bitter cherry	Eastern tent caterpillar (<i>Malacosoma americanum</i>)	I/F	1	[35,37]
		Black knot (<i>Apiosporina morbosa</i>)	D	3	[36]
		Cherry leaf spot (<i>Blumeriella jaapii</i>)	D	1	[36]
		Leaf curl and witches broom (<i>Taphrina wiesneri</i>)	D	1	[36]
		Tent caterpillar (<i>Malacosoma</i> spp.)	I/F	1	[35,37]
<i>Prunus hortulana</i>	hotulan plum	Bacterial spot (<i>Xanthomonas campestris</i> pv. <i>pruni</i>)	D	1	[36]
		Black knot (<i>Apiosporina morbosa</i>)	D	1	[36]
		Cherry leaf spot (<i>Blumeriella jaapii</i>)	D	1	[36]
		Eastern tent caterpillar (<i>Malacosoma americanum</i>)	I/F	1	[35,37]
		Black knot (<i>Apiosporina morbosa</i>)	D	1	[36]
<i>Prunus mexicana</i>	Mexicana plum	Cherry leaf spot (<i>Blumeriella jaapii</i>)	D	1	[36]
		Eastern tent caterpillar (<i>Malacosoma americanum</i>)	I/F	1	[35,37]
		Black knot (<i>Apiosporina morbosa</i>)	D	1	[36]
		Cherry leaf spot (<i>Blumeriella jaapii</i>)	D	1	[36]
		Eastern tent caterpillar (<i>Malacosoma americanum</i>)	I/F	1	[35,37]
<i>Prunus nigra</i>	Canada plum	Black knot (<i>Apiosporina morbosa</i>)	D	1	[36]
		Cherry leaf spot (<i>Blumeriella jaapii</i>)	D	1	[36]
		Eastern tent caterpillar (<i>Malacosoma americanum</i>)	I/F	1	[35,37]
		Black knot (<i>Apiosporina morbosa</i>)	D	1	[36]
		Cherry leaf spot (<i>Blumeriella jaapii</i>)	D	1	[36]
<i>Prunus pensylvanica</i>	pin cherry	Eastern tent caterpillar (<i>Malacosoma americanum</i>)	I/F	1	[35,37]
		Black knot (<i>Apiosporina morbosa</i>)	D	1	[1,36]
		Cherry leaf spot (<i>Blumeriella jaapii</i>)	D	1	[36]
		Leaf curl and witches broom (<i>Taphrina wiesneri</i>)	D	1	[36]
		Tent caterpillar (<i>Malacosoma</i> spp.)	I/F	1	[1,35,37]
<i>Prunus serotina</i>	black cherry	Bacterial spot (<i>Xanthomonas campestris</i> pv. <i>pruni</i>)	D	1	[36]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Prunus umbellata</i>	flatwoods plum	Black knot (<i>Apiosporina morbosa</i>)	D	1	[1,36]
		Cherry leaf spot (<i>Blumeriella jaapii</i>)	D	1	[36]
		Eastern tent caterpillar (<i>Malacosoma americanum</i>)	I/F	1	[1,35,37]
		Black knot (<i>Apiosporina morbosa</i>)	D	1	[36]
		Cherry leaf spot (<i>Blumeriella jaapii</i>)	D	1	[36]
		Eastern tent caterpillar (<i>Malacosoma americanum</i>)	I/F	1	[35,37]
<i>Prunus virginiana</i>	chokecherry	Prunus X-disease (<i>Phytoplasma</i>)	D	5	[36,172,173]
		Black knot (<i>Apiosporina morbosa</i>)	D	3	[36]
		Tent caterpillar (<i>Malacosoma</i> spp.)	I/F	3	[35,37]
		Cherry leaf spot (<i>Blumeriella jaapii</i>)	D	1	[36]
<i>Pseudotsuga macrocarpa</i>	bigcone Douglas-fir	Flatheaded fir borer (<i>Melanophila drummondi</i>)	I/B	3	[174]
		Laminated root rot (<i>Phellinus sulphurascens</i>)	D	3	[20]
		California flatheaded borer (<i>Melanophila californica</i>)	I/B	1	[1]
		Douglas-fir beetle (<i>Dendroctonus pseudotsugae</i>)	I/B	8	[37]
<i>Pseudotsuga menziesii</i>	Douglas-fir	Laminated root rot (<i>Phellinus sulphurascens</i>)	D	8	[20,36]
		Armillaria root disease (<i>Armillaria</i> spp.)	D	5	[36]
		Douglas-fir tussock moth (<i>Orgyia pseudotsugata</i>)	I/F	5	[1]
		Western spruce budworm (<i>Choristoneura freemani</i>)	I/F	5	[37]
<i>Psorothamnus spinosus</i>	smokethorn	none	.	.	.
<i>Quercus acerifolia</i>	mapleleaf oak	Oak wilt (<i>Bretziella fagacearum</i>)*	D	5	[13,34,175]
<i>Quercus agrifolia</i>	California live oak	Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Hypoxylon canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
		Oak wilt (<i>Bretziella fagacearum</i>)*	D	8	[34]
		Sudden oak death (<i>Phytophthora ramorum</i>)*	D	8	[36]
		Goldspotted oak borer (<i>Agrilus auroguttatus</i>)	I/B	5	[176]
		Oak pit scale (<i>Asterolecanium</i> spp.)*	I/S	3	[177]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Quercus alba</i>	white oak	Hypoxylon canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
		Gypsy moth (<i>Lymantria dispar</i> <i>dispar</i>)*	I/F	3	[34]
		Oak wilt (<i>Bretziella fagacearum</i>)*	D	3	[1,34,175]
		Twolined chestnut borer (<i>Agrilus bilineatus</i>)	I/B	3	[1,37]
		Armillaria root disease (<i>Armillaria</i> spp.)	D	1	[36]
<i>Quercus arizonica</i>	Arizona white oak	Hypoxylon canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
		Oak wilt (<i>Bretziella fagacearum</i>)*	D	5	[34]
		Armillaria root disease (<i>Armillaria</i> spp.)	D	1	[36]
		Canker rot of oak (<i>Inonotus andersonii</i>)	D	1	[36]
		Hypoxylon canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
<i>Quercus arkansana</i>	Arkansas oak	Oak wilt (<i>Bretziella fagacearum</i>)*	D	8	[13,34,175]
		Gypsy moth (<i>Lymantria dispar</i> <i>dispar</i>)*	I/F	3	[34]
		Twolined chestnut borer (<i>Agrilus bilineatus</i>)	I/B	3	[37]
		Armillaria root disease (<i>Armillaria</i> spp.)	D	1	[36]
		Hypoxylon canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
<i>Quercus bicolor</i>	swamp white oak	Oak wilt (<i>Bretziella fagacearum</i>)*	D	5	[1,34,175]
		Gypsy moth (<i>Lymantria dispar</i> <i>dispar</i>)*	I/F	3	[34]
		Twolined chestnut borer (<i>Agrilus bilineatus</i>)	I/B	3	[37]
		Armillaria root disease (<i>Armillaria</i> spp.)	D	1	[1,36]
		Hypoxylon canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
<i>Quercus boyntonii</i>	Boynton oak	Oak wilt (<i>Bretziella fagacearum</i>)*	D	8	[13,34,175]
		Gypsy moth (<i>Lymantria dispar</i> <i>dispar</i>)*	I/F	3	[34]
		Twolined chestnut borer (<i>Agrilus bilineatus</i>)	I/B	3	[37]
		Armillaria root disease (<i>Armillaria</i> spp.)	D	1	[36]
		Hypoxylon canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Quercus buckleyi</i>	Buckley oak	Oak wilt (<i>Bretziella fagacearum</i>)*	D	8	[34,175]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Twolined chestnut borer (<i>Agrilus bilineatus</i>)	I/B	3	[37]
		Armillaria root disease (<i>Armillaria</i> spp.)	D	1	[36]
		Hypoxylon canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
<i>Quercus chrysolepis</i>	canyon live oak	Oak wilt (<i>Bretziella fagacearum</i>)*	D	8	[34]
		Sudden oak death (<i>Phytophthora ramorum</i>)*	D	8	[117]
		Oak pit scale (<i>Asterolecanium</i> spp.)*	I/S	3	[178]
		Goldspotted oak borer (<i>Agrilus auroguttatus</i>)	I/B	1	[176]
		Hypoxylon canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36,178]
<i>Quercus coccinea</i>	scarlet oak	Oak wilt (<i>Bretziella fagacearum</i>)*	D	8	[1,34,175]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Forest tent caterpillar (<i>Malacosoma disstria</i>)	I/F	1	[1]
		Hypoxylon canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
		Walkingstick (<i>Diapheromera femorata</i>)	I/F	1	[1]
<i>Quercus douglasii</i>	blue oak	Oak wilt (<i>Bretziella fagacearum</i>)*	D	5	[34]
		Oak pit scale (<i>Asterolecanium</i> spp.)*	I/S	3	[177,178]
		Armillaria root disease (<i>Armillaria</i> spp.)	D	1	[1,36]
		Canker rot of oak (<i>Inonotus andersonii</i>)	D	1	[36]
		Hypoxylon canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36,178]
<i>Quercus ellipsoidalis</i>	northern pin oak	Oak wilt (<i>Bretziella fagacearum</i>)*	D	8	[34,175]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Twolined chestnut borer (<i>Agrilus bilineatus</i>)	I/B	3	[37]
		Armillaria root disease (<i>Armillaria</i> spp.)	D	1	[36]
		Hypoxylon canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Quercus emoryi</i>	Emory oak	Oak wilt (<i>Bretziella fagacearum</i>)*	D	8	[34,165]
		Canker rot of oak (<i>Inonotus andersonii</i>)	D	1	[36]
		Goldspotted oak borer (<i>Agrilus auroguttatus</i>)	I/B	1	[179]
		Hypoxylon canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
<i>Quercus engelmannii</i>	Engelmann oak	Oak wilt (<i>Bretziella fagacearum</i>)*	D	5	[34]
		Oak pit scale (<i>Asterolecanium</i> spp.)*	I/S	3	[178]
		Armillaria root disease (<i>Armillaria</i> spp.)	D	1	[36]
		Goldspotted oak borer (<i>Agrilus auroguttatus</i>)	I/B	1	[176]
		Hypoxylon canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
<i>Quercus falcata</i>	southern red oak	Oak wilt (<i>Bretziella fagacearum</i>)*	D	8	[1,34,175]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Sudden oak death (<i>Phytophthora ramorum</i>)*	D	3	[180]
		Twolined chestnut borer (<i>Agrilus bilineatus</i>)	I/B	3	[7,37]
		Hypoxylon canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
<i>Quercus gambelii</i>	Gambel oak	Oak wilt (<i>Bretziella fagacearum</i>)*	D	5	[34]
		Gambel oak borer (<i>Agrilus quercicola</i>)	I/B	3	[181]
		Canker rot of oak (<i>Inonotus andersonii</i>)	D	1	[7]
		Hypoxylon canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
<i>Quercus garryana</i>	Oregon white oak	Oak wilt (<i>Bretziella fagacearum</i>)*	D	5	[34]
		Oak pit scale (<i>Asterolecanium</i> spp.)*	I/S	3	[178]
		Armillaria root disease (<i>Armillaria</i> spp.)	D	1	[1,36]
		Eastern mistletoe (<i>Phoradendron leucarpum</i>)	P	1	[36]
		Hypoxylon canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
<i>Quercus georgiana</i>	Georgia oak	Oak wilt (<i>Bretziella fagacearum</i>)*	D	8	[13,34,175]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Quercus graciliformis</i>	slender oak	Hypoxyton canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
		Red oak borer (<i>Enaphalodes rufulus</i>)	I/B	1	[182]
		Oak wilt (<i>Bretziella fagacearum</i>)*	D	8	[13,34,175]
<i>Quercus gravesii</i>	Graves oak	Hypoxyton canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
		Oak wilt (<i>Bretziella fagacearum</i>)*	D	8	[13,34,175]
		Hypoxyton canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
<i>Quercus grisea</i>	gray oak	Red oak borer (<i>Enaphalodes rufulus</i>)	I/B	1	[182]
		Oak wilt (<i>Bretziella fagacearum</i>)*	D	5	[13,34,175]
		Hypoxyton canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
<i>Quercus hypoleucoides</i>	silverleaf oak	Oak wilt (<i>Bretziella fagacearum</i>)*	D	8	[34]
		Canker rot of oak (<i>Inonotus andersonii</i>)	D	1	[36]
		Goldspotted oak borer (<i>Agrilus auroguttatus</i>)	I/B	1	[179]
<i>Quercus ilicifolia</i>	scrub oak	Hypoxyton canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
		Oak wilt (<i>Bretziella fagacearum</i>)*	D	8	[34]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
<i>Quercus imbricaria</i>	shingle oak	Hypoxyton canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
		Oak wilt (<i>Bretziella fagacearum</i>)*	D	8	[34]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
<i>Quercus incana</i>	bluejack oak	Bacterial leaf scorch (<i>Xylella fastidiosa</i>)	D	1	[36]
		Endothia canker (<i>Endothia gyrosa</i>)	D	1	[36]
		Hypoxyton canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
<i>Quercus incana</i>	bluejack oak	Oak wilt (<i>Bretziella fagacearum</i>)*	D	8	[34]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Twolined chestnut borer (<i>Agrilus bilineatus</i>)	I/B	3	[37]
		Bacterial leaf scorch (<i>Xylella fastidiosa</i>)	D	1	[36]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Quercus kelloggii</i>	California black oak	Hypoxyton canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
		Oak wilt (<i>Bretziella fagacearum</i>)*	D	8	[34]
		Sudden oak death (<i>Phytophthora ramorum</i>)*	D	8	[36]
		Goldspotted oak borer (<i>Agrilus auroguttatus</i>)	I/B	5	[176]
		Oak pit scale (<i>Asterolecanium</i> spp.)*	I/S	3	[1,178]
<i>Quercus laceyi</i>	Lacey oak	Hypoxyton canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
		Oak wilt (<i>Bretziella fagacearum</i>)*	D	5	[13,34,175]
		Twolined chestnut borer (<i>Agrilus bilineatus</i>)	I/B	3	[37]
<i>Quercus laevis</i>	turkey oak	Hypoxyton canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
		Oak wilt (<i>Bretziella fagacearum</i>)*	D	8	[1,34]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Bacterial leaf scorch (<i>Xylella fastidiosa</i>)	D	1	[36]
		Canker rot of oak (<i>Inonotus andersonii</i>)	D	1	[36]
<i>Quercus laurifolia</i>	laurel oak	Hypoxyton canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
		Oak wilt (<i>Bretziella fagacearum</i>)*	D	8	[34]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Twolined chestnut borer (<i>Agrilus bilineatus</i>)	I/B	3	[37]
		Bacterial leaf scorch (<i>Xylella fastidiosa</i>)	D	1	[36]
<i>Quercus lobata</i>	California white oak	Hypoxyton canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
		Oak wilt (<i>Bretziella fagacearum</i>)*	D	5	[34]
		Oak pit scale (<i>Asterolecanium</i> spp.)*	I/S	3	[177,178]
		Canker rot of oak (<i>Inonotus andersonii</i>)	D	1	[36]
		Hypoxyton canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36,178]
<i>Quercus lyrata</i>	overcup oak	Oak wilt (<i>Bretziella fagacearum</i>)*	D	8	[34]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Quercus macrocarpa</i>	bur oak	Twolined chestnut borer (<i>Agrilus bilineatus</i>)	I/B	3	[37]
		Endothia canker (<i>Endothia gyrosa</i>)	D	1	[36]
		Hypoxylon canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
		Oak wilt (<i>Bretziella fagacearum</i>)*	D	5	[1,34,175]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Twolined chestnut borer (<i>Agrilus bilineatus</i>)	I/B	3	[37]
		Bacterial leaf scorch (<i>Xylella fastidiosa</i>)	D	1	[36]
<i>Quercus margarettae</i>	dwarf post oak	Bur oak blight (<i>Tubakia dryina</i>)	D	1	[183]
		Oak wilt (<i>Bretziella fagacearum</i>)*	D	5	[34]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Twolined chestnut borer (<i>Agrilus bilineatus</i>)	I/B	3	[37]
		Hypoxylon canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
<i>Quercus marilandica</i>	blackjack oak	Oak wilt (<i>Bretziella fagacearum</i>)*	D	8	[34,175]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Twolined chestnut borer (<i>Agrilus bilineatus</i>)	I/B	3	[37]
		Endothia canker (<i>Endothia gyrosa</i>)	D	1	[36]
		Hypoxylon canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
		Oak wilt (<i>Bretziella fagacearum</i>)*	D	5	[34]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
<i>Quercus michauxii</i>	swamp chestnut oak	Twolined chestnut borer (<i>Agrilus bilineatus</i>)	I/B	3	[37]
		Endothia canker (<i>Endothia gyrosa</i>)	D	1	[36]
		Hypoxylon canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
		Oak wilt (<i>Bretziella fagacearum</i>)*	D	5	[34]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Twolined chestnut borer (<i>Agrilus bilineatus</i>)	I/B	3	[37]
		Hypoxylon canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
<i>Quercus minima</i>	dwarf live oak	Red oak borer (<i>Enaphalodes rufulus</i>)	I/B	1	[1]
		Oak wilt (<i>Bretziella fagacearum</i>)*	D	5	[34]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Twolined chestnut borer (<i>Agrilus bilineatus</i>)	I/B	3	[37]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Quercus muehlenbergii</i>	chinkapin oak	Hypoxylon canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
		Oak wilt (<i>Bretziella fagacearum</i>)*	D	8	[1,34]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Twolined chestnut borer (<i>Agrilus bilineatus</i>)	I/B	3	[37]
<i>Quercus myrtifolia</i>	myrtle oak	Hypoxylon canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
		Oak wilt (<i>Bretziella fagacearum</i>)*	D	5	[13,34,175]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Hypoxylon canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
<i>Quercus nigra</i>	water oak	Red oak borer (<i>Enaphalodes rufulus</i>)	I/B	1	[182]
		Oak wilt (<i>Bretziella fagacearum</i>)*	D	8	[34]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Twolined chestnut borer (<i>Agrilus bilineatus</i>)	I/B	3	[37]
<i>Quercus oblongifolia</i>	Mexican blue oak	Hispidus canker (<i>Inonotus hispidus</i>)	D	1	[36]
		Hypoxylon canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
		Oak wilt (<i>Bretziella fagacearum</i>)*	D	8	[34]
		Canker rot of oak (<i>Inonotus andersonii</i>)	D	1	[36]
<i>Quercus oglethorpeensis</i>	Oglethorpe oak	Eastern mistletoe (<i>Phoradendron leucarpum</i>)	P	1	[36]
		Hypoxylon canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
		Oak wilt (<i>Bretziella fagacearum</i>)*	D	5	[34]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
<i>Quercus pagoda</i>	cherrybark oak	Hypoxylon canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
		Oak wilt (<i>Bretziella fagacearum</i>)*	D	8	[1,34]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Twolined chestnut borer (<i>Agrilus bilineatus</i>)	I/B	3	[37]
		Hispidus canker (<i>Inonotus hispidus</i>)	D	1	[36]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Quercus palustris</i>	pin oak	Hypoxyton canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
		Oak wilt (<i>Bretziella fagacearum</i>)*	D	8	[1,34]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Endothia canker (<i>Endothia gyrosa</i>)	D	1	[36]
<i>Quercus phellos</i>	willow oak	Hypoxyton canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
		Red oak borer (<i>Enaphalodes rufulus</i>)	I/B	1	[182]
		Oak wilt (<i>Bretziella fagacearum</i>)*	D	5	[34]
		Fall cankerworm (<i>Alsophila pometaria</i>)	I/F	3	[184]
<i>Quercus polymorpha</i>	Mexican white oak	Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Endothia canker (<i>Endothia gyrosa</i>)	D	1	[36]
		Hispidus canker (<i>Inonotus hispidus</i>)	D	1	[36]
		Oak wilt (<i>Bretziella fagacearum</i>)*	D	5	[13,34,175]
<i>Quercus prinoides</i>	dwarf chinquapin oak	Hypoxyton canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
		Oak wilt (<i>Bretziella fagacearum</i>)*	D	5	[34]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Hypoxyton canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
<i>Quercus prinus</i>	chestnut oak	Oak wilt (<i>Bretziella fagacearum</i>)*	D	5	[1,34]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Twolined chestnut borer (<i>Agrilus bilineatus</i>)	I/B	3	[37]
		Endothia canker (<i>Endothia gyrosa</i>)	D	1	[36]
<i>Quercus pungens</i>	sandpaper oak	Hypoxyton canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
		Oak wilt (<i>Bretziella fagacearum</i>)*	D	5	[13,34,175]
		Hypoxyton canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
		Oak wilt (<i>Bretziella fagacearum</i>)*	D	8	[1,34,175]
<i>Quercus rubra</i>	northern red oak	Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
		Red oak borer (<i>Enaphalodes rufulus</i>)	I/B	3	[182]
		Twolined chestnut borer (<i>Agrilus bilineatus</i>)	I/B	3	[37]
		Endothia canker (<i>Endothia gyrosa</i>)	D	1	[36]
<i>Quercus rugosa</i>	netleaf oak	Oak wilt (<i>Bretziella fagacearum</i>)*	D	5	[13,34,175]
		Hypoxylon canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
<i>Quercus shumardii</i>	Shumard oak	Oak wilt (<i>Bretziella fagacearum</i>)*	D	8	[1,34,175]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Hypoxylon canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
		Red oak borer (<i>Enaphalodes rufulus</i>)	I/B	1	[1]
		Twolined chestnut borer (<i>Agrilus bilineatus</i>)	I/B	1	[1]
<i>Quercus similis</i>	Delta post oak	Oak wilt (<i>Bretziella fagacearum</i>)*	D	5	[34]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Hypoxylon canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
		Red oak borer (<i>Enaphalodes rufulus</i>)	I/B	1	[182]
		Twolined chestnut borer (<i>Agrilus bilineatus</i>)	I/B	1	[185]
<i>Quercus sinuata</i>	Durand oak	Oak wilt (<i>Bretziella fagacearum</i>)*	D	5	[34]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Twolined chestnut borer (<i>Agrilus bilineatus</i>)	I/B	3	[37]
		Hypoxylon canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
<i>Quercus stellata</i>	post oak	Oak wilt (<i>Bretziella fagacearum</i>)*	D	5	[1,34]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Twolined chestnut borer (<i>Agrilus bilineatus</i>)	I/B	3	[37]
		Armillaria root disease (<i>Armillaria</i> spp.)	D	1	[36]
		Red oak borer (<i>Enaphalodes rufulus</i>)	I/B	1	[182]
<i>Quercus texana</i>	Nuttall oak	Oak wilt (<i>Bretziella fagacearum</i>)*	D	8	[1]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Quercus tomentella</i>	island live oak	Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Twolined chestnut borer (<i>Agrilus bilineatus</i>)	I/B	3	[37]
		Butt rot (<i>Hericium erinaceus</i>)	D	1	[36]
		Hypoxylon canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
		Oak wilt (<i>Bretziella fagacearum</i>)*	D	8	[13,34,175]
		Oak pit scale (<i>Asterolecanium</i> spp.)*	I/S	3	[178]
<i>Quercus velutina</i>	black oak	Hypoxylon canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
		Oak wilt (<i>Bretziella fagacearum</i>)*	D	8	[1,34,175]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Twolined chestnut borer (<i>Agrilus bilineatus</i>)	I/B	3	[37]
		Hypoxylon canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
		Red oak borer (<i>Enaphalodes rufulus</i>)	I/B	1	[1]
<i>Quercus virginiana</i>	live oak	Oak wilt (<i>Bretziella fagacearum</i>)*	D	8	[1,34,175]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Twolined chestnut borer (<i>Agrilus bilineatus</i>)	I/B	3	[37]
		Bacterial leaf scorch (<i>Xylella fastidiosa</i>)	D	1	[186]
		Bot canker (<i>Botryosphaeria corticola</i>)	D	1	[186]
		Oak wilt (<i>Bretziella fagacearum</i>)*	D	8	[34]
<i>Quercus wislizeni</i>	interior live oak	Goldspotted oak borer (<i>Agrilus auroguttatus</i>)	I/B	3	[179]
		Oak pit scale (<i>Asterolecanium</i> spp.)*	I/S	3	[178]
		Hypoxylon canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	D	1	[36]
		Wood borer (<i>Sphaeroma terebrans</i>)*	I/B	5	[187]
		Gall disease (<i>Cylindrocarpon didymum</i>)	D	1	[188,189]
		Wood borer (<i>Coccotrypes rhizophorae</i>)*	I/B	1	[188,190]
<i>Rhizophora mangle</i>	American mangrove	Fusarium wilt (<i>Fusarium oxysporum</i> f. sp. <i>rhois</i>)	D	3	[36]
<i>Rhus typhina</i>	staghorn sumac				

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Robinia neomexicana</i>	New Mexico locust	Heart rot fungi (<i>Phellinus robiniae</i>)	D	1	[36]
		Locust borer (<i>Megacyllene robiniae</i>)	I/B	1	[37]
		Locust leafminer (<i>Odontota dorsalis</i>)	I/F	1	[1]
<i>Robinia pseudoacacia</i>	black locust	Heart rot fungi (<i>Phellinus robiniae</i>)	D	1	[36]
		Locust borer (<i>Megacyllene robiniae</i>)	I/B	1	[1]
		Locust leafminer (<i>Odontota dorsalis</i>)	I/F	1	[1]
		Locust twig borer (<i>Ecdytolopha insiticiana</i>)	I/B	1	[1]
		Stunt nematode (<i>Tylenchorhynchus</i> spp.)	D	1	[36]
<i>Roystonea regia</i>	Florida royal palm	Thielaviopsis trunk rot (<i>Thielaviopsis paradoxa</i>)	D	3	[191]
		Ganoderma butt rot (<i>Ganoderma zonatum</i>)	D	1	[191]
		Royal palm bug (<i>Xylastodoris luteolus</i>)	I/S	1	[191]
<i>Sabal mexicana</i>	Mexican palmetto	Texas Phoenix palm decline (<i>Candidatus Phytoplasma palmae</i>)	D	5	[192]
		Thielaviopsis trunk rot (<i>Thielaviopsis paradoxa</i>)	D	5	[192]
		White mottled rot (<i>Ganoderma</i> spp.)	D	1	[192]
<i>Sabal palmetto</i>	cabbage palmetto	Texas Phoenix palm decline (<i>Candidatus Phytoplasma palmae</i>)	D	5	[192]
		Thielaviopsis trunk rot (<i>Thielaviopsis paradoxa</i>)	D	5	[192]
		White mottled rot (<i>Ganoderma</i> spp.)	D	1	[192]
<i>Salix alaxensis</i>	feltleaf willow	Scab and black canker of willow (<i>Venturia saliciperda/Glomerella miyabeana</i>)*	D	3	[34]
<i>Salix amygdaloides</i>	peachleaf willow	Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	1	[34]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Scab and black canker of willow (<i>Venturia saliciperda/Glomerella miyabeana</i>)*	D	3	[36]
		Artist's conk (<i>Ganoderma applanatum</i>)	D	1	[36]
		Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	1	[34]
<i>Salix arbusculoides</i>	peachleaf willow	Scab and black canker of willow (<i>Venturia saliciperda/Glomerella miyabeana</i>)*	D	3	[34]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Salix bebbiana</i>	Bebb willow	Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	1	[34]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Scab and black canker of willow (<i>Venturia saliciperda/Glomerella miyabeana</i>)*	D	3	[36]
		Artist's conk (<i>Ganoderma applanatum</i>)	D	1	[36]
		Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	1	[34]
		Willow witches'-broom (<i>Phytoplasma</i> , members of the yellows group)	D	1	[36]
<i>Salix bonplandiana</i>	Bonpland willow	Scab and black canker of willow (<i>Venturia saliciperda/Glomerella miyabeana</i>)*	D	3	[34]
		Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	1	[34]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
<i>Salix caroliniana</i>	coastal plain willow	Scab and black canker of willow (<i>Venturia saliciperda/Glomerella miyabeana</i>)*	D	3	[34]
		Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	1	[34]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
<i>Salix laevigata</i>	red willow	Scab and black canker of willow (<i>Venturia saliciperda/Glomerella miyabeana</i>)*	D	3	[34]
		Polyphagous shot hole borer (<i>Euwallacea spp.</i>)*	I/B	5	[193]
		Artist's conk (<i>Ganoderma applanatum</i>)	D	1	[36]
		Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	1	[34]
		Polyphagous shot hole borer (<i>Euwallacea spp.</i>)*	I/B	5	[162]
		Scab and black canker of willow (<i>Venturia saliciperda/Glomerella miyabeana</i>)*	D	3	[34]
<i>Salix lasiolepis</i>	arroyo willow	Artist's conk (<i>Ganoderma applanatum</i>)	D	1	[36]
		Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	1	[34]
		Polyphagous shot hole borer (<i>Euwallacea spp.</i>)*	I/B	5	[162]
		Scab and black canker of willow (<i>Venturia saliciperda/Glomerella miyabeana</i>)*	D	3	[34]
		Artist's conk (<i>Ganoderma applanatum</i>)	D	1	[36]
		Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	1	[34]
<i>Salix lucida</i> ssp. <i>lasiandra</i>	Pacific willow	Scab and black canker of willow (<i>Venturia saliciperda/Glomerella miyabeana</i>)*	D	3	[34]
		Artist's conk (<i>Ganoderma applanatum</i>)	D	1	[36]
		Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	1	[34]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Salix nigra</i>	black willow	Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Polyphagous shot hole borer (<i>Euvallacea</i> spp.)*	I/B	3	[162]
		Scab and black canker of willow (<i>Venturia saliciperda/Glomerella miyabeana</i>)*	D	3	[36]
		Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	1	[34]
		Willow witches'-broom (<i>Phytoplasma</i> , members of the yellows group)	D	1	[36]
<i>Salix pyrifolia</i>	balsam willow	Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Scab and black canker of willow (<i>Venturia saliciperda/Glomerella miyabeana</i>)*	D	3	[34]
		Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	1	[34]
<i>Salix scouleriana</i>	Scouler's willow	Scab and black canker of willow (<i>Venturia saliciperda/Glomerella miyabeana</i>)*	D	3	[34]
		Artist's conk (<i>Ganoderma applanatum</i>)	D	1	[36]
		Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	1	[34]
<i>Salix taxifolia</i>	yewleaf willow	Scab and black canker of willow (<i>Venturia saliciperda/Glomerella miyabeana</i>)*	D	3	[34]
		Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	1	[34]
<i>Sapindus drummondii</i>	western soapberry	Soapberry borer (<i>Agrilus prionurus</i>)*	I/B	8	[34,194]
<i>Sassafras albidum</i>	sassafras	Laurel wilt (<i>Raffaelea lauricola</i>)/redbay ambrosia beetle (<i>Xyleborus glabratus</i>)*	C	8	[34,37,195]
		Ganoderma rot of hardwoods (<i>Ganoderma lucidum</i>)	D	1	[36]
		Japanese beetle (<i>Popillia japonica</i>)*	I/F	1	[1]
		Verticillium wilt (<i>Verticillium albo-atrum</i>)	D	1	[35]
		Witches'-broom (<i>Apiosporina collinsii</i>)	D	1	[36]
<i>Sequoia sempervirens</i>	coast redwood	Brown cubical rot (<i>Poria sequoiae</i>)	D	1	[1]
		Cercospora blight (<i>Asperisporium sequoiae</i>)	D	1	[13,36,124]
		Heterobasidion root disease (<i>Heterobasidion</i> spp.)	D	1	[36]
		Sudden oak death (<i>Phytophthora ramorum</i>)*	D	1	[36]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Sequoiadendron giganteum</i>	giant sequoia	White ring rot (<i>Obba rivulosa</i>) Cercospora blight (<i>Asperisporium sequoiae</i>) Heterobasidion root disease (<i>Heterobasidion</i> spp.)	D D D	1 1 1	[1] [13,36,124] [36]
<i>Sideroxylon foetidissimum</i>	false mastic	False-mastic psylla (<i>Ceropsylla sideroxyli</i>) Lobate lac scale (<i>Paratachardina pseudolobata</i>)*	I/S I/S	1 1	[196] [97,197]
<i>Sideroxylon lanuginosum</i>	gum bumelia	none	.	.	.
<i>Sideroxylon lycioides</i>	buckthorn bumelia	none	.	.	.
<i>Sideroxylon salicifolium</i>	white bully	Lobate lac scale (<i>Paratachardina pseudolobata</i>)*	I/S	1	[97,197]
<i>Simarouba glauca</i>	paradise-tree	none	.	.	.
<i>Sophora affinis</i>	Texas sophra	none	.	.	.
<i>Sorbus americana</i>	American mountain-ash	Fire blight (<i>Erwinia amylovora</i>) Gypsy moth (<i>Lymantria dispar dispar</i>)*	D I/F	3 3	[36] [34]
<i>Sorbus decora</i>	northern mountain-ash	Fire blight (<i>Erwinia amylovora</i>) Gypsy moth (<i>Lymantria dispar dispar</i>)*	D I/F	3 3	[36] [34]
<i>Stewartia ovata</i>	mountain camellia	none	.	.	.
<i>Swietenia mahagoni</i>	West Indian mahogany	Mahogany shoot borer (<i>Hypsipyla grandella</i>) Nectria canker (<i>Nectria ditissima</i>)	I/B D	1 1	[37] [36]
<i>Taxodium ascendens</i>	pondcypress	Bagworm (<i>Thyridopteryx ephemeraeformis</i>) Cypress looper (<i>Anacamptodes pergracilus</i>) Southern cypress bark beetle (<i>Phloeosinus taxodii</i>)	I/F I/F I/B	1 1 1	[37] [1,198] [1]
<i>Taxodium distichum</i>	baldcypress	Baldcypress leafroller (<i>Archips goyerana</i>) Cercospora blight (<i>Asperisporium sequoiae</i>) Cypress looper (<i>Anacamptodes pergracilus</i>) Southern cypress bark beetle (<i>Phloeosinus taxodii</i>)	I/F D I/F I/B	1 1 1 1	[1] [36,124] [1,198] [1]
<i>Taxodium mucronatum</i>	Montezuma baldcypress	Southern cypress bark beetle (<i>Phloeosinus taxodii</i>)	I/B	1	[1]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Taxus brevifolia</i>	Pacific yew	White mottled rot (<i>Ganoderma</i> spp.)	D	1	[36]
		Black vine weevil (<i>Otiorhynchus sulcatus</i>)*	I/F	1	[34]
		Heterobasidion root disease (<i>Heterobasidion</i> spp.)	D	1	[36]
		Port-Orford-cedar root disease (<i>Phytophthora lateralis</i>)*	D	1	[35,90]
		Black vine weevil (<i>Otiorhynchus sulcatus</i>)*	I/F	1	[34]
<i>Taxus floridana</i>	Florida yew	Heterobasidion root disease (<i>Heterobasidion</i> spp.)	D	1	[36]
<i>Thrinax morrisii</i>	key thatch palm	Boisduval Scale (<i>Diaspis boisduvalii</i>)	I/S	1	[199]
<i>Thrinax radiata</i>	Florida thatch palm	Boisduval Scale (<i>Diaspis boisduvalii</i>)	I/S	1	[199]38
<i>Thuja occidentalis</i>	northern white-cedar	Heterobasidion root disease (<i>Heterobasidion</i> spp.)	D	1	[36]
<i>Thuja plicata</i>	western redcedar	Seiridium canker (<i>Seiridium cardinale</i>)	D	1	[36]
		Heterobasidion root disease (<i>Heterobasidion</i> spp.)	D	3	[1]
		Cercospora blight (<i>Asperisporium sequoiae</i>)	D	1	[36,124]
		Didymascella leaf blight (<i>Didymascella thujina</i>)	D	1	[36]
		Heterobasidion root disease (<i>Heterobasidion</i> spp.)	D	1	[36]
<i>Tilia americana</i>	American basswood	Seiridium canker (<i>Seiridium cardinale</i>)	D	1	[36]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Japanese beetle (<i>Popillia japonica</i>)*	I/F	1	[34]
<i>Tilia americana</i> var. <i>caroliniana</i>	Carolina basswood	Nectria canker (<i>Nectria ditissima</i>)	D	1	[36]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Japanese beetle (<i>Popillia japonica</i>)*	I/F	1	[34]
<i>Tilia americana</i> var. <i>heterophylla</i>	white basswood	Nectria canker (<i>Nectria ditissima</i>)	D	1	[36]
		Gypsy moth (<i>Lymantria dispar dispar</i>)*	I/F	3	[34]
		Japanese beetle (<i>Popillia japonica</i>)*	I/F	1	[34]
<i>Torreya californica</i>	California torreya (nutmeg)	Nectria canker (<i>Nectria ditissima</i>)	D	1	[36]
		none	.	.	.

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Torreya taxifolia</i>	Florida torreya	Canker disease (<i>Fusarium</i> spp.)	D	10	[200]
<i>Tsuga canadensis</i>	eastern hemlock	Hemlock woolly adelgid (<i>Adelges tsugae</i>)*	I/S	10	[37]
		Elongate hemlock scale (<i>Fiorinia externa</i>)*	I/S	3	[34]
		Hemlock borer (<i>Melanophila fulvoguttata</i>)	I/B	1	[1]
		Hemlock looper (<i>Lambdina fiscellaria</i>)	I/F	1	[1,37]
		Spruce budworm (<i>Choristoneura fumiferana</i>)	I/F	1	[1]
<i>Tsuga caroliniana</i>	Carolina hemlock	Hemlock woolly adelgid (<i>Adelges tsugae</i>)*	I/S	10	[37]
		Elongate hemlock scale (<i>Fiorinia externa</i>)*	I/S	3	[34]
		Hemlock borer (<i>Melanophila fulvoguttata</i>)	I/B	1	[201]
		Hemlock looper (<i>Lambdina fiscellaria</i>)	I/F	1	[202]
<i>Tsuga heterophylla</i>	western hemlock	Laminated root rot (<i>Phellinus sulphurascens</i>)	D	5	[20,203]
		Hemlock dwarf mistletoe (<i>Arceuthobium tsugense</i>)	P	5	[36]
		Heterobasidion root disease (<i>Heterobasidion</i> spp.)	D	3	[36]
		Schweinitzii root and butt rot (<i>Phaeolus schweinitzii</i>)	D	3	[1]
		Western hemlock looper (<i>Lambdina fiscellaria lugubrosa</i>)	I/F	3	[1]
<i>Tsuga mertensiana</i>	mountain hemlock	Laminated root rot (<i>Phellinus sulphurascens</i>)	D	5	[1,20]
		Hemlock dwarf mistletoe (<i>Arceuthobium tsugense</i>)	P	5	[1]
		Heterobasidion root disease (<i>Heterobasidion</i> spp.)	D	3	[1]
		Schweinitzii root and butt rot (<i>Phaeolus schweinitzii</i>)	D	3	[1]
<i>Ulmus alata</i>	winged elm	Dutch elm disease (<i>Ophiostoma novo-ulmi</i>)*	D	8	[36]
		Elm yellows (<i>Candidatus Phytoplasma ulmi</i>)	D	5	[36,204]
		Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	3	[34]
<i>Ulmus americana</i>	American elm	Dutch elm disease (<i>Ophiostoma novo-ulmi</i>)*	D	8	[36]
		Elm yellows (<i>Candidatus Phytoplasma ulmi</i>)	D	5	[36,204]
		Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	3	[34]

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

Host species scientific name	Host species common name	Agent	T/G	Sev.	Sources
<i>Ulmus crassifolia</i>	cedar elm	Banded elm bark beetle (<i>Scolytus schevyrewi</i>)*	I/B	3	[34,35]
		Pit canker (<i>Phytophthora inflata</i>)	D	1	[36]
		Dutch elm disease (<i>Ophiostoma novo-ulmi</i>)*	D	8	[36]
		Elm yellows (<i>Candidatus Phytoplasma ulmi</i>)	D	5	[36,204]
<i>Ulmus rubra</i>	slippery elm	Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	3	[34]
		Dutch elm disease (<i>Ophiostoma novo-ulmi</i>)*	D	5	[36]
		Elm yellows (<i>Candidatus Phytoplasma ulmi</i>)	D	5	[36,204]
		Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	3	[34]
<i>Ulmus serotina</i>	September elm	Dutch elm disease (<i>Ophiostoma novo-ulmi</i>)*	D	8	[36]
		Elm yellows (<i>Candidatus Phytoplasma ulmi</i>)	D	5	[36]
		Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	3	[34]
		Dutch elm disease (<i>Ophiostoma novo-ulmi</i>)*	D	8	[36]
<i>Ulmus thomasii</i>	rock elm	Elm yellows (<i>Candidatus Phytoplasma ulmi</i>)	D	5	[13]
		Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	I/B	3	[34]
		Banded elm bark beetle (<i>Scolytus schevyrewi</i>)*	I/B	3	[36]
		Laurel wilt (<i>Raffaelea lauricola</i>)/redbay ambrosia beetle (<i>Xyleborus glabratus</i>)*	C	8	[205]
<i>Umbellularia californica</i>	California-laurel	Bacterial leaf blight (<i>Pseudomonas lauraceum</i>)	D	1	[36]
		Sudden oak death (<i>Phytophthora ramorum</i>)*	D	1	[36]
		White mottled rot (<i>Ganoderma</i> spp.)	D	1	[36]
		Giant palm borer (<i>Dinapate wrightii</i>)	I/B	3	[7]
<i>Washingtonia filifera</i>	California fan palm	False smut of palms (<i>Graphioa phoenicis</i>)	D	1	[36]
		Wood rotting (<i>Phanerochaete tuberculata</i>)	D	1	[206]
<i>Yucca brevifolia</i>	Joshua tree	none	.	.	.
<i>Yucca treculeana</i>	Trecul yucca	none	.	.	.
<i>Zanthoxylum clava-herculis</i>	Hercules-club	none	.	.	.
<i>Zanthoxylum fagara</i>	lime prickly-ash	none	.	.	.

T/G=Taxon/Guild. Taxon: I=insect, D=disease, P=parasitic plant, C=insect and disease multi-agent complex; Guild (for insects): B=phloem- or wood-borers, F=foliage-feeders, S=sap-feeders. Sev.=Severity. Nonnative invasive agents are indicated with *.

References

1. *Silvics of North America*. Burns, R.M.; Honkala, B.H., Eds.; United States Department of Agriculture, Forest Service: Washington, D.C., 1990; p 877.
2. Wilken, D. Plant Guide: Pacific Silver Fir. Available online: https://plants.usda.gov/plantguide/pdf/pg_abam.pdf (accessed on February 26, 2019).
3. Ragenovich, I.R.; Mitchell, R.G. *Balsam Woolly Adelgid*; Forest Insect & Disease Leaflet 118; U.S. Department of Agriculture, Forest Service: Washington, D.C., 2006; p 11.
4. Thies, W.G.; Sturrock, R.N. *Laminated root rot in western North America*; Gen. Tech. Rep. PNW-GTR-349; United States Department of Agriculture, Forest Service, Pacific Northwest Research Station: Portland, Oregon, 1995; p 32.
5. Global Invasive Species Database. Adelges piceae. Available online: <http://www.iucngisd.org/gisd/species.php?sc=226> (accessed on March 1, 2019).
6. Hansen, E.M.; Lewis, K.J.; American Phytopathological Society. *Compendium of conifer diseases*; APS Press, the American Phytopathological Society: St. Paul, MN, 1997; pp. 101.
7. United States Department of Agriculture Forest Service. Fire Effects Information System (FEIS). Available online: <https://www.feis-crs.org/feis/> (accessed on February 26, 2019).
8. Popp, J.B.; Lundquist, J.E. *Pathology collection of the Rocky Mountain Research Station*; RMRS GTR-167WWW; United States Department of Agriculture, Forest Service, Rocky Mountain Research Station: Fort Collins, Colorado, 2006; p 183.
9. Scharpf, R.F.; Staley, J.; Hawksworth, F.G. A needle cast, the first known disease of Bristlecone Fir in California. *Plant Dis. Rep.* **1970**, *54*, 275-277.
10. Scharpf, R.F. *Diseases of Pacific Coast Conifers*; Agriculture Handbook 521; United States Department of Agriculture, Forest Service, Pacific Southwest Research Station: Albany, California, 1993; p 199.
11. Furniss, R.L.; Carolin, V.M.; United States. Forest Service. *Western Forest Insects*; Miscellaneous Publication 1339; United States Department of Agriculture, Forest Service: Washington D.C., 1977; p 654.
12. United States Department of Agriculture Forest Service. *Field Guide to Diseases and Insects of the Rocky Mountain Region*; Gen. Tech. Report RMRS-GTR-241; United States Department of Agriculture, Forest Service, Rocky Mountain Research Station: Fort Collins, Colorado, 2010; p 336.
13. Horst, R.K. *Westcott's Plant Disease Handbook*, Eighth edition ed.; SpringerReference: Dordrecht, Netherlands, 2013; pp. 826 pages.
14. United States Department of Agriculture Natural Resources Conservation Service. Plant Fact Sheet: White Fir. Available online: https://plants.usda.gov/factsheet/pdf/fs_abco.pdf (accessed on March 1, 2019).
15. Wickman, B.E.; Mason, R.R.; Trostle, G.C. *Douglas-Fir Tussock Moth*; Forest Insect & Disease Leaflet 86; United States Department of Agriculture, Forest Service: Washington, D.C., 1981; p 10.
16. Otrosina, W.J.; Scharpf, R.F. *Proceedings of the Symposium on Research and Management of Annosus Root Disease (*Heterobasidion annosum*) in Western North America*; Gen. Tech. Rep. PSW-GTR-116; United States Department of Agriculture, Forest Service, Pacific Southwest Forest and Range Experiment Station: Berkeley, California, 1989; p 177.
17. Talgo, V.; Chastagner, G. *Phytophthora on Abies*; JKI Data Sheets – Plant Diseases and Diagnosis 77; Julius Kühn-Institut, Federal Research Centre for Cultivated Plants: Quedlinburg, Germany, 2013; p 11.
18. Hagle, S.K. Management Guide for Armillaria Root Disease In *Forest Insect and Disease Management Guide for the Northern and Central Rocky Mountains*, United States Department of Agriculture, Forest Service, Northern and Intermountain Regions, State and Private Forestry: Missoula, Montana, and Boise, Idaho, 2008; p. 23.
19. Ferrell, G.T. *Fir Engraver*; Forest Insect & Disease Leaflet 13; United States Department of Agriculture, Forest Service: Washington, D.C., 1986; p 8.
20. Hagle, S.K. Management Guide for Laminated Root Rot. In *Forest Insect and Disease Management Guide for the Northern and Central Rocky Mountains*, United States Department of Agriculture, Forest Service, Northern and Intermountain Regions, State and Private Forestry: Missoula, Montana, and Boise, Idaho, 2009; p. 20.

21. Gonthier, P.; Nicolotti, G. *Infectious Forest Diseases*; CAB International: Wallingford, Oxfordshire, United Kingdom, 2013.
22. Nesom, G. Plant Guide: Subalpine Fir. Available online: https://plants.usda.gov/plantguide/pdf/pg_abla.pdf (accessed on February 27, 2019).
23. Spiegel, L.H.; Chadwick, K.; Mehmel, C. Assessment of balsam woolly adelgid damage to eastern Washington and Oregon subalpine fir (Project WC-F-07-01). In *Forest Health Monitoring: National Status, Trends, and Analysis 2010*, Potter, K.M., Conkling, B.L., Eds. United States Department of Agriculture Forest Service, Southern Research Station: Asheville, North Carolina, 2013; Vol. Gen. Tech. Rep. SRS-GTR-176, pp. 129-134.
24. Kegley, S. Management Guide for Western Balsam Bark Beetle. In *Forest Insect and Disease Management Guide for the Northern and Central Rocky Mountains*, United States Department of Agriculture, Forest Service, Northern and Intermountain Regions, State and Private Forestry: Missoula, Montana, and Boise, Idaho, 2006; p. 4.
25. Fellin, D.G., Dewey, J.E. *Western Spruce Budworm*; Forest Insect & Disease Leaflet 53; United States Department of Agriculture, Forest Service: Washington, D.C., 1982; p 10.
26. Fairweather, M.L.; Rogers, J.M.T.; Conklin, D.; Fitzgibbon, B. *Field Guide to Insects and Diseases of Arizona and New Mexico Forests*; MR-R3-16-3; United States Department of Agriculture Forest Service, Southwestern Region: Albuquerque, New Mexico, 2006; p 271.
27. Randall, C.B. Management Guide for Fir Engraver In *Forest Insect and Disease Management Guide for the Northern and Central Rocky Mountains*, United States Department of Agriculture, Forest Service, Northern and Intermountain Regions, State and Private Forestry: Missoula, Montana, and Boise, Idaho, 2006; p. 7.
28. Filip, G.M.; Goheen, D.J.; Kimmey, J.W. *Rust-red stringy rot caused by the Indian Paint Fungus*; Forest Insect & Disease Leaflet 93; United States Department of Agriculture, Forest Service, Pacific Northwest Region: Portland, Oregon, 2009; p 11.
29. Goheen, E.M.; Willhite, E.A. *Field Guide to Common Diseases and Insect Pests of Oregon and Washington Conifers*; R6-NR-FID-PR-01-06; United States Forest Service, Pacific Northwest Region: Portland, Oregon, 2006; p 327.
30. Filip, G.M.; Beatty, J.S.; Mathiasen, R.L. Fir dwarf mistletoe. In *Forest Insect & Disease Leaflet 89*, United States Department of Agriculture, Forest Service: Washington, D.C., 2000.
31. Filip, G.M.; Schmitt, C.L. *R_x for Abies: Silvicultural Options for Diseased Firs in Oregon and Washington*; Gen. Tech. Rep. PNW-252; United States Department of Agriculture, Forest Service, Pacific Northwest Research Station: Portland, Oregon, 1990.
32. Parrotta, J.A. *Acacia farnesiana* (L.) Willd., *Aroma, huisache* : Leguminosae (Mimosoideae), legume family; SO-ITF-SM-49; U.S. Forest Service, Southern Forest Experiment Station, Institute of Tropical Forestry: Rio Piedras, Puerto Rico, 1992; p 5.
33. Francis, J.K. *Wildland Shrubs of the United States and its Territories: Thamnic Descriptions: Volume 1*; Gen. Tech. Rep. IITF-GTR-26; United States Department of Agriculture, Forest Service, Rocky Mountain Research Station: Rio Piedras, Puerto Rico, 2004; p 830.
34. United States Department of Agriculture Forest Service. Alien Forest Pests and Their Hosts Available online: http://www.nrs.fs.fed.us/tools/afpe/data/tree_list/ (accessed on February 25, 2019).
35. Center for Invasive Species and Ecosystem Health. Forest Pests of North America. Available online: https://www.pests.org/host_trees.html (accessed on February 25, 2019).
36. Sinclair, W.A.; Lyon, H.H. *Diseases of trees and shrubs*, 2nd ed.; Comstock Pub. Associates/Cornell University Press: Ithaca, N.Y., 2005; pp. xiii, 660 p.
37. Ciesla, W.M. *Forest entomology : a global perspective*; Wiley-Blackwell: Chichester, West Sussex ; Hoboken, NJ, 2011; pp. viii, 400 p., [440] p. of col. plates.
38. Eskalen, A.; Stouthamer, R.; Lynch, S.C.; Rugman-Jones, P.F.; Twizeyimana, M.; Gonzalez, A.; Thibault, T. Host Range of Fusarium Dieback and Its Ambrosia Beetle (Coleoptera: Scolytinae) Vector in Southern California. *Plant Dis.* **2013**, 97, 938-951, doi: <https://doi.org/10.1094/PDIS-11-12-1026-RE>.

39. Garbelotto, M.; Davidson, J.M.; Ivors, K.; Maloney, P.E.; Hüberli, D.; Koike, S.T.; Rizzo, D.M. Non-oak native plants are main hosts for sudden oak death pathogen in California. *Calif. Agr.* **2003**, *57*, 18-23, doi: <https://doi.org/10.3733/ca.v057n01p18>.
40. Eatough Jones, M.; Paine, T.D. Potential pesticides for control of a recently introduced ambrosia beetle (*Euwallacea* sp.) in southern California. *J. Pest Sci.* **2018**, *91*, 237-246, doi: <https://doi.org/10.1007/s10340-017-0866-8>.
41. Kessler, K.J.; Hadfield, J.S. *Eutypella Canker of Maple*; Forest Pest Leaflet 136; United States Department of Agriculture, Forest Service: Washington, D.C., 1972; p 6.
42. Pijut, P.M. *Diseases in Hardwood Tree Plantings*; FNR-221; Hardwood Tree Improvement and Regeneration Center, Purdue University and United States Department of Agriculture Forest Service, Northern Research Station: West Lafayette, Indiana, 2006; p 16.
43. Dickerson, J. Plant Fact Sheet: Red Maple. Available online: https://plants.usda.gov/factsheet/pdf/fs_acru.pdf (accessed on March 1, 2019).
44. Gould, A.B.; Lashomb, J.H. Bacterial leaf scorch (BLS) of shade trees. *The Plant Instructor* **2007**, *10.1094/PHI-I-2007-0403-07*, doi: 10.1094/PHI-I-2007-0403-07.
45. Heimann, M.F.; Worf, G.L. *Maple and Other Trees Disorder: Verticillium Wilt*; A2537; University of Wisconsin-Extension, Cooperative Extension: Madison, Wisconsin, 1997; p 2.
46. United States Department of Agriculture Animal and Plant Health Inspection Service. *Japanese Beetle (*Popillia japonica Newman*)*; Program Aid No. 1881; United States Department of Agriculture, Animal and Plant Health Inspection Service: 2007; p 2.
47. Berlanger, I.; Powelson, M.L. Verticillium wilt. *The Plant Health Instructor* **2005**.
48. Broschat, T.K. *Acoelorrhaphes wrightii: Paurotis Palm*; ENH-217; University of Florida, Institute of Food and Agricultural Sciences Extension: Gainesville, Florida, 2014; p 3.
49. Inserra, R.N.; Dunn, R.A.; Vovlas, N. Host Response of Ornamental Palms to *Rotylenchulus reniformis*. *J. Nematol.* **1994**, *26*, 737-743.
50. Purcell, A.H.; Saunders, S.R. Fate of Pierce's Disease Strains of *Xylella fastidiosa* in Common Riparian Plants in California. *Plant Dis.* **1999**, *83*, 825-830, doi: <https://doi.org/10.1094/PDIS.1999.83.9.825>.
51. Rizzo, D.M.; Garbelotto, M.; Davidson, J.M.; Slaughter, G.M.; Koike, S.T. *Phytophthora ramorum* and sudden oak death in California: I. Host relationships. In *Proceedings of the Fifth Symposium on Oak Woodlands: Oaks in California's Challenging Landscape*, Standiford, R.B., Ed. United States Department of Agriculture, Forest Service, Pacific Southwest Research Station: Albany, California, 2002; Vol. Gen. Tech. Rep. PSW-GTR-184, pp. 733-740.
52. Santamour, F.S.J.; Riedel, L.G.H. Susceptibility of various landscape trees to root-knot nematodes. *Journal of Arboriculture* **1993**, *19*, 257-259.
53. DePamphilis, C.W.; Neufeld, H.S. Phenology and ecophysiology of *Aesculus sylvatica*, a vernal understory tree. *Botany* **1989**, *67*, 2161-2167, doi: <https://doi.org/10.1139/b89-274>.
54. Sims, L.; Goheen, E.; Kanaskie, A.; Hansen, E. Alder Canopy Dieback and Damage in Western Oregon Riparian Ecosystems. *Northwest Sci.* **2015**, *89*, 34-46, doi: <https://doi.org/10.3955/046.089.0103>.
55. Worrall, J.J.; Adams, G.C.; Tharp, S.C. Summer heat and an epidemic of cytospora canker of *Alnus*. *Can. J. Plant Pathol.* **2010**, *32*, 376-386, doi: <https://doi.org/10.1080/07060661.2010.499265>.
56. Kruse, J.J.; Zogas, K.; Hard, J.; Lisuzzo, N. *Pest Alert New Pest in Alaska and Washington: The Green Alder Sawfly - Monsoma pulveratum (Retzius)*; R10-PR-022; United States Department of Agriculture, Forest Service, State and Private Forestry: Anchorage, Alaska, 2010; p 2.
57. United States Department of Agriculture Forest Service. *Forest Health Conditions in Alaska 2010: A Forest Health Protection Report*; Forest Protection Report R10-PR-23; United States Department of Agriculture, Forest Service, Alaska Region: Anchorage, Alaska, 2010; p 78.
58. United States Department of Agriculture Forest Service. *Forest Health Conditions in Alaska 2017: A Forest Health Protection Report*; Forest Protection Report R10-PR-43; United States Department of Agriculture, Forest Service, Alaska Region: Anchorage, Alaska, 2018; p 64.
59. Batzer, H.O.; Morris, R.C. *Forest Tent Caterpillar*; Forest Insect & Disease Leaflet 9; United States Department of Agriculture, Forest Service: Washington, D.C., 1978; p 8.

60. Nesom, G. Plant Guide: Downy Serviceberry. Available online: https://plants.usda.gov/plantguide/pdf/pg_amar3.pdf (accessed on February 27, 2019).
61. Gilman, E.F.; Watson, D.G. *Amelanchier arborea: Downy Serviceberry*; ENH-232; University of Florida, Institute of Food and Agricultural Sciences Extension: Gainesville, Florida, 2006; p 3.
62. Moorman, G.W. Amelanchier Diseases. Available online: <https://extension.psu.edu/amelanchier-diseases> (accessed on March 1, 2019).
63. Niemiec, S.S.; Ahrens, G.R.; Willits, S.; Hibbs, D.E. *Hardwoods of the Pacific Northwest*; Research Contribution 8; Oregon State University, College of Forestry, Forest Research Laboratory: Corvallis, Oregon, 1995; p 115.
64. Bennett, M.; Shaw, D. *Diseases and Insect Pests of Pacific Madrone*; EC 1619-E; Oregon State University, Extension Service: Corvallis, Oregon, 2008; p 5.
65. Immel, D.L. Plant Guide: Pacific Madrone. Available online: https://plants.usda.gov/plantguide/pdf/cs_arne.pdf (accessed on March 1, 2019).
66. Elliott, M.; Edmonds, R.L. Systematic Study of *Nattrassia mangiferae*, the Cause of Madrone Canker In *Proceedings of the 51st Western International Forest Disease Work Conference*, Geils, B.W., Ed. United States Department of Agriculture, Forest Service, Rocky Mountain Research Station: Flagstaff, Arizona, 2003; pp. 59-62.
67. Layne, D.R. The Pawpaw [*Asimina triloba* (L.) Dunal]: A New Fruit Crop for Kentucky and the United States. *HortScience* **1996**, *31*, 777-784.
68. Gilbert, G.S.; Sousa, W.P. Host Specialization among Wood-Decay Polypore Fungi in a Caribbean Mangrove Forest. *Biotropica* **2002**, *34*, 396-404.
69. Muilenburg, V.L.; Herms, D.A. A Review of Bronze Birch Borer (Coleoptera: Buprestidae) Life History, Ecology, and Management. *Environ. Entomol.* **2012**, *41*, 1372-1385, doi: <https://doi.org/10.1603/en12238>.
70. Katovich, S.A.; Munson, A.S.B., J.; McCullough, D. *Bronze Birch Borer*; Forest Insect & Disease Leaflet 111; United States Department of Agriculture, Forest Service: Washington, D.C., 2000; p 8.
71. McCormick, M.A.; Cubeta, M.A.; Grand, L.F. Geography and hosts of the wood decay fungi *Fomes fasciatus* and *Fomes fomentarius* in the United States. *North American Fungi* **2013**, *8*, 1-53, doi: <http://dx.doi.org/10.2509/naf2013.008.002>.
72. Wennerberg, S. Plant Guide: Gray Birch. Available online: https://plants.usda.gov/plantguide/pdf/pg_bepo.pdf (accessed on February 27, 2019).
73. Santamour, F.S.J. Differential feeding by adult Japanese beetles on foliage of birch (*Betula*) species and hybrids. *Journal of Arboriculture* **2001**, *27*, 18-23.
74. Caldwell, D.L.; Palmateer, A.J. Gumbo-Limbo Trees Struggle With Three New Dieback Plagues: Puzzling Witches'-broom Symptoms, Croton Scale, and Rugose Spiraling Whitefly. *Proc. Fl. State Hortic.* **2014**, *127*, 158-159.
75. Smith, T.F.; Rizzo, D.M.; North, M. Patterns of mortality in an old-growth mixed-conifer forest of the southern Sierra Nevada, California. *For. Sci.* **2005**, *51*, 266-275.
76. Hoover, G.A. *Entomological Notes: Juniper Scale*; EV-7; The Pennsylvania State University, College of Agricultural Sciences, Cooperative Extension: University Park, Pennsylvania, 2002; p 1.
77. Pierson, E.A.; Turner, R.M. An 85-Year Study of Saguaro (*Carnegiea gigantea*) Demography. *Ecology* **1998**, *79*, 2676-2693, doi: <https://doi.org/10.2307/176509>.
78. Kelly, J.; Olsen, M.W. *Problems and Pests of Agave, Aloe, Cactus and Yucca*; AZ 1399; The University of Arizona, Cooperative Extension: Tucson, Arizona, 2011; p 9.
79. Davidson, C.B.; Gottschalk, K.W.; Johnson, J.E. Tree Mortality Following Defoliation by the European Gypsy Moth (*Lymantria dispar* L.) in the United States: A Review. *Forest Sci.* **1999**, *45*, 74-84.
80. Grand, L.; Vernia, C. New taxa and hosts of poroid wood-decay fungi in North Carolina. *Castanea* **2002**, *67*, 193-200.
81. Grand, L.F.; Vernia, C.S. Biogeography and hosts of poroid wood decay fungi in North Carolina: species of *Fomes*, *Fomitopsis*, *Fomitella* and *Ganoderma*. *Mycotaxon* **2005**, *94*, 231-234.
82. McCracken, F.I. *Canker-rots in Southern Hardwoods*; Forest Insect & Disease Leaflet 33; United States Department of Agriculture, Forest Service: Washington, D.C., 1978; p 4.
83. Forcella, F. Why twig-girdling beetles girdle twigs. *Naturwissenschaften* **1982**, *69*, 398-400, doi: 10.1007/bf00396699.

84. Anagnostakis, S.L. The Effect of Multiple Importations of Pests and Pathogens on a Native Tree. *Biol. Invasions* **2001**, *3*, 245–254, doi: 10.1023/a:1015205005751.
85. Case, A.E.; Mayfield, I.I.I.A.E.; Clark, S.L.; Schlarbaum, S.E.; Reynolds, B.C. Abundance and Frequency of the Asiatic Oak Weevil (Coleoptera: Curculionidae) and Defoliation on American, Chinese, and Hybrid Chestnut (*Castanea*). *J. Insect Sci.* **2016**, *16*, 29–29, doi: 10.1093/jisesa/iew012.
86. Payne, J.A.; Miller, G.; Johnson, G.P.; Senter, S.D. *Castanea pumila* (L.) Mill.: An Underused Native Nut Tree. *HortScience* **1994**, *29*, 62–131.
87. Geyer, W.A.; Broyles, P.J.; Row, J.M. Plant Guide: Southern Catalpa. Available online: https://plants.usda.gov/plantguide/pdf/pg_cabi8.pdf (accessed on February 27, 2019).
88. Geyer, W.A.; Broyles, P.J. Plant Guide: Northern Catalpa. Available online: https://plants.usda.gov/plantguide/pdf/pg_casp8.pdf (accessed on February 29, 2019).
89. Furniss, M.M.; Ferguson, D.C.; Voget, K.W.; Burkhardt, J.W.; Tiedemann, A.R.; Oldemeyer, J.L. *Taxonomy, life history, and ecology of a mountain-mahogany defoliator, Stamnodes animata (Pearsall)*, in Nevada; Fish and Wildlife Research 3; U.S. Dept. of the Interior, Fish and Wildlife Service: Washington, D.C., 1988; pp ix, 26 pages.
90. Bettjewski, F.; Goheen, D.J.; Angwin, P.A.; Snieszko, R.A. *Port-Orford-Cedar Root Disease*; Forest Insect & Disease Leaflet 131; United Department of Agriculture, Forest Service, Pacific Northwest Region: Portland, OR, 2011; p 12.
91. Thiemann, D.; Lopez, V.; Ray, A.M.; Cipollini, D. The History of Attack and Success of Emerald Ash Borer (Coleoptera: Buprestidae) on White Fringetree in Southwestern Ohio. *Environ. Entomol.* **2016**, *45*, 961–966, doi: <https://doi.org/10.1093/ee/nvw073>.
92. Cipollini, D. White Fringetree as a Novel Larval Host for Emerald Ash Borer. *J. Econ. Entomol.* **2015**, *108*, 370–375, doi: <https://doi.org/10.1093/jee/tou026>.
93. Saavedra, A.; Hansen, E.M.; Goheen, D.J. *Phytophthora cambivora* in Oregon and its pathogenicity to *Chrysophyilla chrysophylla*. *Forest Pathol.* **2007**, *37*, 409–419, doi: <https://doi.org/10.1111/j.1439-0329.2007.00515.x>.
94. Hamon, A.; Fasulo, T.R.; Buss, L.J. *Cardin's Whitefly, Metaleurodinus cardini (Back)* (Insecta: Hemiptera: Aleyrodidae: Aleurodicinae); EENY-15; University of Florida, Institute of Food and Agricultural Sciences Extension Service: Gainesville, Florida, 2014; p 4.
95. Kern, W.H. *Fiddlewood Leafroller, Seagrape Moth (suggested common names) Epicorsia oedipodalis* (Guenée, 1854) (Lepidoptera: Ditrysia: Pyraloidea: Pyralidae: Pyraustinae); ENH-617; University of Florida, Institute of Food and Agricultural Sciences Extension: Gainesville, Florida, 2015; p 4.
96. Gilman, E.F.; Watson, D.G. *Cladrastis kentukea: American Yellowwood*; ENH329; University of Florida, Institute of Food and Agricultural Sciences Extension: Gainesville, Florida, 2014; p 3.
97. Howard, F.W.; Pemberton, R.; Hamon, A.; Hodges, G.S.; Steinberg, B.; Mannion, C.M.; McLean, D.; Wofford, J. *Lobate Lac Scale, Paratachardina lobata lobata (Chamberlin)* (Hemiptera: Sternorrhyncha: Coccoidea: Kerriidae); EENY-276; University of Florida, Institute of Food and Agricultural Sciences Extension: Gainesville, Florida, 2017; p 14.
98. Schoch, C.L.; Crous, P.W.; Witthuhn, R.C.; Cronwright, G.; El-Gholl, N.E.; Wingfield, B.D. Recombination in *Calonectria morganii* and Phylogeny with Other Heterothallic Small-Spored Calonectria Species. *Mycologia* **2000**, *92*, 665–673, doi: <https://doi.org/10.2307/3761424>.
99. Neal, A. *Sri Lankan Weevil Myllocerus undecimpustulatus undatus Marshall*; EENY579; University of Florida, Institute of Food and Agricultural Sciences Extension: Gainesville, Florida, 2013; p 4.
100. Gilman, E.F.; Watson, D.G. *Cordia sebestena: Geiger Tree*; ENH-341; University of Florida, Institute of Food and Agricultural Sciences Extension: Gainesville, Florida, 1993; p 2.
101. Linsky, J.; Manco, B.N. *Cordia sebestena* var. *sebestena*. The IUCN Red List of Threatened Species. Version 2018-2. Available online: <http://dx.doi.org/10.2305/IUCN.UK.2014-1.RLTS.T56497523A56504011.en> (accessed on March 2, 2019).
102. Palmateer, A.J.; Pérez, J.M.; Cating, R.A.; Ploetz, R.C.; Hoy, M.A. First Report of Tar Spot on Orange Geiger, *Cordia sebestena*, Caused by *Diatractium cordianum* in Florida. *Plant Dis.* **2008**, *92*, 1250–1250, doi: <https://doi.org/10.1094/PDIS-92-8-1250B>.

103. Gyeltshen, J.; Hedges, A. *Dogwood Borer, Synanthonedon scitula (Harris) (Insecta: Lepidoptera: Sesiidae)*; EENY-374; University of Florida, Institute of Food and Agricultural Sciences Extension: Gainesville, Florida, February 2, 2016, 2006; p 3.
104. Gilman, E.F.; Watson, D.G. *Cotinus obovatus: American Smoketree*; ENH-367; University of Florida, Institute of Food and Agricultural Sciences Extension: Gainesville, Florida, 2006; p 3.
105. Carroll, J.; Rosa, L.D. Cherry Bark Tortrix Moth. Available online: <https://nysipm.cornell.edu/agriculture/fruits/invasive-species-exotic-pests/> (accessed on February 26, 2019).
106. Lorenzo, A.B. Plant Guide: Brainerd's Hawthorn. Available online: https://plants.usda.gov/plantguide/pdf/pg_crbr3.pdf (accessed on February 27, 2019).
107. Lorenzo, A.B. Plant Guide: Pear Hawthorn. Available online: https://plants.usda.gov/plantguide/pdf/pg_crca.pdf (accessed on February 27, 2019).
108. Tobin, P.C.; Bai, B.B.; Eggen, D.A.; Leonard, D.S. The ecology, geopolitics, and economics of managing *Lymantria dispar* in the United States. *Int. J. Pest Manage.* **2012**, *58*, 195-210, doi: <https://doi.org/10.1080/09670874.2011.647836>.
109. Lorenzo, A.B. Plant Guide: Cockspur Hawthorn. Available online: https://plants.usda.gov/plantguide/pdf/pg_crcr2.pdf (accessed on February 28, 2019).
110. Lorenzo, A.B. Plant Guide: Fireberry Hawthorn. Available online: https://plants.usda.gov/plantguide/pdf/pg_crch.pdf (accessed on February 27, 2019).
111. Lorenzo, A.B. Plant Guide: Fanleaf Hawthorn. Available online: https://plants.usda.gov/plantguide/pdf/pg_crlf.pdf (accessed on February 27, 2019).
112. Cale, J.A.; McNulty, S.A. Not dead yet: Beech trees can survive nearly three decades in the aftermath phase of a deadly forest disease complex. *For. Ecol. Manage.* **2018**, *409*, 372-377, doi: <https://doi.org/10.1016/j.foreco.2017.11.044>.
113. Cale, J.A.; Garrison-Johnston, M.T.; Teale, S.A.; Castello, J.D. Beech bark disease in North America: Over a century of research revisited. *For. Ecol. Manage.* **2017**, *394*, 86-103, doi: <https://doi.org/10.1016/j.foreco.2017.03.031>.
114. Hedges, G.S.; Hodgson, C.J. *Phalacrocooccus howertoni*, a new genus and species of soft scale (Hemiptera: Coccoidea) from Florida. *Fla. Entomol.* **2010**, *93*, 8-23, doi: <https://doi.org/10.1653/024.093.0102>.
115. Habeck, D.H.; Mead, F.W. *Edwards Wasp Moth, Lymire edwardsii (Grote) (Insecta: Lepidoptera: Arctiidae: Ctenuchinae)*; EENY-182; University of Florida, Institute of Food and Agricultural Sciences Extension: Gainesville, Florida, 2017.
116. Legaspi, J.C.; Mannion, C.; Amalin, D.; Jr., B.C.L. Biology, ecology and control of the Ficus whitefly, *Singhiella simplex* (Hemiptera: Aleyrodidae). In *Potential Invasive Pests of Agricultural Crops*, Pena, J.E., Ed. CABI: Wallingford, Oxfordshire, United Kingdom, and Boston, Massachusetts, 2013; pp. 363-372.
117. Henricot, B.; Prior, C. *Phytophthora ramorum*, the cause of sudden oak death or ramorum leaf blight and dieback. *Mycologist* **2004**, *18*, 151-156, doi: <https://doi.org/10.1017/S0269915X04004148>.
118. Tanis, S.R.; McCullough, D.G. Host Resistance of Five *Fraxinus* Species to *Agrilus planipennis* (Coleoptera: Buprestidae) and Effects of Paclobutrazol and Fertilization. *Environ. Entomol.* **2015**, *44*, 287-299, doi: <https://doi.org/10.1093/ee/nvu005>.
119. Garden, M.B. Ash Yellows and Decline. Available online: <http://www.missouribotanicalgarden.org/gardens-gardening/your-garden/help-for-the-home-gardener/advice-tips-resources/pests-and-problems/diseases/viruses/ash-yellows.aspx> (accessed on March 2, 2019).
120. University of Illinois Extension. White Ash: *Fraxinus americana*. Available online: <http://extension.illinois.edu/hortanswers/plantdetail.cfm?PlantID=209&PlantTypeID=7> (accessed on March 2, 2019).
121. University of Illinois Extension. Fungal Disease: Verticillium Wilt [Shrubs and Trees]: *Verticillium albo-atrum* and *V. dahliae*. Available online: <http://extension.illinois.edu/hortanswers/detailProblem.cfm?PathogenID=26> (accessed on March 2, 2019).
122. Gilman, E.F.; Watson, D.G. *Fraxinus pennsylvanica: Green Ash*; ENH-425; University of Florida, Institute of Food and Agricultural Sciences Extension: Gainesville, Florida, 2006; p 3.

123. Griffiths, H.M.; Sinclair, W.A.; Smart, C.D.; Davis, R.E. The phytoplasma associated with ash yellows and lilac witches'-broom: '*Candidatus Phytoplasma fraxini*'. *Int. J. Syst. Evol. Micr.* **1999**, *49*, 1605-1614, doi: <https://doi.org/10.1099/00207713-49-4-1605>.
124. Sutton, B.C.; Hodges, C.S. Revision of *Cercospora*-like fungi on *Juniperus* and related conifers. *Mycologia* **1990**, *82*, 313-325, doi: <https://doi.org/10.2307/3759902>.
125. Leahy, R.M. *Cercosporidium Blight of Leyland Cypress and Related Conifers*; Plant Pathology Circular No. 397; Florida Department of Agriculture and Consumer Services, Division of Plant Industry: Gainesville, Florida, 2000; p 4.
126. Geils, B.W.; Moody, B.H.; Tovar, J.C. *Mistletoes of North American Conifers*; United States Department of Agriculture, Forest Service, Rocky Mountain Research Station: Ogden, Utah, 2002; p 123.
127. Hodges, G.S.; Ruter, J.M.; Braman, S.K. Susceptibility of *Ilex* Species, Hybrids and Cultivars to Florida Wax Scale (*Ceroplastes floridensis* Comstock). *J. Environ. Hortic.* **2001**, *19*, 32-36.
128. Conrad, A.O.; Taylor, N.J.; Bonello, P.E. *Thousand Cankers Disease*; PLPATH-TREE-07; The Ohio State University, Ohio State University Extension: Columbus, Ohio, 2016.
129. Seybold, S.; Haugen, D.; O'Brien, J.; Graves, A. *Pest Alert: Thousand Cankers Disease*; NA-PR-02-10; United States Department of Agriculture, Forest Service, Northeastern Area State and Private Forestry: Newtown Square, Pennsylvania, 2013; p 2.
130. United States Department of Agriculture Forest Service; United States Department of Agriculture Plant Protection and Quarantine. Thousand Cankers Disease Survey Guidelines for 2018. Available online: https://www.aphis.usda.gov/plant_health/plant_pest_info/tcd/downloads/TCD_Survey_Guidelines.pdf (accessed on March 2, 2019).
131. Linit, M. Insects Pests Of Black Walunt. In *Nut Production Handbook for Eastern Black Walnut*, Jones, J.E., Mueller, R., Van Sambeek, J.W., Eds. Southwest Missouri Resources, Conservation & Development (RC&D), Inc.: St. Paul, Minnesota, 1998; pp. 120-123.
132. Moffis, B.L.; Arthurs, S.P. *Bagworm, Thyridopteryx ephemeraeformis Haworth (Insecta: Lepidoptera: Psychidae)*; EENY-548; University of Florida, Institute of Food and Agricultural Sciences Extension: Gainesville, FL, 2016.
133. Missouri Botanical Garden. *Juniperus ashei*. Available online: <http://www.missouribotanicalgarden.org/PlantFinder/PlantFinderDetails.aspx?taxonid=279618&isprofile=0&> (accessed on March 2, 2019).
134. DeGomez, T.; Hayes, C.J.; Anhold, J.A.; McMillin, J.D.; Clancy, K.M. Evaluation of insecticides for protecting Arizona cypress (*Cupressus arizonica*) and one-seed juniper (*Juniperus monosperma*) from attack by *Phloeosinus* bark beetles. *Arboriculture and Urban Forestry* **2007**, *33*, 162-167.
135. Schmitt, C.L.; Parmeter, J.R.; Kliejunas, J.T. *Annosus Root Disease of Western Conifers*; Forest Insect & Disease Leaflet 172; United States Department of Agriculture, Forest Service: Washington, D.C., 2000; p 9.
136. Weber, D.J.; Bunderson, E.D.; Davis, J.N.; Nelson, D.L.; Hreha, A. Diseases and Environmental Factors of the Pinyon-Juniper Communities In *Proceedings: ecology and management of pinyon-juniper communities within the Interior West*, Monsen, S.B., Stevens, R., Eds. United States Department of Agriculture, Forest Service, Rocky Mountain Research Station: Ogden, Utah, 1999; Vol. Proceedings RMRS-P-9, pp. 118-121.
137. Durham, G. Juniper Pocket Rot (*Pyrofomes demidoffii*). Available online: <http://forestry.nv.gov/forestry-resources/forest-health/juniper-pocket-rot/> (accessed on March 2, 2019).
138. Kim, M.-S.; Klopfenstein, N.B.; McDonald, G.I. Effects of Forest Management Practices and Environment on Occurrence of *Armillaria* Species. *Journal of the Korean Forest Society* **2010**, *99*, 251-257.
139. Jetton, R.M. Defoliation tolerance and resistance to the forest tent caterpillar, *Malacosoma disstria* Hbn., in sweetgum, *Liquidambar styraciflua* L. North Carolina State University, Raleigh, NC, 2002.
140. Jetton, R.M.; Robison, D.J. Effects of artificial defoliation on growth and biomass accumulation in short-rotation sweetgum (*Liquidambar styraciflua*) in North Carolina. *J. Insect. Sci.* **2014**, *14*, 107, doi: <https://doi.org/10.1673/031.014.107>.
141. Stocks, I. *The Mexican Lac Scale Tachardiella mexicana* (Comstock) (Hemiptera: Kerriidae); EENY-552; University of Florida, Institute of Food and Agricultural Sciences Extension: Gainesville, Florida, 2016; p 3.

142. *A Compendium of Verticillium Wilts in Tree Species*. Hiemstra, J.A.; Harris, D.C., Eds.; Commision of European Communities: Wageningen, The Netherlands, 1998; p 80.
143. United States Department of Agriculture Natural Resources Conservation Service. Plant Fact Sheet: Southern Crab Apple. Available online: https://plants.usda.gov/factsheet/pdf/fs_maan3.pdf (accessed on March 2, 2019).
144. United States Department of Agriculture Natural Resources Conservation Service. Plant Fact Sheet: Swamp Tupelo. Available online: http://plants.usda.gov/factsheet/pdf/fs_nybi.pdf (accessed on February 28, 2019).
145. Dixon, W.N.; Woodruff, R.E.; Foltz, J.L. *Black Twig Borer, Xylosandrus compactus (Eichhoff)* (Insecta: Coleoptera: Curculionidae: Scolytinae; EENY-311; University of Florida, Institute of Food and Agricultural Sciences: Gainesville, Florida, 2005; p 5.
146. Culin, J.D.; Gorsuch, C.S.; Pizzuto, T.M. Natural History and Recommendations for Control of the Rhododendron Stem Borer, *Oberea myops*. *Journal American Rhododendron Society* **1993**, *47*, 206-209.
147. Martin, C.A. *Parkinsonia florida* (formerly *Cercidium floridum*). Available online: <http://www.public.asu.edu/~camartin/plants/Plant%20html%20files/parkinsoniaflorida.html> (accessed on February 27, 2019).
148. Schuch, U.; Kelly, J. *Mesquite and Palo Verde Trees for the Urban Landscape*; AZ1429; The University of Arizona Cooperative Extension: Tucson, Arizona, 2007; p 9.
149. Shields, J.; Jose, S.; Freeman, J.; Bunyan, M.; Celis, G.; Hagan, D.; Morgan, M.; Pietersen, E.C.; Zak, J. Short-Term Impacts of Laurel Wilt on Redbay (*Persea borbonia* [L.] Spreng.) in a Mixed Evergreen-Deciduous Forest in Northern Florida. *J. Forest.* **2011**, *109*, 82-88.
150. McDonald, G.I.; Martin, N.E.; Harvey, A.E. *Armillaria in the northern Rockies : pathogenicity and host susceptibility on pristine and disturbed sites*; Research Note INT-371; United States Department of Agriculture, Forest Service, Intermountain Research Station: Ogden, UT, 1987; p 12.
151. Arno, S.F.; Hoff, R.J. *Silvics of whitebark pine (Pinus albicaulis)*; Gen. Tech. Rep. INT-GTR-253; United States Department of Agriculture, Forest Service, Intermountain Research Station: Ogden, Utah, 1989.
152. Coop, J.D.; Schoettle, A.W. Fire and high-elevation, five-needle pine (*Pinus aristata* & *P. flexilis*) ecosystems in the southern Rocky Mountains: What do we know? In *The future of high-elevation, five-needle white pines in Western North America: Proceedings of the High Five Symposium*, Keane, R.E., Tombak, D.F., Murray, M.P., Smith, C.M., Eds. United States Department of Agriculture, Forest Service, Rocky Mountain Research Station: Fort Collins, CO, 2011; Vol. Proceedings RMRS-P-63, pp. 164-173.
153. Lanier, G.N. Biosystematics of North American Ips (Coleoptera: Scolytidae): Hopping's Group IX. *Can. Entomol.* **1970**, *102*, 1139-1163, doi: <https://doi.org/10.4039/Ent1021139-9>.
154. Gomez, D.F.; Hulcr, J. *Southern pine beetle, Dendroctonus frontalis Zimmermann* (Insecta: Coleoptera: Scolytidae); EENY-176; University of Florida, Institute of Food and Agricultural Sciences Extension: Gainesville, Florida, 2019; p 8.
155. Coyle, D.R.; Self, A.B.; Floyd, J.D.; Riggins, J.J. *Ips Bark Beetles in the Southesastern U.S.*; SREF-FH-002 Southern Regional Extension Forestry: Athens, Georgia, 2016; p 9.
156. Miller, D.R.; Borden, J.H. Life History and Biology of *Ips latidens* (LeConte) (Coleoptera: Scolytidae). *Can. Entomol.* **1985**, *117*, 859-871, doi: <https://doi.org/10.4039/Ent117859-7>.
157. Camilli, K.S.; Marshall, J.; Owen, D.; Gordon, T.; Wood, D. *Pitch Canker Disease in California*; Tree Note 32; California Department of Forestry and Fire Protection: Sacramento, California, 2013; p 9.
158. Virginia Department of Forestry. *Forest Health Review*; Virginia Department of Forestry: Charlottesville, Virginia, 2016; p 11.
159. Russell, L.M. Notes on *Euphalerus nidifex* Schwarz and Related Nest-Making, New World Psyllids (Homoptera: Psyllidae). *Fla. Entomol.* **1971**, *54*, 3-12.
160. Townsend, K.R. The Dutch elm disease. Boston University, Boston, Massachusetts, 1950.
161. University of Illinois Extension. Dutch Elm Disease. Available online: <http://extension.illinois.edu/focus/index.cfm?problem=dutch-elm-disease> (accessed on March 1, 2019).
162. Boland, J.M. The impact of an invasive ambrosia beetle on the riparian habitats of the Tijuana River Valley, California. *PeerJ* **2016**, *4*, e2141, doi: <https://doi.org/10.7717/peerj.2141>.

163. Solomon, J.D. *Guide to Insect Borers in North American Broadleaf Trees and Shrubs.*; Agriculture Handbook 706; United States Department of Agriculture, Forest Service: Washington, D.C., 1995.
164. Olsen, M. Diseases of Urban Plants in Arizona. Available online: <http://warnerstreesurgery.com/pdf/Diseases%20of%20Urban%20Plants.pdf> (accessed on August 12).
165. Van Driesche, R.G.; LaForest, J.H.; Bargeron, C.T.; Reardon, R.C.; Herlihy, M. *Forest Pest Insects in North America: A Photographic Guide*; FHTET-2012-02; United States Department of Agriculture, Forest Service, Forest Health Technology Enterprise Team: Morgantown, West Virginia, 2013; p 702.
166. Trouillas, F.P.; Úrbez-Torres, J.R.; Gubler, W.D. Diversity of diatrypaceous fungi associated with grapevine canker diseases in California. *Mycologia* **2010**, *102*, 319-336, doi: <https://doi.org/10.3852/08-185>.
167. Blackford, D. Management Guide for Forest Tent Caterpillar. In *Forest Insect and Disease Management Guide for the Northern and Central Rocky Mountains*, United States Department of Agriculture, Forest Service, Northern and Intermountain Regions, State and Private Forestry: Missoula, Montana, and Boise, Idaho, 2005; p. 4.
168. Worrall, J.J.; Egeland, L.; Eager, T.; Mask, R.A.; Johnson, E.W.; Kemp, P.A.; Shepperd, W.D. Rapid mortality of *Populus tremuloides* in southwestern Colorado, USA. *Forest Ecol. Manag.* **2008**, *255*, 686-696, doi: <https://doi.org/10.1016/j.foreco.2007.09.071>.
169. Zegler, T.J.; Moore, M.M.; Fairweather, M.L.; Ireland, K.B.; Fulé, P.Z. *Populus tremuloides* mortality near the southwestern edge of its range. *Forest Ecol. Manag.* **2012**, *282*, 196-207, doi: <https://doi.org/10.1016/j.foreco.2012.07.004>.
170. Marchetti, S.B.; Worrall, J.J.; Eager, T. Secondary insects and diseases contribute to sudden aspen decline in southwestern Colorado, USA. *Can. J. Forest Res.* **2011**, *41*, 2315-2325, doi: <https://doi.org/10.1139/x11-106>.
171. Trummer, L.; Hennon, P. *Hazard Trees in Alaska: A Guide to the Identification and Management of Hazard Trees in Alaska*; R10-TP-142; United States Forest Service, Alaska Region, State and Private Forestry: Anchorage, Alaska, 2009; p 62.
172. Guo, Y.H.; Cheng, Z.-M.; Walla, J.A.; Zhang, Z. Large-scale Screening for X-disease Phytoplasma Infection in Chokecherry. *HortScience* **1998**, *33*, 293-295.
173. Guo, Y.H.; Walla, J.A.; Cheng, Z.-M.; Lee, I.-M. X-Disease Confirmation and Distribution in Chokecherry in North Dakota. *Plant Dis.* **1996**, *80*, 95-102, doi: <https://doi.org/10.1094/PD-80-0095>.
174. Gibson, K. Management Guide for Flatheaded Fir Borers In *Forest Insect and Disease Management Guide for the Northern and Central Rocky Mountains*, United States Department of Agriculture, Forest Service, Northern and Intermountain Regions, State and Private Forestry: Missoula, Montana, and Boise, Idaho, 2004; p. 2.
175. O'Brien, J.; Mielke, M.; Starkey, D.; Juzwik, J. *How to Identify, Prevent, and Control Oak Wilt*; NA-FR-01-11; United States Department of Agriculture, Forest Service, Northern Area State & Private Forestry: Newtown Square, Pennsylvania, 2000; p 31.
176. Coleman, T.W.; Graves, A.D.; Hodde, M.; Heath, Z.; Chen, Y.; Flint, M.L.; Seybold, S.J. Forest stand composition and impacts associated with *Agrilus auroguttatus* Schaeffer (Coleoptera: Buprestidae) and *Agrilus coxalis* Waterhouse in oak woodlands. *Forest Ecol. Manag.* **2012**, *276*, 104-117, doi: <https://doi.org/10.1016/j.foreco.2012.03.011>.
177. Pritchard, E.A.; Beer, R.E. Biology and Control of *Asterolecanium* Scales on Oaks in California. *J. Econ. Entomol.* **1950**, *43*, 494-497, doi: <https://doi.org/10.1093/jee/43.4.494>.
178. Swiecki, T.J.; Bernhardt, E. *A field guide to insects and diseases of California oaks*; Gen. Tech Rep. PSW-GTR-197; United States Department of Agriculture, Forest Service, Pacific Southwest Research Station: Albany, CA, 2006; p 151.
179. Coleman, T.W.; Seybold, S.J. Collection History and Comparison of the Interactions of the Goldspotted Oak Borer, *Agrilus auroguttatus* Schaeffer (Coleoptera: Buprestidae), with Host Oaks in Southern California and Southeastern Arizona, U.S.A. *Coleopts. Bull.* **2011**, *65*, 93-108, doi: <https://doi.org/10.1649/072.065.0224>.
180. Brasier, C.M.; Denman, S.; Rose, J.; Kirk, S.A.; Hughes, K.J.D.; Griffin, R.L.; Lane, C.R.; Inman, A.J.; Webber, J.F. First report of ramorum bleeding canker on *Quercus falcata*, caused by *Phytophthora ramorum*. *Plant Pathol.* **2004**, *53*, 804-804, doi: <https://doi.org/10.1111/j.1365-3059.2004.01079.x>.

181. Sever, A.; Cranshaw, W.; Brudenell, R. *Agrilus quercicola* (Fisher) (Coleoptera: Buprestidae), the Gambel Oak Borer, as a Pest of *Quercus* spp. *Southwest. Entomol.* **2012**, *37*, 147-150, doi: <https://doi.org/10.3958/059.037.0207>.
182. Donley, D.E.; Acciavatti, R.E. *Red Oak Borer*; Forest Insect & Disease Leaflet 163; United States Department of Agriculture, Forest Service, Northern Area State & Private Forestry: Newtown Square, Pennsylvania, 1980; p 9.
183. Harrington, T.C.; McNew, D.; Yun, H.Y. Bur oak blight, a new disease on *Quercus macrocarpa* caused by *Tubakia iowensis* sp. nov. *Mycologia* **2012**, *104*, 79-92, doi: <https://doi.org/10.3852/11-112>.
184. Walter, J.A.; Finch, F.T.; Johnson, D.M. Re-evaluating fall cankerworm management thresholds for urban and suburban forests. *Agr. Forest Entomol.* **2016**, *18*, 145-150, doi: <https://doi.org/10.1111/afe.12147>.
185. Haack, R.A.; Acciavatti, R.E. *Twolined Chestnut Borer*; Forest Insect & Disease Leaflet 168; United States Department of Agriculture, Forest Service: Washington, D.C., 1992; p 12.
186. Goldman, D.H. Plant Guide: Live Oak. Available online: https://plants.usda.gov/plantguide/pdf/pg_quvi.pdf (accessed on February 28, 2019).
187. Rehm, A.; Humm, H.J. *Sphaeroma terebrans*: A Threat to the Mangroves of Southwestern Florida. *Science* **1973**, *182*, 173-174, doi: <https://doi.org/10.1126/science.182.4108.173>.
188. Pasiecznik, N. Invasive Species Compendium: *Rhizophora mangle* (Red Mangrove). Available online: <https://www.cabi.org/isc/datasheet/47509> (accessed on March 1, 2019).
189. Olexa, M.T.; Freeman, T.E. A Gall Disease of Red Mangrove Caused by *Cylindrocarpion didymum*. *Plant Dis. Rep.* **1978**, *62*, 283-286.
190. Woodruff, R.E. A Mangrove Borer, *Poecilips rhizophorae* (Hopkins)(Coleoptera: Scolytidae). *Entomology Circular* **1970**, *98*.
191. Broschat, T.K. *Roystonea regia*: Royal Palm; ENH-732; University of Florida, Institute of Food and Agricultural Sciences Extension: Gainesville, Florida, 2014; p 4.
192. Broschat, T.K. *Sabal palmetto*: Sabal or Cabbage Palm; ENH-733; University of Florida, Institute of Food and Agricultural Sciences Extension: Gainesville, Florida, 2013; p 5.
193. Eskalen, A.; R., S.; S.C., L.; Rugman-Jones, P.F.; Twizeyimana, M.; A., G.; T., T. Shot Hole Borer Reproductive Hosts. Available online: https://ucanr.edu/sites/pshb/overview/SHB_Reproductive_Hosts/ (accessed on March 1, 2019).
194. Billings, R.F.; Grosman, D.M.; Pase, H.A. Soapberry Borer, *Agrilus prionurus* (Coleoptera: Buprestidae): An Exotic Pest Threatens Western Soapberry in Texas. *Southeast. Nat.* **2014**, *13*, 105-116, doi: <https://doi.org/10.1656/058.013.s510>.
195. Kendra, P.E.; Montgomery, W.S.; Niogret, J.; Epsky, N.D. An Uncertain Future for American Lauraceae: A Lethal Threat from Redbay Ambrosia Beetle and Laurel Wilt Disease (A Review). *Am. J. Plant Sci.* **2013**, Vol.04No.03, *12*, doi: <https://doi.org/10.4236/ajps.2013.43A092>.
196. Mead, F.W. *False-Mastic Psylla, Ceropsylla sideroxyli Riley* (Insecta: Hemiptera: Psyllidae); EENY-370; University of Florida, Institute of Food and Agricultural Sciences Extension: Gainesville, Florida, 2015; p 3.
197. Howard, F.W.; Pemberton, R.W.; Hodges, G.S.; Steinberg, B.; McLean, D.; Liu, H. Host plant range of lobate lac scale, *Paratachardina*, in Florida. *Proc. Fl. State Hortic.* **2006**, *119*, 398-408.
198. Dixon, W.N. *Cypress Looper Anacampodes pergracilis* (Hulst) (Insecta: Lepidoptera: Geometridae); EENY-303; University of Florida, Institute of Food and Agricultural Sciences Extension: Gainesville, Florida, 2016; p 3.
199. Espinosa, A.; Bowman, H.; Hodges, A.; Hodges, G. *Boisduval Scale, Diaspis boisduvalii Signoret* (Insecta: Hemiptera: Diaspididae); EENY-467; University of Florida, Institute of Food and Agricultural Sciences Extension: Gainesville, Florida, 2016; p 5.
200. Smith, J.A.; O'Donnell, K.; Mount, L.L.; Shin, K.; Peacock, K.; Trulock, A.; Spector, T.; Cruse-Sanders, J.; Determann, R. A Novel Fusarium Species Causes a Canker Disease of the Critically Endangered Conifer, *Torreya taxifolia*. *Plant Dis.* **2011**, *95*, 633-639, doi: <https://doi.org/10.1094/PDIS-10-10-0703>.
201. United States Department of Agriculture Forest Service. *Pest Alert: Hemlock Borer*; NA-PR-03-00; United States Department of Agriculture, Forest Service, Northeastern Area State and Private Forestry: 2000; p 2.
202. Wilson, C.; Vendettuoli, J.; Orwig, D.; Preisser, E. Impact of an Invasive Insect and Plant Defense on a Native Forest Defoliator. *Insects* **2016**, *7*, 45, doi: <https://doi.org/10.3390/insects7030045>.

203. Hadfield, J.S.; Goheen, D.J.; Filip, G.M.; Schmitt, C.L.; Harvey, R.D. *Root Diseases in Oregon and Washington Conifers*; R6-FPM-250-86; United States Department of Agriculture, Forest Service, Pacific Northwest Region: Portland, Oregon, 1986.
204. University of Amsterdam. Dutch elm disease on CD-ROM. Available online: <http://www.dutchelmdisease.org/EXPERT/DED/CORE/00/00/80.HTML> (accessed on February 28, 2019).
205. Fraedrich, S.W. California Laurel Is Susceptible to Laurel Wilt Caused by *Raffaelea lauricola*. *Plant Dis.* **2008**, *92*, 1469-1469, doi: <https://doi.org/10.1094/PDIS-92-10-1469A>.
206. Gilbertson, R.L. Wood-Rotting Fungi of North America. *Mycologia* **1980**, *72*, 1-49, doi: <https://doi.org/10.2307/3759417>.

Table S2. The 339 insect, disease, and parasitic higher plant agents identified as among the most important threats to each of 419 North American tree species. Nonnative invasive agents are indicated with *. The list includes the guilds of the insect agents, and the number of host tree species identified for each agent.

Agent	Guild	Hosts
Insects		
Alder bark beetle (<i>Alniphagus aspericollis</i>)	Phloem- or Wood-Borers	1
Asian longhorned beetle (<i>Anoplophora glabripennis</i>)*	Phloem- or Wood-Borers	43
Asiatic oak weevil (<i>Cyrtepistomus castaneus</i>)*	Foliage-Feeders	3
Bagworm (<i>Thyridopteryx ephemeraeformis</i>)	Foliage-Feeders	4
Baldcypress leafroller (<i>Archips goyerana</i>)	Foliage-Feeders	1
Balsam woolly adelgid (<i>Adelges piceae</i>)*	Sap-Feeders	5
Banded elm bark beetle (<i>Scolytus schevyrewi</i>)*	Phloem- or Wood-Borers	2
Bark beetle (<i>Ips spinifer</i>)	Phloem- or Wood-Borers	1
Bark beetle (<i>Liparthrum arizonicum</i>)	Phloem- or Wood-Borers	1
Bark beetle (<i>Orthotomicus spinifer</i>)	Phloem- or Wood-Borers	1
Birch leafminer (<i>Fenusia pusilla</i>)*	Foliage-Feeders	3
Black stem borer (<i>Xylosandrus germanus</i>)*	Phloem- or Wood-Borers	1
Black twig borer (<i>Xylosandrus compactus</i>)*	Phloem- or Wood-Borers	2
Black vine weevil (<i>Otiorhynchus sulcatus</i>)*	Foliage-Feeders	2
Black-horned juniper borer (<i>Callidium texanum</i>)	Phloem- or Wood-Borers	1
Blue spruce engraver (<i>Ips hunteri</i>)	Phloem- or Wood-Borers	1
Boisduval Scale (<i>Diaspis boisduvalii</i>)	Sap-Feeders	2
Borer (<i>Megacyllene antennata</i>)	Phloem- or Wood-Borers	3
Bronze birch borer (<i>Agrilus anxius</i>)	Phloem- or Wood-Borers	4
Bronze poplar borer (<i>Agrilus liragus</i>)	Phloem- or Wood-Borers	1
Cactus longhorn beetle (<i>Moneilema gigas</i>)	Phloem- or Wood-Borers	1
Calico scale (<i>Eulecanium cerasorum</i>)*	Sap-Feeders	3
California fivespined ips (<i>Ips paraconfusus</i>)	Phloem- or Wood-Borers	1
California flatheaded borer (<i>Melanophila californica</i>)	Phloem- or Wood-Borers	1
California oakworm (<i>Phryganidea californica</i>)	Foliage-Feeders	1
Cardin's whitefly (<i>Metaleurodicus cardini</i>)*	Sap-Feeders	1
Catalpa sphinx moth (<i>Ceratomia catalpae</i>)	Foliage-Feeders	2
Cedar bark beetle (<i>Phloeosinus spp.</i>)	Phloem- or Wood-Borers	7
Cherry bark tortrix (<i>Enarmonia formosana</i>)*	Phloem- or Wood-Borers	8
Chestnut gall wasp (<i>Dryocosmus kuriphilus</i>)*	Foliage-Feeders	1
Cottonwood borer (<i>Plectrodera scalator</i>)	Phloem- or Wood-Borers	2
Cottonwood leaf beetle (<i>Chrysomela scripta</i>)	Foliage-Feeders	4
Cypress looper (<i>Anacamptodes pergracilis</i>)	Foliage-Feeders	2
Dogwood borer (<i>Synanthedon scitula</i>)	Phloem- or Wood-Borers	1
Dogwood clubgall midge (<i>Resseliella clavula</i>)	Foliage-Feeders	1
Dogwood twig borer (<i>Oberea tripunctata</i>)	Phloem- or Wood-Borers	1
Douglas-fir beetle (<i>Dendroctonus pseudotsugae</i>)	Phloem- or Wood-Borers	1
Douglas-fir tussock moth (<i>Orgyia pseudotsugata</i>)	Foliage-Feeders	2

Agent	Guild	Hosts
Eastern juniper bark beetle (<i>Phloeosinus dentatus</i>)	Phloem- or Wood-Borers	2
Eastern tent caterpillar (<i>Malacosoma americanum</i>)	Foliage-Feeders	7
Edwards wasp moth (<i>Lymire edwardsii</i>)	Foliage-Feeders	2
Elongate hemlock scale (<i>Fiorinia externa</i>)*	Sap-Feeders	2
Emerald ash borer (<i>Agrilus planipennis</i>)*	Phloem- or Wood-Borers	15
European pine sawfly (<i>Neodiprion sertifer</i>)*	Foliage-Feeders	1
Fall cankerworm (<i>Alsophila pometaria</i>)	Foliage-Feeders	2
False-mastic psylla (<i>Ceropsylla sideroxyli</i>)	Sap-Feeders	1
Ficus whitefly (<i>Singhiella simplex</i>)*	Sap-Feeders	2
Fiddlewood leafroller (<i>Epicorsia oedipodalis</i>)	Foliage-Feeders	2
Fir engraver (<i>Scolytus ventralis</i>)	Phloem- or Wood-Borers	4
Flatheaded apple tree borer (<i>Chrysobothris femorata</i>)	Phloem- or Wood-Borers	1
Flatheaded fir borer (<i>Melanophila drummondi</i>)	Phloem- or Wood-Borers	1
Florida wax scale (<i>Ceroplastes floridensis</i>)	Sap-Feeders	2
Forest tent caterpillar (<i>Malacosoma disstria</i>)	Foliage-Feeders	28
Gambel oak borer (<i>Agrilus quercicola</i>)	Phloem- or Wood-Borers	1
Geiger tortoise beetle (<i>Eurypepla calochroma</i>)	Foliage-Feeders	1
Geometrid moth (<i>Stamnodes animata</i>)*	Foliage-Feeders	1
Giant palm borer (<i>Dinapate wrightii</i>)	Phloem- or Wood-Borers	1
Goldspotted oak borer (<i>Agrilus auroguttatus</i>)	Phloem- or Wood-Borers	7
Green alder sawfly (<i>Monsoma pulveratum</i>)*	Foliage-Feeders	2
Gypsy moth (<i>Lymantria dispar dispar</i>)*	Foliage-Feeders	65
Hackberry butterfly (<i>Asterocampa celtis</i>)	Foliage-Feeders	1
Hackberry engraver (<i>Scolytus muticus</i>)	Phloem- or Wood-Borers	1
Hackberry nipplegall maker (<i>Pachypsylia celtidismamma</i>)*	Foliage-Feeders	1
Hemlock borer (<i>Melanophila fulvoguttata</i>)	Phloem- or Wood-Borers	2
Hemlock looper (<i>Lambdina fiscellaria</i>)	Foliage-Feeders	2
Hemlock woolly adelgid (<i>Adelges tsugae</i>)*	Sap-Feeders	2
Hickory bark beetle (<i>Scolytus quadrispinosus</i>)	Phloem- or Wood-Borers	8
Hickory borer (<i>Goes pulcher</i>)	Phloem- or Wood-Borers	5
Hickory spiral borer (<i>Agrilus arcuatus torquatus</i>)	Phloem- or Wood-Borers	2
Honeylocust agrilus (<i>Agrilus difficilis</i>)	Phloem- or Wood-Borers	1
Jack pine budworm (<i>Choristoneura pinus</i>)	Foliage-Feeders	1
Japanese beetle (<i>Popillia japonica</i>)*	Foliage-Feeders	14
Jeffrey pine beetle (<i>Dendroctonus jeffreyi</i>)	Phloem- or Wood-Borers	1
Juniper borer (<i>Atimia huachucae</i>)	Phloem- or Wood-Borers	1
Juniper scale (<i>Carulaspis juniperi</i>)*	Sap-Feeders	16
Juniper twig pruner (<i>Styloxus bicolor</i>)	Phloem- or Wood-Borers	1
Lac scale (<i>Tachardiella mexicana</i>)	Sap-Feeders	1
Larch casebearer (<i>Coleophora laricella</i>)*	Foliage-Feeders	2
Larch sawfly (<i>Pristiphora erichsonii</i>)	Foliage-Feeders	3
Linden looper (<i>Erranis tiliaria</i>)	Foliage-Feeders	1
Lobate lac scale (<i>Paratachardina pseudolobata</i>)*	Sap-Feeders	6

Agent	Guild	Hosts
Locust borer (<i>Megacyllene robiniae</i>)	Phloem- or Wood-Borers	2
Locust leafminer (<i>Odontota dorsalis</i>)	Foliage-Feeders	2
Locust twig borer (<i>Ecdytolopha insiticiana</i>)	Phloem- or Wood-Borers	1
Madrone thrips (<i>Thrips madronii</i>)	Foliage-Feeders	1
Magnolia scale (<i>Neolecanium cornuparvum</i>)	Sap-Feeders	3
Mahogany shoot borer (<i>Hypsipyla grandella</i>)	Phloem- or Wood-Borers	1
Mimosa webworm (<i>Homadaula anisocentra</i>)*	Foliage-Feeders	1
Monterey pine weevil (<i>Pissodes radiatae</i>)*	Phloem- or Wood-Borers	2
Mountain mahogany bark beetle (<i>Chaetophloeus heterodoxus</i>)	Phloem- or Wood-Borers	1
Mountain pine beetle (<i>Dendroctonus ponderosae</i>)	Phloem- or Wood-Borers	10
Mountain-mahogany looper (<i>Iridopsis clivinaria</i>)	Foliage-Feeders	1
North American pine engraver (<i>Ips pini</i>)	Phloem- or Wood-Borers	1
Oak pit scale (<i>Asterolecanium spp.</i>)*	Sap-Feeders	9
Oystershell scale (<i>Lepidosaphes ulmi</i>)*	Sap-Feeders	21
Palo verde root borers (<i>Derobrachus geminatus</i>)	Phloem- or Wood-Borers	1
Pecan carpenterworm (<i>Cossula magnifica</i>)	Phloem- or Wood-Borers	1
Pine engraver beetles (<i>Ips spp.</i>)	Phloem- or Wood-Borers	17
Pine shoot beetle (<i>Tomicus piniperda</i>)*	Phloem- or Wood-Borers	1
Pink hibiscus mealybug (<i>Maconellicoccus hirsutus</i>)*	Sap-Feeders	2
Pinon twig beetle (<i>Pityophthorus punctifrons</i>)	Phloem- or Wood-Borers	1
Pinyon ips (<i>Ips confusus</i>)	Phloem- or Wood-Borers	4
Pinyon needle scale (<i>Matsucoccus acalyptus</i>)	Sap-Feeders	6
Pinyon sawfly (<i>Neodiprion edulicolus</i>)	Foliage-Feeders	3
Polyphagous shot hole borer (<i>Euwallacea spp.</i>)*	Phloem- or Wood-Borers	6
Poplar borer (<i>Saperda calcarata</i>)	Phloem- or Wood-Borers	7
Poplar-and-willow borer (<i>Cryptorhynchus lapathi</i>)*	Phloem- or Wood-Borers	2
Psyllid (<i>Euphalerus nidifex</i>)	Foliage-Feeders	1
Red oak borer (<i>Enaphalodes rufulus</i>)	Phloem- or Wood-Borers	10
Red pine scale (<i>Matsucoccus resinosae</i>)*	Sap-Feeders	1
Redheaded pine sawfly (<i>Neodiprion lecontei</i>)	Foliage-Feeders	1
Rhododendron stem borer (<i>Oberea myops</i>)	Phloem- or Wood-Borers	1
Roundheaded pine beetle (<i>Dendroctonus adjunctus</i>)	Phloem- or Wood-Borers	1
Royal palm bug (<i>Xylastodoris luteolus</i>)	Sap-Feeders	1
Rugose spiraling whitefly (<i>Aleurodicus rugioperculatus</i>)*	Sap-Feeders	1
Sawfly (<i>Neodiprion spp.</i>)	Foliage-Feeders	4
Scale insect (<i>Phalacrocooccus howertoni</i>)	Sap-Feeders	3
Silver fir beetle (<i>Pseudohylesinus sericeus</i>)	Phloem- or Wood-Borers	1
Soapberry borer (<i>Agrilus prionurus</i>)*	Phloem- or Wood-Borers	1
Southern cypress bark beetle (<i>Phloeosinus taxodii</i>)	Phloem- or Wood-Borers	3
Southern pine beetle (<i>Dendroctonus frontalis</i>)	Phloem- or Wood-Borers	12
Spruce beetle (<i>Dendroctonus rufipennis</i>)	Phloem- or Wood-Borers	6
Spruce budworm (<i>Choristoneura fumiferana</i>)	Foliage-Feeders	5
Sri Lankan weevil (<i>Myllocerus undecimpustulatus undatus</i>)*	Foliage-Feeders	1

Agent	Guild	Hosts
Sycamore borer (<i>Synanthedon resplendens</i>)	Phloem- or Wood-Borers	1
Sycamore lace bug (<i>Corythucha ciliata</i>)	Foliage-Feeders	1
Tent caterpillar (<i>Malacosoma</i> spp.)	Foliage-Feeders	6
Thousand cankers disease (<i>Geosmithia morbida</i> / <i>Pityophthorus juglandis</i>)	Phloem- or Wood-Borers	6
True fir bark beetles (<i>Scolytus dentatus</i>)	Phloem- or Wood-Borers	1
Tuliptree scale (<i>Toumeyella liriodendri</i>)	Sap-Feeders	2
Tupelo leafminer (<i>Antispila nysaefoliella</i>)	Foliage-Feeders	4
Twig beetle (<i>Pityophthorus boycei</i>)	Phloem- or Wood-Borers	1
Twig beetles (<i>Pityophthorus</i> spp.)	Phloem- or Wood-Borers	1
Twig girdler (<i>Oncideres cingulata</i>)	Phloem- or Wood-Borers	9
Twig girdler (<i>Oncideres pustulatus</i>)	Phloem- or Wood-Borers	1
Twolined chestnut borer (<i>Agrilus bilineatus</i>)	Phloem- or Wood-Borers	31
Two-spotted spider mite (<i>Tetranychus urticae</i>)	Sap-Feeders	1
Virginia pine sawfly (<i>Neodiprion pratti</i>)	Foliage-Feeders	1
Walkingstick (<i>Diapheromera femorata</i>)	Foliage-Feeders	1
Walnut caterpillar (<i>Datana integerrima</i>)	Foliage-Feeders	1
Western balsam bark beetle (<i>Dryocoetes confusus</i>)	Phloem- or Wood-Borers	2
Western hemlock looper (<i>Lambdina fiscellaria lugubrosa</i>)	Foliage-Feeders	1
Western pine beetle (<i>Dendroctonus brevicomis</i>)	Phloem- or Wood-Borers	1
Western spruce budworm (<i>Choristoneura freemani</i>)	Foliage-Feeders	7
White pine weevil (<i>Pissodes strobi</i>)	Phloem- or Wood-Borers	6
Winter moth (<i>Operophtera brumata</i>)*	Foliage-Feeders	8
Wood borer (<i>Coccotrypes rhizophorae</i>)*	Phloem- or Wood-Borers	1
Wood borer (<i>Sphaeroma terebrans</i>)*	Phloem- or Wood-Borers	1
Wood-boring beetle (<i>Dicerca horni</i>)	Phloem- or Wood-Borers	1
Yellow poplar weevil (<i>Odontopus calceatus</i>)	Foliage-Feeders	1

Diseases

Alder canker (<i>Valsa melanodiscus</i>)	.	3
Anthracnose (<i>Apiognomonia veneta</i>)	.	3
Apple mosaic virus (ApMV)	.	2
Apple scab (<i>Venturia inaequalis</i>)	.	4
Arbutus canker (<i>Nattrassia mangiferae</i>)	.	2
Armillaria root disease (<i>Armillaria</i> spp.)	.	35
Artist's cork (<i>Ganoderma applanatum</i>)	.	11
Ash yellows (<i>Candidatus Phytoplasma fraxini</i>)	.	14
Bacterial leaf blight (<i>Pseudomonas lauraceum</i>)	.	1
Bacterial leaf scorch (<i>Xylella fastidiosa</i>)	.	15
Bacterial necrosis (<i>Erwinia cacticida</i>)	.	1
Bacterial spot (<i>Xanthomonas campestris</i> pv. <i>pruni</i>)	.	6
Basal canker and butt rot (<i>Kretzschmaria deusta</i>)	.	3
Beech bark disease (<i>Neonectria</i> spp./ <i>Cryptococcus fagisuga</i>)*	.	1
Birch rust (<i>Melampsoridium betulinum</i>)	.	1

Potter *et al.*, Important insect and disease threats to United States tree species, Table S2, page 5

Agent	Guild	Hosts
Black knot (<i>Apiosporina morbosa</i>)	.	12
Black root rot (<i>Thielaviopsis basicola</i>)	.	1
Blackline (Cherry leafroll virus)	.	1
Black-stain root disease (<i>Leptographium wageneri</i>)	.	1
Blister canker (<i>Biscogniauxia marginata</i>)	.	3
Bot canker (<i>Botryosphaeria corticola</i>)	.	1
Botryosphaeria canker (<i>Botryosphaeria ribis</i>)	.	2
Broom rust (<i>Gymnosporangium libocedri</i>)	.	1
Brown cubical rot (<i>Poria sequoiae</i>)	.	1
Brown spot needle blight (<i>Mycosphaerella dearnessii</i>)	.	1
Brown trunk rot (<i>Fomitopsis officinalis</i>)	.	1
Bunch disease (Phytoplasma strains in subgroup of Prunus X-disease group)	.	2
Bur oak blight (<i>Tubakia dryina</i>)	.	1
Butt rot (<i>Hericium erinaceus</i>)	.	2
Butt rot fungi (<i>Phellinus swieteniae</i>)	.	1
Butternut canker (<i>Sirococcus clavigignenti-juglandacearum</i>)*	.	1
Canadian pine scale (<i>Matsucoccus macrocicatrices</i>)/ <i>Caliciopsis pinea</i> complex	.	1
Canker (<i>Cryptosphaeria pullmanensis</i>)	.	1
Canker (<i>Phytophthora cambivora</i>)	.	1
Canker disease (<i>Fusarium</i> spp.)	.	1
Canker rot of oak (<i>Inonotus andersonii</i>)	.	8
Canker-rot (<i>Inonotus</i> spp.)	.	1
Catalpa midge (<i>Cecidomyia catalpae</i>)	.	2
Cercospora blight (<i>Asperisporium sequoiae</i>)	.	7
Cercospora blight of juniper (<i>Pseudocercospora juniperi</i>)	.	3
Cherry leaf spot (<i>Blumeriella jaapii</i>)	.	12
Chestnut blight (<i>Cryphonectria parasitica</i>)*	.	3
Clitocybe root rot (<i>Armillaria tabescens</i>)	.	1
Collar rot (<i>Phytophthora siskiyouensis</i>)	.	2
Comandra blister rust (<i>Cronartium comandrae</i>)	.	5
Coral spot fungus (<i>Nectria cinnabarina</i>)	.	2
Cotton root rot (<i>Phymatotrichopsis omnivora</i>)	.	2
Crown canker and root rot (<i>Phytophthora cactorum</i>)	.	4
Crown gall (<i>Agrobacterium tumefaciens</i>)	.	2
Cytospora canker (<i>Cytospora chrysosperma</i>)	.	2
Cytospora canker of spruce (<i>Cytospora kunzei</i>)	.	1
Decay (<i>Trametes versicolor</i>)	.	2
Didymascella leaf blight (<i>Didymascella thujina</i>)	.	1
Dogwood anthracnose (<i>Discula destructiva</i>)*	.	3
Dutch elm disease (<i>Ophiostoma novo-ulmi</i>)*	.	7
Elm yellows (<i>Candidatus Phytoplasma ulmi</i>)	.	6
Elytroderma needle cast (<i>Elytroderma deformans</i>)	.	4
Endothia canker (<i>Endothia gyrosa</i>)	.	7

Agent	Guild	Hosts
European larch canker (<i>Lachnellula willkommii</i>)*	.	1
Eutypella canker (<i>Eutypella parasitica</i>)	.	3
False smut of palms (<i>Graphiloa phoenicis</i>)	.	2
Fire blight (<i>Erwinia amylovora</i>)	.	11
Foliage diseases (<i>Bifusella</i> spp.)	.	1
Fusarium wilt (<i>Fusarium oxysporum</i> f. sp. <i>rhois</i>)	.	1
Fusiform rust (<i>Cronartium quercuum</i> f. sp. <i>fusiforme</i>)	.	5
Gall disease (<i>Cylindrocarpon didymum</i>)	.	1
Gall rust (<i>Cronartium quercuum</i>)	.	3
Gall rust of Virginia pine (<i>Cronartium quercuum</i> f. sp. <i>virginianae</i>)	.	1
Ganoderma butt rot (<i>Ganoderma zonatum</i>)	.	3
Ganoderma root and butt rot (<i>Ganoderma australe</i>)	.	1
Ganoderma root and butt rot (<i>Ganoderma meredithae</i>)	.	1
Ganoderma rot of hardwoods (<i>Ganoderma lucidum</i>)	.	4
Guignardia blotch (<i>Guignardia aesculi</i>)	.	2
Hackberry witches-broom complex (<i>Podosphaera phytoptophila/Aceria celtis</i>)	.	2
Hawthorn rust (<i>Gymnosporangium globosum</i>)	.	1
Heart rot (<i>Flammulina velutipes</i>)	.	1
Heart rot (<i>Inonotus glomeratus</i>)	.	2
Heart rot fungi (<i>Phellinus robiniae</i>)	.	2
Heterobasidion root disease (<i>Heterobasidion</i> spp.)	.	21
Heterobasidion root disease P-type (<i>Heterobasidion irregularare</i>)	.	12
Hispidus canker (<i>Inonotus hispidus</i>)	.	4
Hornbeam anthracnose (<i>Apiosporopsis carpinea</i>)	.	1
Hypoxylon canker of aspen (<i>Entoleuca mammata</i>)	.	1
Hypoxylon canker of oak (<i>Biscogniauxia atropunctata</i> var. <i>atropunctata</i>)	.	55
Indian paint fungus (<i>Echinodontium tinctorium</i>)	.	4
Juniper pocket rot (<i>Pyrofomes demidoffii</i>)	.	2
Laminated root rot (<i>Phellinus sulphurascens</i>)	.	8
Laminated root rot (<i>Phellinus weiri</i>)	.	1
Larch needle blight (<i>Hypodermella laricis</i>)	.	1
Laurel wilt (<i>Raffaelea lauricola</i>)/redbay ambrosia beetle (<i>Xyleborus glabratus</i>)*	.	3
Leaf blight and fruit rot (<i>Monilinia johnsonii</i>)	.	6
Leaf curl and witches broom (<i>Taphrina wiesneri</i>)	.	2
Leaf spot disease (<i>Cristulariella depraedans</i>)	.	1
Limb rust (<i>Cronartium arizonicum</i>)	.	1
Limb rust (<i>Cronartium</i> sp.)	.	1
Lophodermium needle cast (<i>Lophodermium</i> spp.)	.	1
Madrone canker (<i>Neofusicoccum arbuti</i>)	.	2
Melampsora leaf rust (<i>Melampsora medusae</i>)	.	1
Nectria canker (<i>Nectria ditissima</i>)	.	18
Nectria canker (<i>Nectria magnoliae</i>)	.	1
Nectria canker (<i>Nectria</i> sp.)	.	1

Agent	Guild	Hosts
Nectriella gall (<i>Nectriella pironii</i>)	.	1
Needle cast (<i>Lirula nervisequia</i> var. <i>conspicua</i>)	.	1
Nematode (<i>Rotylenchulus reniformis</i>)	.	1
Oak bracket (<i>Inonotus dryadeus</i>)	.	4
Oak wilt (<i>Bretziella fagacearum</i>)*	.	61
Persimmon wilt (<i>Acremonium diospyri</i>)	.	1
Pezicula trunk and branch cankers (<i>Pezicula subcarnea</i>)	.	1
Phytophthora disease of alders (<i>Phytophthora</i> sp.)	.	3
Phytophthora root rot (<i>Phytophthora cinnamomi</i>)*	.	8
Pinyon blister rust (<i>Cronartium occidentale</i>)	.	6
Pit canker (<i>Phytophthora inflata</i>)	.	1
Pitch canker (<i>Fusarium circinatum</i>)*	.	10
Pocket dry rot (<i>Tyromyces amarus</i>)	.	1
Poplar rust (<i>Melampsora abietis-canadensis</i>)	.	4
Poplar rust (<i>Melampsora occidentalis</i>)	.	1
Port-Orford-cedar root disease (<i>Phytophthora lateralis</i>)*	.	2
Powdery mildew (<i>Erysiphe flexuosa</i>)	.	3
Powdery mildew (<i>Phyllactinia guttata</i>)	.	1
Prunus X-disease (<i>Phytoplasma</i>)	.	2
Red heart fungus (<i>Stereum sanguinolentum</i>)	.	1
Red ring rot (<i>Phellinus pini</i> var. <i>cancriformans</i>)	.	1
Red ring rot (<i>Phellinus pini</i>)	.	3
Red ring rot canker (<i>Phenillus cancriformans</i>)	.	1
Red root and butt rot (<i>Onnia tomentosa</i>)	.	7
Red-belted fungus (<i>Fomitopsis pinicola</i>)	.	2
Root rot (<i>Cylindrocladium scoparium</i>)	.	1
Root-knot nematode (<i>Meloidogyne</i> spp.)	.	1
Rust (<i>Gymnosporangium cunninghamianum</i>)	.	3
Rust (<i>Gymnosporangium</i> spp.)	.	8
Rust (<i>Ravenelia</i> spp.)	.	1
Rust (<i>Ravenelia versatilis</i>)	.	1
Rust fungus (<i>Melampsorella caryophyllacearum</i>)	.	1
Scab and black canker of willow (<i>Venturia saliciperda/Glomerella miyabeana</i>)*	.	13
Schweinitzii root and butt rot (<i>Phaeolus schweinitzii</i>)	.	6
Scleroderris canker (<i>Gremmeniella abietina</i>)*	.	1
Seiridium canker (<i>Seiridium cardinale</i>)	.	13
Sooty bark canker (<i>Encoelia pruinosa</i>)	.	1
Sooty canker (<i>Fusicoccum mangiferae</i>)	.	2
Spiculosia canker (<i>Phellinus spiculosus</i>)	.	6
Stem canker (<i>Fusarium solani</i>)	.	5
Sterile conk trunk rot (<i>Inonotus obliquus</i>)	.	3
Stigmina leaf spot (<i>Stigmina palmivora</i>)	.	1
Stunt nematode (<i>Tylenchorhynchus</i> spp.)	.	1

Agent	Guild	Hosts
Sudden oak death (<i>Phytophthora ramorum</i>)*	.	12
Sycamore canker (<i>Ceratocystis fimbriata</i> f. <i>platani</i>)	.	1
Tar spot (<i>Diatractium cordianum</i>)	.	1
Texas Phoenix palm decline (<i>Candidatus Phytoplasma palmae</i>)	.	2
Thielaviopsis trunk rot (<i>Thielaviopsis paradoxa</i>)	.	3
Thyronectria canker (<i>Pleonectria austroamericana</i>)	.	2
Tinder conk (<i>Fomes fomentarius</i>)	.	1
Valsa canker and dieback (<i>Valsa abietis</i>)	.	1
Verticillium wilt (<i>Verticillium albo-atrum</i>)	.	15
Verticillium wilt (<i>Verticillium dahliae</i>)	.	4
Verticillium wilt (<i>Verticillium</i> spp.)	.	5
Western gall rust (<i>Endocronartium harknessii</i>)	.	4
White mottled rot (<i>Ganoderma</i> spp.)	.	13
White pine blister rust (<i>Cronartium ribicola</i>)*	.	9
White pine needle cast (<i>Lophodermella arcuata</i>)	.	1
White ring rot (<i>Obba rivulosa</i>)	.	1
White rot (<i>Bjerkandera adusta</i>)	.	1
White rot (<i>Phellinus badius</i>)	.	1
White rot (<i>Phellinus weiriatus</i>)	.	1
White spongy rot (<i>Phellinus igniarius</i>)	.	6
Willow witches'-broom (<i>Phytoplasma</i> , members of the yellows group)	.	2
Witches broom (<i>Phytoplasma</i>)	.	1
Witches'-broom (<i>Apiosporina collinsii</i>)	.	3
Wood decay (<i>Phellinus gilvus</i>)	.	2
Wood rotting (<i>Phanerochaete tuberculata</i>)	.	1
<i>Parasitic Higher Plants</i>		
Apache dwarf mistletoe (<i>Arceuthobium apachecum</i>)	.	1
Coastal dwarf mistletoe (<i>Arceuthobium littorum</i>)	.	1
Dwarf mistletoe (<i>Arceuthobium californicum</i>)	.	1
Dwarf mistletoe (<i>Arceuthobium gillii</i>)	.	1
Dwarf mistletoe (<i>Arceuthobium monticola</i>)	.	1
Dwarf mistletoe (<i>Arceuthobium occidentale</i>)	.	2
Dwarf mistletoe (<i>Arceuthobium</i> spp.)	.	6
Eastern dwarf mistletoe (<i>Arceuthobium pusillum</i>)	.	3
Eastern mistletoe (<i>Phoradendron leucarpum</i>)	.	10
Hemlock dwarf mistletoe (<i>Arceuthobium tsugense</i>)	.	2
Juniper mistletoe (<i>Phoradendron juniperinum</i>)	.	5
Larch dwarf mistletoe (<i>Arceuthobium laricis</i>)	.	1
Limber pine dwarf mistletoe (<i>Arceuthobium cyanocarpum</i>)	.	4
Lodgepole pine dwarf mistletoe (<i>Arceuthobium americanum</i>)	.	2
Mistletoe (<i>Phoradendron bolleanum</i>)	.	6
Mistletoe (<i>Phoradendron californicum</i>)	.	8

Agent	Guild	Hosts
Pinyon dwarf mistletoe (<i>Arceuthobium divaricatum</i>)	.	7
Red fir dwarf mistletoe (<i>Arceuthobium abietinum</i> f. sp. <i>magnifica</i>)	.	2
Western dwarf mistletoe (<i>Arceuthobium campylopodum</i>)	.	3
Western spruce dwarf mistletoe (<i>Arceuthobium microcarpum</i>)	.	2




Figure S1. Hotspots of Importance Value at Risk (IVAR) for plots within 834 km² hexagons, encompassing both native and exotic insect and disease agents. G^* scores ≤ -2 (red) are significant clusters of low survivability scores (and therefore hotspots of IVAR, the opposite of survivability); scores > 2 (blue) are significant clusters of high survivability values (coldspots of IVAR). (a) Hotspots of IVAR for insect agents. (b) Hotspots of IVAR for disease agents.




Figure S2. Mean percent host tree Importance Value at Risk (IVAR) for plots within 834 km² hexagons, encompassing insect agents. (a) IVAR for native insects. (b) IVAR for exotic insects.




Figure S3. Mean percent host tree Importance Value at Risk (IVAR) for plots within 834 km² hexagons, encompassing insect agents. (a) IVAR for phloem- or wood-borers. (b) IVAR for sap-feeders. (c) IVAR for foliage-feeders.




Figure S4. Hotspots of Importance Value at Risk (IVAR) for plots within 834 km² hexagons, encompassing insect agents. G_i^* scores ≤ -2 (red) are significant clusters of low survivability scores (and therefore hotspots of IVAR, the opposite of survivability); scores > 2 (blue) are significant clusters of high survivability values (coldspots of IVAR). (a) Hotspots of IVAR for phloem- or wood-borers. (b) Hotspots of IVAR for sap-feeders. (c) Hotspots of IVAR for foliage-feeders.