

Beneficial Microorganisms and Water Stress Influence *Quercus ilex* Seedlings' Response to *Phytophthora cinnamomi* Rands

Roberto J. Cabrera-Puerto, Francisco J. Ruiz-Gómez * and Rafael M. Navarro-Cerrillo

Supplementary material

Table S1. Beneficial microorganisms composition of T-complex.

Species	Concentration UFC/ml	Taxa
<i>Trichoderma viride</i>	2-4 10 ⁵	<i>Fungi</i>
<i>Trichoderma harzianum</i>	2-4 10 ⁵	<i>Fungi</i>
<i>Azospirillum brasilense</i>	10 ⁸ -10 ⁹	<i>Rizobacterias</i>
<i>Rhizobium tropici</i>	10 ⁸ -10 ⁹	<i>Rizobacterias</i>
<i>Pantoea ananatis</i>	10 ⁸ -10 ⁹	<i>Rizobacterias</i>
<i>Pseudomonas putida</i>	10 ⁸ -10 ⁹	<i>Rizobacterias</i>
<i>Pseudomonas fluorescens</i>	10 ⁸ -10 ⁹	<i>Rizobacterias</i>
<i>Pseudomonas brassicacearum</i>	10 ⁸ -10 ⁹	<i>Rizobacterias</i>
<i>Bacillus aryabhaktai</i>	10 ⁸ -10 ⁹	<i>Rizobacterias</i>




Figure S1: Pearson correlations of studied variables. Color intensity and size by importance of Pearson value. Positive correlations in blue and negative correlations in red.