

**Table S1.** Effect of maternal characteristics and severity COVID-19 on oxidative stress markers.

	<b>B</b>	<b>95 CI%</b>	<b>p-value</b>	<b>R2/(R2 adjusted)</b>
<b>Malondialdehyde</b>				
Constant	36.7.	14.9 to 58.5	0.001	
Maternal age	-0.032	-0.434 to 0.370	0.874	0.0338 (-0.03529)
GA at diagnosis	-0.170	-0.528 to 0.188	0.344	
pBMI	-0.141	-0.527 to 0.245	0.468	
Severe COVID-19	-2.17	-6.60 to 2.26	0.330	
<b>Carbonylated proteins</b>				
Constant	12604	5077 to 20132	0.001	
Maternal age	-160	-299 to -21.3	0.025	0.1932 (0.1311)
GA at diagnosis	-1.38	-125 to 122	0.982	
pBMI	-65.5	-199 to 67.8	0.330	
Severe COVID-19	2032	504 to 3561	0.010	
<b>Total antioxidant capacity</b>				
Constant	43.8	-4.24 to 91.8	0.075	
Maternal age	0.052	-0.833 to 0.938	0.906	0.2240 (0.1643)
GA at diagnosis	0.071	-0.714 to 0.861	0.852	
pBMI	-0.213	-1.063 to 0.638	0.618	
Severe COVID-19	18.1	8.26 to 27.8	0.001	

Models included 54 COVID-19 pregnant women; non-severe COVID-19 ( $n = 40$ ) and severe COVID-19 ( $n = 17$ ); GA: Gestational age; pBMI: Pregestational body mass index.