

Supplementary Materials

pH-Responsive Drug Delivery Nanoplatfoms as Smart Carriers of Unsymmetrical Bisacridines for Targeted Cancer Therapy

**Joanna Pilch ^{1,*}, Agnieszka Potęga ¹, Agata Kowalczyk ², Artur Kasprzak ³,
Patrycja Kowalik ^{2,3}, Piotr Bujak ³, Ewa Paluszkiewicz ¹, Ewa Augustin ¹
and Anna M. Nowicka ^{2,*}**

¹ Faculty of Chemistry, Gdańsk University of Technology, Narutowicza Street 11/12,
80-233 Gdańsk, Poland

² Faculty of Chemistry, University of Warsaw, Pasteura Street 1, 02-093 Warsaw, Poland

³ Faculty of Chemistry, Warsaw University of Technology, Noakowskiego Street 3,
00-664 Warsaw, Poland

* Correspondence: joanna.pilch@pg.edu.pl (J.P.); anowicka@chem.uw.edu.pl (A.M.N.)




Figure S1. UV-Vis titration curves obtained for β -CD(C-2028) complex in various pH: 7.4 (A), 6.2 (B), 5.2 (C), and 4.0 (D).




Figure S2. Intracellular fluorescence images of acidic organelles loaded with LysoTracker Red DND-99 in H460, Du-145, LNCaP, MRC-5, and PNT1A cells incubated with QD_{green}, QD_{green}-β-CD-FA, β-CD for 72 h. The scale bar is 20 μm.