

Solid lipid nanoparticles containing Morin: preparation, characterization, and *ex vivo* permeation studies.

Federica De Gaetano¹, Consuelo Celesti², Giuseppe Paladini³, Valentina Venuti⁴, Maria Chiara Cristiano⁵, Donatella Paolino⁶, Daniela Iannazzo², Vincenza Strano⁷, Anna Gueli³, Silvana Tommasini¹, Cinzia Anna Ventura^{1*}, Rosanna Stancanelli¹

¹ Department of Chemical, Biological, Pharmaceutical and Environmental Sciences, University of Messina, V.le Ferdinando Stagno d'Alcontres 31, I-98166 Messina, Italy; fedegaetano@unime.it; stommasini@unime.it; rstancanelli@unime.it

² Department of Engineering, University of Messina, Contrada Di Dio, I-98166 Messina, Italy; ccelesti@unime.it

³ Department of Physics and Astronomy "Ettore Majorana", University of Catania, Via S. Sofia, 64 I-95123 Catania, Italy; anna.gueli@unict.it

⁴ Department of Mathematical and Computer Sciences, Physical Sciences and Earth Sciences, University of Messina, V.le Ferdinando Stagno D'Alcontres 31, I-98166 Messina, Italy; vvenuti@unime.it

⁵ Department of Medical and Surgical Sciences University of Catanzaro "Magna Graecia", V.le Europa s.n.c., I-88100 Catanzaro, Italy; mchiara.cristiano@unicz.it

⁶ Department of Experimental and Clinical Medicine, University of Catanzaro "Magna Graecia", V.le Europa s.n.c., I-88100 Catanzaro, Italy; paolino@unicz.it

⁷ CNR-IMM, Catania (University) Unit, Via S. Sofia 64, I-95123 Catania, Italy; vincenzina.strano@ct.infn.it

* Correspondence: cavenatura@unime.it (C.A.V.)




Figure S1. MRN chromatogram (100 µg/mL concentration). See material and methods section for the operative conditions.




Figure S2. Plot of peak area vs. MRN concentration