


**Supplementary Data:**

**Evaluation of Cytotoxicity, Release Behavior and Phytopathogens Control by Mancozeb-Loaded Guar Gum Nanoemulsions for Sustainable Agriculture**


Ravinder Kumar <sup>1,\*</sup>, Manju Nehra <sup>2</sup>, Dharmender Kumar <sup>3</sup>, Baljeet Singh Saharan <sup>4</sup>, Prince Chawla <sup>5</sup>, Pardeep Kumar Sadh <sup>1</sup>, Anju Manuja <sup>6</sup>, Joginder Singh Duhan <sup>1,\*</sup>

**Table S1.** Runs to optimize the size of guar gum NEs using response surface methodology.


Run	Factor 1	Factor 2	Response
	Concentrations		
	A:Guar gum %	B:Mancozeb %	Particle size
	w/v	w/v	nm
1	1	0.5	356.7
2	1.5	1.5	394.1
3	0.5	0.5	429.9
4	0.5	1	547
5	1	1	208.1
6	1.5	1.5	205.9
7	0.5	1.5	320.3
8	1.5	0.5	368.3
9	0.5	1.5	559
10	1	1	360.9
11	1	1.5	591.5
12	1	1	460.3
13	1.5	1	246.6
14	1	1	460.4
15	1.5	0.5	356.7
16	1	1	368.3
17	0.5	0.5	460.4




(a)



(b)



(c)



(d)

**Figure S1.** In vitro antifungal ability of blank (N3), and fungicide-loaded (N3F) GG NEs against (a) *A. alternata* (b) *S. lycopersici* (c) *A. solani* (d) *S. sclerotiorum*; where: C-control, PC-pure control (media only), N3-blank NEs, N3F-mancozeb (1.0 mg/ml) encapsulated NEs and F- commercial fungicide at 0.5, 1.0 and 1.5 ppm.



**Figure S2.** Disease controlling efficacy of GG NEs on tomato plants (a) early blight (b) leaf spot; on potato plants (c) early blight (d) stem rot; in pot conditions; where C-control (No treatment), CP1-P1, F-fungicide, FP1-fungicide+P1, N3-blank NEs, N3P1-blank NEs+P1, N3F-fungicide-loaded NEs, N3FP1-fungicide loaded NEs+P1, CP2-pathogen P2, FP2-fungicide+P2, N3P2-blank NEs+P2, N3FP2-fungicide-loaded NEs+P2.