

Evaluating the Effects of Climate Change and Human Activities on the Seasonal Trends and Spatial Heterogeneity of Soil Moisture

Ermei Zhang^{1,2}, Yujie Liu^{1,2,*}, Tao Pan^{1,2}, Qinghua Tan^{1,2}, Zhiang Ma¹

¹ Key Laboratory of Land Surface Pattern and Simulation, Institute of Geographic Sciences and Natural Resources Research, Chinese Academy of Sciences, Beijing 100101, China

² University of Chinese Academy of Sciences, Beijing 100049, China

* Corresponding author at: No.11A, Datun Road, Chaoyang District, Beijing, China

E-mail address: liuyujie@igsrr.ac.cn (Yujie Liu)




Figure S1. Land covers on the Tibetan Plateau in 2020. The legend numbers correspond to the land cover codes

Table S1. Lookup table of land cover classes in the Tibetan Plateau

Code	Name	Number in original data	Code	Name	Number in original data
10	Tillage	12	33	Low coverage grassland	33
		12 Dry land			Low coverage grassland

20	Woodland	21 Forested land, 23 Sparse woodland, 24 Other woodland	40	Water	41 River channel, 42 Lake, 43 Reservoir pit, 44 Permanent glacier, 46 Shoal, 64 Wetlands
22	Shrub	22 Shrubbery	50	Construction land	51 Urban land, 52 Rural resident, 53 Other construction land, 61 Sandy land、62 Gobi,
31	High coverage grassland	31 High coverage grassland	60	Sand, Gobi and Bare land	63 Saline-alkali land, 65 Bare land, 66 Bare rock texture
32	Medium coverage grassland	32 Medium coverage grassland	67	Desert	67 Other unused land, including alpine desert, tundra, etc.




Figure S2. Annual variation of soil moisture from 2000 to 2020. * was the significance of 0.05 level.




Figure S3. Spatial distribution of mean annual soil moisture of the Tibetan Plateau from 2000 to 2020.




Figure S4. Spatial distribution of fractional vegetation coverage in the Tibetan Plateau from 2000 to 2020.