

Supplementary table 1. Median values and interquartile ranges for the Psychological General Well-Being (PGWB) and Gastrointestinal Symptoms Rating Scale (GSRS) totals and subscores for the dermatitis herpetiformis (DH) patients and the coeliac disease controls with and without fractures. In the PGWB, a higher score indicates a better quality of life, and in the GSRS, a higher score indicates more severe symptoms.

	DH patients (n=222)			Coeliac disease controls (n=129)		
	With fracture (n=45)	Without fracture (n=177)	p-value	With fracture (n=35)	Without fracture (n=94)	p-value
GSRS						
Total	1.9 (1.3–2.3)	1.6 (1.3–2.1)	0.191	1.9 (1.5–2.5)	1.7 (1.4–2.5)	0.472
Diarrhoea	1.7 (1.0–2.3)	1.3 (1.0–2.0)	0.116	2.0 (1.0–2.7)	1.7 (1.0–2.5)	0.635
Indigestion	2.0 (1.3–2.5)*	1.8 (1.5–2.5)	0.630	2.4 (1.8–2.8)	2.0 (1.5–3.0)	0.343
Constipation	1.3 (1.0–2.3)	1.3 (1.0–2.3)	0.568	1.7 (1.3–2.5)	1.7 (1.0–2.7)	0.572
Pain	1.7 (1.2–2.3)	1.3 (1.0–2.0)	0.130	1.7 (1.3–2.3)	1.7 (1.3–2.3)	0.517
Reflux	1.5 (1.0–2.0)	1.0 (1.0–1.5)	0.012	1.5 (1.0–2.5)	1.0 (1.0–2.0)	0.083
PGWB						
Total	106 (94–113)	112 (101–119)	0.006	110 (93–120)	106 (97–116)	0.629
Anxiety	25 (22–27)	26 (23–28)	0.020	27 (21–29)	25 (23–28)	0.757
Depression	17 (16–18)	18 (16–18)	0.311	17 (15–18)	17 (15–18)	0.948
Well-being	17 (15–19)	18 (16–20)	0.007	18 (15–19)	17 (16–20)	0.941
Self control	16 (15–17)	16 (15–17)	0.052	16 (13–17)	16 (14–17)	0.888
General health	13 (11–15)	15 (13–16)	0.012	13 (11–16)	13 (11–15)	0.712
Vitality	18 (17–21)	20 (17–21)	0.029	19 (17–20)	18 (16–20)	0.665

*p < 0.05 when the DH patients with fractures were compared to the coeliac disease patients with fractures