

Nutrients

Supplementary Material

Preparation of red ginseng marc-derived gintonin and its application as a skin nutrient

Rami Lee ¹, Ji-Hun Kim ¹, Hongik Hwang ², Hyewhon Rhim ³, Sung-Hee Hwang ⁴, Ik-Hyun Cho ⁵, Do-Geun Kim ⁶, Hyoung-Chun Kim ⁷ and Seung-Yeol Nah ^{1,*}

¹ Ginsentology Research Laboratory and Department of Physiology, College of Veterinary Medicine, Konkuk University, Seoul 05029, Republic of Korea; rmlee12@konkuk.ac.kr (R.L.); bioskjh@naver.com (J.-H.K.)

² Department of Life Science, University of Seoul, Seoul 02504, Republic of Korea; hongik@uos.ac.kr

³ Center for Neuroscience, Korea Institute of Science and Technology, Seoul 02792, Republic of Korea; hrhim@kist.re.kr

⁴ Department of Pharmaceutical Engineering, College of Health Sciences, Sangji University, Wonju 26339, Republic of Korea; sunghhwang@sangji.ac.kr

⁵ Department of Convergence Korean Medical Science, College of Korean Medicine, Kyung Hee University, Seoul, 02447, Republic of Korea; ihcho@khu.ac.kr

⁶ Korea Brain Research Institute (KBRI), 61, Cheomdan-ro, Dong-gu, Daegu, 41062, Republic of Korea; kimvet0911@kbri.re.kr

⁷ Neuropsychopharmacology and Toxicology Program, College of Pharmacy, Kangwon National University, Chunchon 24341, Republic of Korea; kimhc@kangwon.ac.kr

*Correspondence: synah@konkuk.ac.kr; Tel.: +82-2-450-4154




Figure S1.

A possible process for the lysophosphatidic acids (LPAs) formation from phospholipids such as phosphatidic acid (PA) or phosphatidylcholine (PC) in KRGM gintonin.