

Lipidome profiling in childhood obesity compared to adults: a pilot study

Andrea Soria-Gondek^{1*}, Pablo Fernández-García^{2*}, Lorena González³, Marjorie Reyes-Farias^{3,4}, Marta Murillo⁵, Aina Valls⁵, Nativitat Real⁶, Silvia Pellitero⁷, Jordi Tarascó⁸, Benjamin Jenkins⁹, María Galán², Francesc Villarroya^{10,11}, Albert Koulman⁹, Patricia Corrales², Antonio Vidal-Puig^{12†}, Rubén Cereijo^{10,11, 13†}, David Sánchez-Infantes^{2,3,11†}

Contact info:

David Sánchez-Infantes; Department of Health Sciences, Campus Alcorcón, University Rey Juan Carlos (URJC), E-28922 Madrid, Spain. Electronic address: david.sanchezi@urjc.es

Supplementary file S6. Heatmap representation of correlation coefficient (r) values for linear regressions between the differences in fatty liver index (FLI) and those of selected top significantly regulated lipids coincident in the pediatric (PE) and adult (AD) cohorts (see main Figure 3) 6 months after metabolic intervention regarding the baseline ($\Delta 6M-0M$). Blue cells indicate positive ($r > 0$) and red cells indicate negative associations ($r < 0$). *: Statistically significant correlation ($p < 0.05$).

