

Biometabolites of *Citrus unshiu* Peel Enhance Intestinal Permeability and Alter Gut Commensal Bacteria

Se-Hui Lee ¹, Dongju Seo ¹, Kang-Hee Lee ¹, So-Jung Park ¹, Sun Park ¹, Hyeyun Kim ¹, Taekyung Kim ², In Hwan Joo ³, Jong-Min Park ³, Yun-Hwan Kang ⁴, Gah-Hyun Lim ¹, Dong Hee Kim ³ and Jin-Young Yang ^{1,*}

¹ Mucosal Immunology Laboratory, Department of Integrated Biological Science, Pusan National University, Busan 46241, Republic of Korea; two_hui@pusan.ac.kr (S.-H.L.); csa83201@pusan.ac.kr (D.S.); rkdgm19056@pusan.ac.kr (K.-H.L.); ahejr26@naver.com (S.-J.P.); sun980610@pusan.ac.kr (S.P.); kelly5060@pusan.ac.kr (H.K.); ghlim16@pusan.ac.kr (G.-H.L.)

² Department of Biology Education, Pusan National University, Busan 46241, Republic of Korea; taekim@pusan.ac.kr

³ Department of Pathology, College of Korean Medicine, Daejeon University, Daejeon 34520, Republic of Korea; jooih@dju.kr (I.H.J.); jmpark@dju.kr (J.-M.P.); dhkim@dju.kr (D.H.K.)

⁴ Department of Industry Promotion, National Institute for Korean Medicine Development, Geongsan 38540, Republic of Korea; k_yunhwan@nikom.or.kr

* Correspondence: jyyang99@pusan.ac.kr; Tel.: +82-51-510-2286; Fax: +82-51-581-2962

Supplementary Table

Table S1. List of commensal bacteria (species). All individual gut microbes shown in the pie chart in Figure 6D are arranged in order of their frequency here.


 <i>Alistioides outredinis</i>	 <i>Alistioides onderdonkii</i>	 <i>Nealecta timonensis</i>
 <i>Hunotatella xylanolytica</i>	 <i>Parabacteroides goldsteinii</i>	 <i>Ruminiclostridium cellulolyticum</i>
 <i>Muribaculum intestinale</i>	 <i>Lawsonibacter asaccharolyticus</i>	 <i>Anaerosporeobacter mobilis</i>
 <i>Kineothrix alysoides</i>	 <i>Oscillibacter valericigenes</i>	 <i>Anaerostipes butyraticus</i>
 <i>Alistipes seneqalensis</i>	 <i>[Clostridium] populeti</i>	 <i>Anaerotruncus rubiinfantis</i>
 <i>Flintibacter butyricus</i>	 <i>Enterocloster aldenensis</i>	 <i>Hydrogenoanaerobacterium saccharovorans</i>
 <i>Lacrimispora saccharolytica</i>	 <i>Petrocella atlantisensis</i>	 <i>Butyrivibrio faecihominis</i>
 <i>Mucispirillum schaedleri</i>	 <i>Parabacteroides merdae</i>	 <i>Roseburia faecis</i>
 <i>Christensenella massiliensis</i>	 <i>Gracilibacter thermotolerans</i>	 <i>Parasutterella excrementihominis</i>
 <i>Duncaniella freteri</i>	 <i>Vallitalea prouvensis</i>	 <i>Acetivibrio straminisolvans</i>
 <i>Lacrimispora aerotolerans</i>	 <i>Parabacteroides distasonis</i>	 <i>[Bacteroides] pectinophilus</i>
 <i>Phocaeicola massiliensis</i>	 <i>Anaeroplasmata abactoclasticum</i>	 <i>Alkalibacter saccharofermentans</i>
 <i>Oscillibacter ruminantium</i>	 <i>Phocaea massiliensis</i>	 <i>Enterocloster asparaqiformis</i>
 <i>Christensenella hongkongensis</i>	 <i>Anaerotruncus propionicum</i>	 <i>Lactobacillus johnsonii</i>
 <i>Lactobacillus apodemi</i>	 <i>[Clostridium] viride</i>	 <i>Lachnoclostridium pacaense</i>
 <i>Clostridium cellulovorans</i>	 <i>[Clostridium] polysaccharolyticum</i>	 <i>Desulfotobacterium hafniense</i>
 <i>Acetivibrio thermocellus</i>	 <i>Eisenbergiella massiliensis</i>	 <i>[Eubacterium] siraeum</i>
 <i>Vampirovibrio chlorellavorus</i>	 <i>Bacteroides nordii</i>	 <i>Caecibacterium sporiformans</i>
 <i>Acetatifactor muris</i>	 <i>Desulfovibrio desulfuricans</i>	 <i>Eubacterium xylanophilum</i>
 <i>Butyrivibrio phoceensis</i>	 <i>Maoeeibacillus indolicus</i>	 ETC (<0.01%)
 <i>Intestinimonas butyriciproducens</i>	 <i>Anaerotaenia torta</i>	

Table S2. List of specific primer sets used for Real-Time PCR in this study.

Target Gene	Forward Sequence	Reverse Sequence
<i>GAPDH</i>	CAATGACCCCTTCATTGACC	GACAAGCTTCCCGTTCTCAG
<i>Occludin</i>	CCTCCAATGGCAAAGTGAAT	CTCCCCACCTGTCGTGTAGT
<i>E-cadherin</i>	CACCTGGAGAGAGGCCATGT	TGGGAAACATGAGCAGCTCT
<i>Claudin-2</i>	TATGTGGTGCCAGCATTGT	TCATGCCCACCACAGAGATA
<i>ZO-1</i>	CCACCTCTGTCCAGCTCTTC	CACCGGAGTGATGGTTTTCT