

**Table S1** Participant flow of enrolled infants with dietary data at baseline and follow-ups.

<b>Number of subjects with dietary data at baseline and follow-ups</b>																	
<b>Age (months)</b>	<b>1*</b>	<b>2*</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>7</b>	<b>8</b>	<b>9</b>	<b>12</b>	<b>18</b>	<b>24</b>	<b>36</b>	<b>48</b>	<b>60</b>	<b>72</b>	<b>96</b>
	1,049	1,213	1,188	1,144	1,131	1,098	1,018	990	971	946	797	820	576	535	492	527	447
<b>Number of subjects excluded due to...</b>																	
<b>Inaccurate eating frequency** or illness</b>	6	1	9	12	18	18	20	25	36	30	28	29	19	7	9	7	3
<b>Breastfeeding***</b>	427	379	340	324	297	269	215	188	152	87	27	13	1	--	--	--	--
<b>Study population N = 1,244</b>																	
<b>Subjects</b>	616	833	839	808	815	812	784	778	783	829	742	778	556	528	483	520	444

\*Infants were enrolled during the first 2 months of life. \*\*Inaccurate eating frequency defined as missing time information or as food records, in which EF was not assigned to one particular point in time, e.g. “during day”. \*\*\* Number of breastfed infants were included in the sub analysis resulting in a total of 1,370 infants and a total of 536 breastfed infants at a given time point.