

Supplementary Materials for:

Fetal alcohol growth restriction is independent of infant feeding practices and postnatal alcohol exposure in a prospective South African birth cohort

Alexia C. Edwards, Sandra W. Jacobson, Marjanne Senekal, Neil C. Dodge, Christopher D. Molteno, Ernesta M. Meintjes, Joseph L. Jacobson, R. Colin Carter

Table S1. Univariate relations between control variables and infant feeding practices^a

	Maternal age	Gravidity	Maternal education	Cigarettes/day	Days/mo marijuana use	Days/mo methamphetamine use	Weeks gestation at delivery	Infant sex
<u>Infant feeding practices</u>								
Weeks breastfed	0.06	0.01	0.04	-0.08	-0.03	-0.01	-0.16 [†]	0.09
Weeks formula fed	-0.01	-0.03	-0.15 [†]	0.04	-0.06	0.01	0.03	-0.06
Weeks complementary foods	-0.08	-0.12	0.12	-0.18*	-0.05	0.02	0.04	-0.03
Weeks exclusively breastfed	0.00	0.04	0.15 [†]	-0.04	0.08	0.03	-0.12	0.04
Weeks exclusively formula fed	-0.06	-0.02	-0.01	0.07	0.07	0.04	0.08	-0.09
Weeks mixed feeding	0.04	-0.04	-0.13	-0.02	-0.11	-0.03	0.01	0.03
Reporting insufficient breastmilk	-0.09	-0.08	-0.11	0.07	-0.05	0.14	-0.08	-0.03
<u>Complementary food provision</u>								
Juice	-0.09	-0.08	0.03	0.031	-0.02	-0.07	0.06	0.01
Porridge	0.04	0.10	-0.14	-0.04	-0.08	-0.01	-0.08	-0.13
Fruits	-0.04	-0.03	-0.01	-0.11	0.05	0.01	0.03	0.02
Vegetables	-0.01	-0.06	-0.04	-0.21*	-0.09	0.09	0.14	-0.03
Eggs	0.04	0.13	-0.01	0.06	-0.04	-0.11	-0.11	0.01
Chicken	0.17 [†]	0.14	-0.15 [†]	-0.05	-0.08	-0.04	-0.07	-0.11
Fries	-0.05	-0.09	-0.07	-0.02	0.07	-0.01	0.00	0.11

^ap < 0.10; *p < 0.05; **p < 0.01; ***p < 0.001.

^aValues are Pearson r's