

Review

# The Diversity of Cyanobacterial Toxins on Structural Characterization, Distribution and Identification: A Systematic Review

Xingde Du <sup>1</sup>, Haohao Liu <sup>1</sup>, Le Yuan <sup>1</sup>, Yueqin Wang <sup>1</sup>, Ya Ma <sup>1</sup>, Rui Wang <sup>1</sup>, Xinghai Chen <sup>2</sup>, Michael D. Losiewicz <sup>2</sup>, Hongxiang Guo <sup>3,\*</sup> and Huizhen Zhang <sup>1,\*</sup>

<sup>1</sup> College of Public Health, Zhengzhou University, Zhengzhou 450001, China; duxd1993@gs.zzu.edu.cn (X.D.); lhl2019@gs.zzu.edu.cn (H.L.); ylsir2018@gs.zzu.edu.cn (L.Y.); wangyueqin@gs.zzu.edu.cn (Y.W.); maya@gs.zzu.edu.cn (Y.M.); wangrui2019@gs.zzu.edu.cn (R.W.)

<sup>2</sup> Department of Chemistry and Biochemistry, St Mary's University, San Antonio, TX 78228, USA; xchen@stmarytx.edu (X.C.); mlosiewicz@stmarytx.edu (M.D.L.)

<sup>3</sup> College of Life Sciences, Henan Agricultural University, Zhengzhou 450002, China

\* Correspondence: huizhenzhang@zsu.edu.cn (H.Z.); guohongxiang@henau.edu.cn (H.G.); Tel.: +86-151-8835-7252 (H.Z.); +86-136-4386-7952 (H.G.)

Received: 27 August 2019; Accepted: 9 September 2019; Published: 12 September 2019



**Abstract:** The widespread distribution of cyanobacteria in the aquatic environment is increasing the risk of water pollution caused by cyanotoxins, which poses a serious threat to human health. However, the structural characterization, distribution and identification techniques of cyanotoxins have not been comprehensively reviewed in previous studies. This paper aims to elaborate the existing information systematically on the diversity of cyanotoxins to identify valuable research avenues. According to the chemical structure, cyanotoxins are mainly classified into cyclic peptides, alkaloids, lipopeptides, nonprotein amino acids and lipoglycans. In terms of global distribution, the amount of cyanotoxins are unbalanced in different areas. The diversity of cyanotoxins is more obviously found in many developed countries than that in undeveloped countries. Moreover, the threat of cyanotoxins has promoted the development of identification and detection technology. Many emerging methods have been developed to detect cyanotoxins in the environment. This communication provides a comprehensive review of the diversity of cyanotoxins, and the detection and identification technology was discussed. This detailed information will be a valuable resource for identifying the various types of cyanotoxins which threaten the environment of different areas. The ability to accurately identify specific cyanotoxins is an obvious and essential aspect of cyanobacterial research.

**Keywords:** cyanobacterial toxins; diversity; structural characterization; distribution; identification

**Key Contribution:** The existing information on the diversity of cyanotoxins was reviewed and the routine and advanced detection methods of cyanotoxins were also discussed.

## 1. Introduction

Cyanobacteria, which have existed for about 3.5 billion years, are believed to be the oldest creatures on Earth. Even though over 2600 cyanobacterial species have been described so far, it is believed that there are still many unknown species [1]. Cyanotoxins are toxic secondary metabolites produced by cyanobacteria, and pose a major threat to the ecological environment and human health. Cyanotoxins are diverse in functional properties and chemical structures. Based on functional properties, cyanotoxins can be mainly classified into hepatotoxins, neurotoxins, dermatotoxins and cytotoxins [2]. According to the chemical structure, cyanotoxins are mainly divided into cyclic peptides, alkaloids, lipopeptides,

non-protein amino acids and lipoglycans. Different chemical structures of cyanotoxins contribute to the diversity of their functional properties. A number of lipopeptide cyanotoxins have been found to have peculiar biological activities [3,4], such as anticancer, antifungal and molluscicidal activities. However, there are still many cyanotoxins that are believed to have special activities that have not yet been identified.

Cyanobacterial toxins are widely distributed in fresh and saltwater, and they have been reported hundreds of times in various countries and regions of the world, even in the Arctic and arid desert areas [5,6]. The extensiveness of the distribution of cyanotoxins is self-evident, however, the characteristics of the distribution and the severity of cyanobacterial toxins vary in different areas. For example, in the Baltic region, cyanobacterial blooms covering an area of 100,000 square kilometers can be formed in the summer [7], and various cyanotoxins produced by this cyanobacteria are seriously threatening the surrounding countries.


The harm of cyanotoxins is obvious, and it has been regarded as a new public health concern by the World Health Organization (WHO) [5]. Common methods such as bioassays, biochemical analysis, chemical assays and molecular assays have been used to detect cyanotoxins. However, because of the diversity of cyanotoxin species and chemical structure, the current routine methods cannot be used to detect all the types and variants of cyanotoxins. Therefore, the development of valuable techniques and methods of detecting cyanotoxins are prerequisite for the accurate identification of cyanobacterial toxins in the environment.

For these reasons, data on the structural characterization and regional distribution of cyanotoxins over the past twenty years were collected for this paper to provide a review of the cyanotoxin diversity in the world. We also summarize and discuss the protocols and advanced detection methods for cyanotoxins reported in recent years.

## 2. Structural Characterization

### 2.1. Cyclic Peptides

A cyclic peptide is a cyclic polypeptide chain formed by a covalent linkage between the amino termini and the carboxyl termini, the amino termini and the side chain, the carboxyl termini and the side chain, or the side chain with another side chain. Microcystins (MCs) and nodularins (NODs) are two classical cyclic peptide cyanobacterial toxins (Figure 1).



**Figure 1.** Structure of cyclic peptide cyanotoxins. **(A)** Microcystin-LR, YR and RR, **(B)** Nodularin.

### 2.1.1. Microcystins

MCs are hepatotoxic cyclic heptapeptides. The general structure of a MC is cyclo (-D-Ala-L-X-D-*erythro*-β-methylAsp-L-Y-Adda-D-Glu-N-methyldehydro-Ala), in which Adda is the unusual C20 amino acid (3-amino-9-methoxy-2,6,8-trimethyl-10-phenyl-4,6-dienoic acid) [8]. Nuclear magnetic resonance (NMR) was used for the structure determination of MCs in the 1990s [9]. In recent years, more sensitive mass spectrometry (MS) techniques have enabled the detection of MCs in trace amounts in cyanobacterial samples. To date, 246 MC congeners have been found [10]. The most common congeners are MC-LR, MC-YR and MC-RR, resulting from the presence of the L-forms of leucine (L), tyrosine (Y) or arginine (R) in positions 2 and 4 (Figure 1A). The structure diversity of MCs results in significant differences in toxicity, and the toxicity of MCs has been demonstrated in 79 countries [11]. As one of the most frequent and toxic variant, MC-LR had been classified as a possible human carcinogen (group 2B) by the International Agency for Research on Cancer (IARC) in 2010 [12], and the WHO has established a provisional guideline of 1 µg/L MCs for human drinking water.

Investigations have found that the distribution of MCs in freshwater habitats is a matter of concern, and MCs are typically produced by several cyanobacterial genera including *Anabaena*, *Dolichospermum*, *Geitlerinema*, *Leptolyngbya*, *Microcystis*, *Nostoc*, *Phormidium* and *Planktothrix* [13]. In addition, MCs have also been found in many marine cyanobacteria [14] such as *Spirulina*, *Synechococcus* and *Trichodesmium* (Table 1).

Biochemical methods, including enzyme-linked immunosorbent assays (ELISA) and phosphatase inhibition assay (PPIA), are well suited for the quick screening of MCs in water samples [15]. However, physicochemical methods are more sensitive and accurate than biochemical analysis and can provide quantitative analysis. For example, ultraviolet (UV) absorbance detection method can detect sub-nanogram amounts of toxin, and major congeners of MCs are easily identified and quantified by reversed-phase high-performance liquid chromatography (HPLC) [16]. In recent years, ultra-HPLC (UHPLC) tandem MS becomes the preferred technique for quantitative analysis and detection of cyanotoxins in water bodies, cyanobacteria and shellfish, due to its high efficiency and high-sensitivity [17].

### 2.1.2. Nodularins

NODs are hepatotoxic cyclic pentapeptides. The structure of NODs are similar to MCs, consisting of D-glutamic acid, N-methyldehydrobutyryne, D-*erythro*-β-methylaspartic acid and L-arginine (Figure 1B) [11]. Since the first description of an NOD in 1988, ten congeners have been characterized [10], and NOD-R is the most often found. NODs are considered to be liver tumor initiators and promoters [18]. According to the IARC evaluation, NODs are not classifiable to their carcinogenicity to humans, because of the lack of exposure data [12]. However, NODs are potent cyanotoxins that have been shown to be associated with death in animals and humans, and appear to be more toxic than MCs.

NODs are widely distributed throughout the temperate and subtropical regions, and are mainly found in the coastal sea water and freshwater. In the past, NODs produced by cyanobacteria had only been found in *Nodularia* [11], but the latest reports found that NODs were also isolated from the newly identified cyanobacteria *Iningainema* (*Scytonemataceae*) in Australia [19] (Table 1).

NODs and MCs are similar in structure, so it is expected that NODs have similar molecular mechanisms of toxicity to MCs [8]. Therefore, the common methods for identifying and detecting MCs are equally applicable to NODs.

**Table 1.** Category, common name, primary toxicity, analogues and producing genera of cyanotoxins.


Category	Common Name	Primary Toxicity	Analogues	Producing Genera	Reference
Cyclic peptides	Microcystins	Hepatotoxicity	246	<i>Anabaena, Aphanizomenon, Dolichospermum, Fischerella, Geitlerinema, Hapalosiphon, Leptolyngbya, Limnothrix, Merismopedia, Microcystis, Nostoc, Oscillatoria, Phormidium, Planktothrix, Pseudanabaena, Radiocystis, Pseudoanabaena, Spirulina, Synechococcus, Trichodesmium, Woronichinia</i>	[6,11,13,14,20–22]
	Nodularins	Hepatotoxicity	10	<i>Iningainema, Nodularia</i>	[10,11,19]
	Cylindrospermopsins	Hepatotoxicity	5	<i>Anabaena, Aphanizomenon, Chrysosporum, Cylindrospermopsis, Dolichospermum, Lyngbya, Oscillatoria, Raphidiopsis, Sphaerospermopsis</i>	[13,23–25]
Alkaloids	Saxitoxins	Neurotoxicity	57	<i>AnabaenaAphanizomenon, Cuspidothrix, Cylindrospermopsis,Dolichospermum, Fischerella, Geitlerinema, Lyngbya, Phormidium, Planktothrix, Raphidiopsis, Scytonema, Tolypothrix</i>	[11,13,21,26–28]
	Anatoxin-a	Neurotoxicity	4	<i>Anabaena, Aphanizomenon, Arthrospira, Cuspidothrix, Cylindrospermum, Dolichospermum, Microcoleus, Microcystis, Oscillatoria, Phormidium, Planktothrix, Pseudanabaena, Raphidiopsis, Tychonema</i>	[13,29,30]
Lipopeptides	Anatoxin-a(s)	Neurotoxicity	-	<i>Anabaena</i>	[22,23]
	Lyngbyatoxins	Dermatoxicity	7	<i>Oscillatoria, Schizothrix, Lyngbya majuscula</i>	[31–35]
	Aplysiatoxins	Dermatoxicity	5	<i>Lyngbya majuscula</i>	[34,36]
	Antillatoxins	Neurotoxicity	2		[37,38]
	Jamaicamides	Neurotoxicity	6		[39,40]
	Kalkitoxins	Neurotoxicity	7		[41,42]
	Barbamides	Molluscicidal	2	<i>Lyngbya majuscula</i>	[43]
	Majusculamides	Cytotoxicity	4		[31,34,44,45]
	Hectochlorins	Cytotoxicity	5		[46,47]
	Curacins	Cytotoxicity	5		[48–51]
Nonprotein amino acid	β-N-methylamino-l-alanine	Neurotoxicity	4	<i>Anabaena, Calothrix, Cyanodictyon, Leptolyngbya, Limnothrix, Microcoleus, Microcystis, Myxosarcina, Nodularia, Nostoc, Oscillatoria, Phormidium, Planktothrix Pseudanabaena, Scytonema, Synechococcus, Trichodesmium</i>	[52–56]
Lipoglycans	Lipopolysaccharides	Endotoxicity	-	All genera of cyanobacteria	[57]

## 2.2. Alkaloids

Alkaloids are basic organic compounds containing nitrogen. They vary in type and structure, so the properties of different alkaloids can greatly vary. Most alkaloids have a complex cyclic structure. Six alkaloid cyanotoxins are introduced here, including cylindrospermopsins (CYNs), saxitoxins (STXs), anatoxin-a (ATX-a), anatoxin-a(s) (ATX-a(s)), lyngbyatoxins and aplsiatoxins.

### 2.2.1. Cylindrospermopsins

A CYN is an alkaloid hepatotoxin that is formed from a tricyclic guanidine moiety combined with hydroxymethyluracil (Figure 2A) [58]. The chemical structure of a CYN was first identified in 1992 [59]. Currently, there are five known CYN analogs, namely CYN, 7-Deoxydesulfo-CYN, 7-deoxydesulfo-12-acetyl-CYN, 7-*epi*-CYN, 7-deoxy-CYN [25]. CYNs primarily target the liver, but it is also a cytotoxin. Recent studies have found that CYNs also cause nervous system damage, and they are more toxic than MCs in terms of neurotoxicity [58,60].



**Figure 2.** Structure of alkaloid cyanotoxins. (A) Cylindrospermopsin, (B) Saxitoxin, (C) Anatoxin-a, (D) Anatoxin-a(s), (E) Lyngbyatoxin, (F) Aplysiatoxin.

CYNs were first reported in 1979 after a hepato-enteritis outbreak which was caused by a *Cylindrospermopsis* bloom in Palm Island (Queensland, Australia) [61]. CYNs are mainly found in fresh water and seawater in tropical, subtropical, and temperate climate zones. CYNs are produced by various cyanobacteria, including *Aphanizomenon*, *Chrysosporum*, *Cylindrospermopsis*, *Oscillatoria* and *Raphidiopsis* [23] (Table 1).

Reverse phase-HPLC coupled with photodiode array detection was the first screening method used for CYNs [62]. Unlike many other cyanotoxins, CYNs are often present in the form of extracellular dissolution [25]. There is little research on the detection of CYNs in drinking water and recreational water, and clearly, an effective method is crucial. Presently, the methods most widely used for the detection and identification of CYNs include bioassays, ELISA, liquid chromatography (LC), capillary electrophoresis (CE) and molecular methods [25,63]. CYNs from multiple types of samples can be quantified by ELISA with high sensitivity. However, it is nonselective for CYNs analogues and cross-reactivity may occur. LC-MS has been proven to be an ideal method for discovering trace CYNs in water samples because of its sensitivity and specificity, and it has been established as the standard method for CYNs detection [64].

## 2.2.2. Saxitoxins

STX, a neurotoxic alkaloid, is the most representative paralytic shellfish poison. The structure of STX, which was finally confirmed in 1971, is composed of a 3,4-perhydropurine tricyclic system with two guanidinium groups (Figure 2B) [65]. Currently, 57 STXs have been described [28]. STX is one of the most lethal toxins, and it has been used in chemical weapons [66]. STXs cause an annually estimated 2000 cases of paralytic shellfish poisoning globally, with a mortality rate of 15% [67]. In reference to previous poisonings, an estimated dose of 0.5–1 mg of STX is considered toxic for humans. The Food and Drug Administration recommends that the concentration of STXs in shellfish food should not exceed 80 µg/100 g [21].

STXs are produced by cyanobacterial and accumulate in shellfish, which can then be ingested by humans. *Anabaena*, *Aphanizomenon*, *Cuspidothrix*, *Cylindrospermopsis*, *Dolichospermum*, *Fischerella* and *Geitlerinema* are some of the potentially STX-producing genera (Table 1).

Common physicochemical methods, toxicology-based biological methods, and biochemical methods are suitable for the detection and quantification of STXs [68]. The previous “gold standard” for STX measurement was the mouse bioassay, which has been refined and standardized by the Association of Official Analytical Chemists to provide quick and adequately accurate measurements. Nowadays, cytotoxicity tests have become an alternative to the mouse bioassay as a routine monitoring method to detect STXs [69]. Immunoassays based on the interactions of an antibody and its target is also commonly used for the detection of STXs. Owing to cross-reactivity within the different derivatives, it is recommended that ELISAs be used as screening tools rather than quantitative assays [68]. In the last decade, LC-MS has been successfully adapted for the analysis of STXs, and hydrophilic interaction liquid chromatography (HILIC) is often used for separation and selective reaction monitoring [70]. If only sensitivity and selectivity are considered, HPLC-fluorescence (HPLC-FL) and HPLC-MS are still the best methods [71].

## 2.2.3. Anatoxin-a

Anatoxins (ATXs) are neurotoxins isolated from cyanobacteria *Anabaena*. ATX-a and ATX-a(s) are the two most important ATXs. ATX-a is a neurotoxic bicyclic alkaloid, containing 2-acetyl-9-azabicycle(4.2.1)non-2-ene (Figure 2C) [72]. ATX-a was the first cyanotoxin to be structurally resolved by X-ray crystallography and confirmed by <sup>1</sup>H-NMR [72]. The four congeners of this compound are ATX-a, homoATX-a, dihydroATX-a and dihydrohomoATX-a [30]. HomoATX-a is a potent analogue of ATX-a. First synthesized in 1992, the structure has a propionyl group in place of the acetyl group [73]. ATX-a and homoATX-a are nicotinic agonists and have been shown to bind to nicotinic acetylcholine receptors [52]. In sufficiently high concentrations, ATXs can paralyze the nervous system and cause death. However, these neurotoxins are unstable and have a short half-life [74]. They can be rapidly degraded into nontoxic products, especially in light and high pH conditions.

ATX-a was first isolated from the freshwater cyanobacteria *Anabaena flos-aquae* in the 1970s. To date, ATX-a has been reported to exist in many cyanobacterial genera, such as *Anabaena*, *Aphanizomenon*, *Arthrospira*, *Cuspidothrix*, *Cylindrospermum*, *Dolichospermum*, *Oscillatoria* and *Phormidium* (Table 1).

Apart from bioassays, most reported methods for detecting ATX-a are based on chemical assays. HPLC coupled with UV or fluorescence detection, gas chromatography/Mass Spectroscop (GC/MS), and HPLC-MS are becoming the most common techniques [75]. Among them, LC-MS/MS offers perhaps the best analytical approach to determining ATX-a, homoATX-a, their degradation products and analogues [74]. Furthermore, qPCR allows the quantification of ATX-a gene (*anaC*) copy numbers in environmental samples. Recently, a new tested PCR-based method to detect ATX-a in aquatic ecosystems was developed, improving the detection level of *anaC* in environmental samples [76].

#### 2.2.4. Anatoxin-a(s)

ATX-a(s) is a unique guanidinium methyl phosphate ester that is also a neurotoxic alkaloid [77]. The structure of ATX-a(s) was determined by NMR and MS in 1989 [78], and its structure was determined to be (5S)-2-amino-1-((hydroxymethoxyphosphinyl) oxy)-N,N-dimethyl-4,5-dihydro-1*H*-imidazole-5-methanamine (Figure 2D) [22]. To date, no analogues of ATX-a(s) have been found. ATX-a(s) is also a natural organic phosphorus compound [79]. It is similar to organophosphorus and carbamate insecticides, and it may act as a potent irreversible acetylcholinesterase inhibitor to cause toxicity. ATX-a(s) is produced only by *Anabaena* and is not found as frequently as other cyanobacterial toxins [22] (Table 1).

Few studies have been done on the analysis of ATX-a(s). Bioassays have been used to detect ATX-a(s) in the environment, but such assays are limited by several issues, such as the sensitivity and specificity of detection [80]. Because of the lack of chromophores on this organophosphorus molecule, HPLC-UV is not suitable for its detection [81]. The AChE inhibition assay is useful for detecting ATX-a(s), but other organophosphorus compounds may inhibit the activity of this enzyme, so this method is not always dependable [82]. Due to these limitations, it is more effective to combine multiple methods than to rely on a single approach. To ensure reliability, the ATX-a(s) in the samples should be reconfirmed by LC-MS/MS.

#### 2.2.5. Lyngbyatoxins

Lyngbyatoxins are dermatotoxic alkaloids. Chemical and spectral data show that the lyngbyatoxin molecule is comprised of an indolactam ring, with an attached linalyl side group at C-7 (Figure 2E) [83]. So far, seven natural lyngbyatoxin analogues have been identified, namely lyngbyatoxin A-C, 12-*epi*-lyngbyatoxin A, 2-oxo-3(*R*)-hydroxy-lyngbyatoxin A, 2-oxo-3(*R*)-hydroxy-13-N-desmethyllyngbyatoxin A and 2,3-seco-2,3-dioxolyngbyatoxin A [32,34,35]. Lyngbyatoxins are highly inflammatory and vesicatory substances derived from marine cyanobacteria [84]. In another study, lyngbyatoxins were reported as a causative agent of human skin irritation and marine turtle poisoning [83,85]. Lyngbyatoxins are also considered to be an effective tumor promoter that induces protein kinase C (PKC) activity [86].

Lyngbyatoxins were first isolated from the lipid extract of a Hawaiian shallow-water variety of *Lyngbya majuscula* in 1979 [83]. Lyngbyatoxins have thus far been detected in the blooms of *Lyngbya*, *Oscillatoria* and *Schizothrix* [32,33]. A recent study found that it was also present in the Campania coast of southern Italy [87] (Table 1).

Chemical analysis methods are often used to detect lyngbyatoxins in environmental samples, such as the liquid chromatogram-high resolution mass spectrum (LC-HRMS/MS) combined with bioinformatics analysis on molecular network [87]. Also, bioanalysis is an important method for the toxicity evaluation of lyngbyatoxins.

#### 2.2.6. Aplysiatoxins

Aplysiatoxins are considered to be dermatotoxic alkaloids [77]. The molecular architecture of aplysiatoxins are bislactones of 3,4-dihydroxyvaleric acid and 4,6,6,10,12-pentamethyl-3,7,9,11,15-tetraoxy-15-phenylpentadecanoic acid (Figure 2F) [88]. Five analogues of this compound have been identified by NMR and MS, namely aplysiatoxin, debromo-aplysiatoxin, anhydro-aplysiatoxin,


3-methoxydebromo-aplysiatoxin, oscillatoxin A and 31-noroscillatoxin B [36]. Similar to lyngbyatoxins, aplysiatoxins induce contact dermatitis through the activation of PKC [23]. Aplysiatoxin and debromo-aplysiatoxin were also recently found to be potent tumor promoters in two-stage carcinogenesis in mouse skin [89]. Another study has showed that aplysiatoxin and its debrominated analogue induced expression of latent HIV-1 provirus in both cell line and primary cell models [90]. Aplysiatoxins are produced by the marine cyanobacteria *Lyngbya majuscula* in estuarine and coastal waters under tropical and subtropical climates [34] (Table 1).

Chemical analysis and bioanalysis are suitable for detecting aplysiatoxins in the environment and evaluating its toxic effects, respectively. It has been shown that LC-MS/MS-based molecular networking approach is a rapid analytical technique, which can be used to detect aplysiatoxins from environmental samples [36].

### 2.3. Lipopeptides

Lipopeptides are a class of compounds consisting of aliphatic chains and peptide chains, which have a wide range of biological activities. Most lipopeptides are derived from metabolites of microorganisms (including cyanobacteria).

Seven lipopeptide cyanotoxins from marine cyanobacteria are described as antillatoxins, jamaicamides, kalkitoxins, barbamides, majuscumamides, hectochlorin and curacins, respectively (Figure 3).



**Figure 3.** Structure of lipopeptide cyanotoxins. (A) Antillatoxin, (B) Jamaicamide A-C, (C) Kalkitoxin, (D) Barbamide, (E) Majuscumamides A and B, (F) Hectochlorin, (G) Curacin A.

### 2.3.1. Antillatoxins

Antillatoxin A is a neurotoxic lipopeptide that was isolated from the tropical marine cyanobacteria *Lyngbya majuscula* in Curaçao [38]. Antillatoxin B is a *N*-methylhomophenylalanine analogue also produced by *Lyngbya majuscula* that has been found in Puerto Rico and Florida [37]. The original planar structure of antillatoxins was determined by standard spectroscopic techniques. Antillatoxins are composed of a tripeptide that forms both ester and amide linkages with a highly methylated lipid section (Figure 3A), and their structural characteristics is a 9-tert-butyl-6,8-dimethyl-6,8-diene unit attached to the C5 of the cyclic peptide backbone [38,91]. Antillatoxins exhibit significant sodium channel activating and neurotoxicity that activates voltage-gated sodium channels leading to sodium influx in cerebellar granule neurons and cerebrocortical neurons [37,92]. Antillatoxins also have potent cytotoxicity toward Neuro 2a mouse neuroblastoma cells [93].

### 2.3.2. Jamaicamides

Jamaicamides, a kind of neurotoxic lipopeptide, contain an alkynyl bromide, vinyl chloride,  $\beta$ -methoxy enone system and pyrrolinone ring (Figure 3B). In 2004, three jamaicamides were isolated from the marine cyanobacteria *Lyngbya majuscula*, and were named as jamaicamide A-C [40]. The only difference in their structures was at the terminal of their polyketide aliphatic chains [94]. In 2015, three new analogues (Jamaicamide D-F) were discovered in the crude extract of *Lyngbya majuscula* by LC-MS and molecular networking analysis [39]. These jamaicamides have unusual biological activities, including sodium channel blocking, and arthropod and fish toxicity [23,40]. As a sodium channel blocker in neocortical neurons, jamaicamide A is approximately two to three times more potent than jamaicamide B and F [39].

### 2.3.3. Kalkitoxins

The neurotoxic lipopeptide kalkitoxin was first isolated from the marine cyanobacteria *Lyngbya majuscula* in the Caribbean Sea [42]. Natural kalkitoxins possess a 2,4-disubstituted thiazoline, a lipophilic chain and an unsaturated  $\text{CH}_2=\text{CH}_2$  unit (Figure 3C) [95]. Based on the evidence by 1D and 2D NMR spectra, six kalkitoxin isomers (3-*epi*-kalkitoxin, 7-*epi*-kalkitoxin, 8-*epi*-kalkitoxin, 10-*epi*-kalkitoxin, 10-*nor*-kalkitoxin and 16-*nor*-kalkitoxin) have been synthesized [41]. Kalkitoxins are strongly ichthyotoxic, and toxic to brine shrimp. They inhibit cell division, suppress inflammation and potently block voltage-sensitive sodium channels in murine neuro-2a cells [42]. In humans, Kalkitoxins have a neurotoxicity that acts on sodium channels and mitochondria [96].

### 2.3.4. Barbamides

Barbamide is a recently discovered lipopeptide with molluscicidal activity and it is isolated from the marine cyanobacteria *Lyngbya majuscula* [97]. The structure of barbamide was determined by spectroscopic methods, and the several unique structural features include a trichloromethyl group and the methyl enol ether of a  $\beta$ -keto amide (Figure 3D) [98]. Heterologous expression of the barbamide biosynthetic gene cluster resulted in the production of a new barbamide analogue named as 4-O-demethylbarbamide [43]. This new compound is several-fold more potent than barbamide as a molluscicide. If it lacks other toxicities, it may be a superb candidate for treating snail-infested waterways which pose health risks for human populations [43].

### 2.3.5. Majusculamides

There are four Majusculamides (majusculamides A-D) that have been found. Majusculamides A and B are cytotoxic lipopeptides isolated from the marine cyanobacteria *Lyngbya majuscula* with antilarval settlement activities [31,34]. Their structures were elucidated using high-resolution fast atom bombardment mass spectrometry (HR-FAB-MS), and 1D and 2D NMR analyses [34]. Majusculamide A was identified as *N*-((2*R*)-2-methyl-3-oxodecanoyl)-*D*-N,O-dimethyltyrosyl-L-

N-methylvalinamide [45]. Majusculamide B is an epimer of majusculamide A (Figure 3E). Majusculamide C was previously found in cyanobacteria from the Marshall Islands and was reported to have antifungal activity [45]. Majusculamide D was also originally isolated from *Lyngbya majuscula*. The absolute configuration of the 1,3-dimethyloctanamide motif was determined by the synthesis of this fragment via zirconium-catalyzed asymmetric carboalumination chemistry [44]. Majusculamides possess potent cancer cell toxicity, exhibiting selective cytotoxicity toward both pancreatic (PANC-1) and glioblastoma (U251N) cancer cell lines [44].

### 2.3.6. Hectochlorins

Hectochlorin is a cytotoxic lipopeptide isolated from marine cyanobacteria *Lyngbya majuscula* in Jamaica and Panama [99]. The planar structure of this compound was determined by 1D and 2D spectroscopy, and X-ray crystallography was used to determine the absolute stereochemistry. Hectochlorin resembles lyngbyatoxin in structure (Figure 3F) [47]. In 2015, three hectochlorin analogs were identified from mass spectroscopy based-molecular networking, respectively (Hectochlorin B-D) [39]. Hectochlorin has the ability to promote actin polymerization and shows cytotoxicity and potent inhibitory activity toward the fungus *Candida albican* [47,99]. Hectochlorin has also shown great potency against several cancer cell lines, namely colon, melanoma, ovary, and kidney [100]. Deacetylhectochlorin, a derivative of hectochlorin, was isolated from the Thai sea hare (*Bursatella leachii*). This cognate is also a potent stimulator of actin assembly, and exhibits more potent toxicity than hectochlorin against several human carcinoma cell lines [47]. The artificial synthesis of hectochlorin is the focus of current researches [99,101].

### 2.3.7. Curacins


Curacin A is a marine cytotoxic lipopeptide found in strains of the tropical marine cyanobacteria *Lyngbya majuscula*, in Curacao [48]. NMR analysis showed that its structure contains one cysteine, ten acetate units and two S-adenosyl methionine-derived methyl groups (Figure 3G). There are four isomers of curacin A with similar biological activity and structure, namely (curacin B-E). Curacins B and C were isolated from a Curagao collection of *Lyngbya majuscula* [49]. Curacin D was found in *Caldora penicillata* and *Lyngbya majuscula* from the Pacific islands and the Virgin Islands. The structural elucidation of curacin D was accomplished through multidimensional NMR and GC/MS [50,51]. Curacin E was isolated from *Ophiocoma scolopendrina* at Kabira Reef on Ishigaki Island. NMR data and the GC spectrum showed that curacin E has an ethylcarbonyl terminus in its side chain [102]. Curacins exert their potent cell toxicity through interaction with the colchicine drug binding site on microtubules [103,104], and it is also a potent cancer cell toxin and antimitotic agent of cyanobacteria origin [105].

## 2.4. Nonprotein Amino Acids

Nonprotein amino acids are natural compounds containing amino and carboxyl groups which do not constitute proteins. They are important bioactive substances mainly in microorganisms and plants.  $\beta$ -N-methylamino-L-alanine (BMAA, Figure 4) produced by cyanobacteria, is a type of nonprotein amino acids that was found to be a cyanotoxin. BMAA is composed of carbon with a carboxyl group, an amino group and a methylamino side chain [11,106]. Three biologically relevant structural isomers of BMAA have been identified [53], including 2,4-diaminobutyric acid (2,4-DAB), *N*-2-aminoethylglycine (AEG) and  $\beta$ -amino-N-methyl lanine (BAMA).

Research on BMAA has intensively increased over the past decade, and now BMAA appears to be a cause of the Guamanian ALS and Parkinsonism Dementia Complex (PDC) [107]. BMAA was first isolated from a cycad on the island of Guam in 1967. Symbiotic cyanobacteria produces BMAA in the coraloid root of the cycad, and then BMAA is transferred to and accumulated in the seeds. BMAA was once believed to occur only in cycads. However, it is now known to be produced by a range of cyanobacteria, diatoms and dinoflagellates across the world in freshwater, saltwater and terrestrial

ecosystems [108,109]. BMAA is typically detected in strain cultures of cyanobacterial species belonging to *Anabaena*, *Calothrix*, *Microcystis*, *Nostoc*, *Scytonema*, *Synechococcus* and *Trichodesmium* [52] (Table 1).




**Figure 4.** Structure of  $\beta\text{-N-methylamino-L-alanine}$ .

The analysis of BMAA in native form is complicated, because of the physicochemical properties and the absence of a chromophore or fluorophore. An HPLC-FL method was developed and used for amino acid analysis in 2003 [110]. In recent years, more specific methods have been developed that have been used for BMAA analysis, such as amino acid analyzers, UPLC-UV, UPLC-MS and LC-MS/MS [111,112]. Amino acid analyzers are often used in hospital settings for analysis of amino acid contents in physiological fluids [113]. Presently, LC-MS/MS method is the most widely used analytical method to determine the content of BMAA sensitively and accurately [111]. CE-MS/MS has been developed as an alternative method for the quantitative determination of free BMAA [114]. This method displays excellent resolution of amino acid isomers and shows no interference from matrix components.

## 2.5. Lipoglycans

Lipopolsaccharides (LPS), also known as lipoglycans and endotoxin, are large molecules consisting of a lipid and a polysaccharide. LPS are major components of cyanobacterial cell walls and cover 75% of the surface area of the outer membrane [57]. In general, the fatty acid component (lipid-A) of LPS is responsible for its toxic actions. It is an irritant and can trigger allergenic responses in human and animal tissues [115].

The structure of cyanobacterial LPS was determined by NMR, MS, GC and matrix-assisted laser desorption/ionization (MALDI)-MS [116]. The structure of LPS consist of an internal acylated glycolipid (lipid-A), a core domain (an oligosaccharide), and an outer polysaccharide chain (O-antigen) (Figure 5).



**Figure 5.** Structure of Lipid-A. Reproduced from [115]. Copyright 2005, Elsevier.

The toxicity of cyanobacterial LPS have been found to be less toxic than those from heterotrophic bacteria. It has been suggested that cyanobacterial LPS may cause or contribute to human illness, particularly in causing epidermal allergic reactions [116]. However, there is insufficient evidence in the scientific literature to confirm that skin contact with cyanobacterial LPS causes a skin rash. It is worth noting that cyanobacterial LPS are indirectly involved in various human diseases, including skin diseases, gastrointestinal issues, respiratory diseases, fever, allergies and headaches [117]. LPS have been shown to modulate biotransformation and toxicity of cyanotoxins and may worsen the hepatic damage induced by hepatotoxins [118].

LPS are believed to be produced by all cyanobacteria and gram-negative bacteria [57]. The production of this toxin has been reported in several cyanobacteria such as *Agmenellum*, *Anabaena*, *Anacystis*, *Microcystis*, *Oscillatoria*, *Phormidium*, *Schizothrix* and *Spirulina* [116].

Various LPS have a similar mode of biological action (toxicity) but have diverse chemical structures in different cyanobacterial strains [119]. Therefore, traditional methods such as chromatography are often not used in the analysis of LPS. Isolated LPS have been characterized and analyzed by other approaches, such as the in vivo toxicity test, the standardized pyrogenicity test with Limulus amoebocyte lysate, the advanced pyrogenicity test using the PyroGene rFC endotoxin system, or by testing the potency to activate human leukocytes [120,121]. In a recent study, a porous silicon membrane (pSiM)-based electrochemical biosensor was developed for a direct and sensitive detection of LPS [122].

### 3. Distribution of Cyanotoxins

The global distribution of cyanobacteria toxins is presented in Figure 6. Common cyanotoxins (MCs, NODs, CYNs, STXs, ATXs and BMAA) are found in every continent except Antarctica (no BMAA and ATXs have been found in Antarctica). However, cyanotoxins are more frequently reported in Europe and North America. There are fewer reports from Asia, South America and Africa. Marine cyanotoxins are mainly distributed in the western Atlantic and central Pacific oceans.

#### 3.1. Asia

Six common cyanobacterial toxins (MCs, CYNs, NODs, STXs, ATXs and BMAA) have been found in Asia (Table 2). These cyanotoxins were mainly distributed in temperate and tropical coastal areas of Asia, as well as inland lakes and rivers in various countries. Among them, MCs are the most widely distributed, which have been reported in China [123–126], Japan [127], Korea [128], Bangladesh [129], Singapore [130], Saudi Arabia [131], India [132], Philippines [133], Thailand [134], Vietnam [135,136], Israel [137] and Turkey [138]. CYNs and their homologues were mainly distributed in China [125,139], Japan [140], Saudi Arabia [141], Vietnam [142], Thailand [134,143], Israel [144] and Turkey [145]. NODs were reported only in China [146] and Turkey [138]. STXs were reported in China [147], Korea [148], India [149], Singapore [150], Bangladesh [151] and Turkey [152]. ATXs have been found in Korea [153], Qatar [154], India [149], China, Japan and Turkey [138]. And BMAA was reported in China [155], Japan [156] and Qatar [111,157].

#### 3.2. Africa

There are six common cyanobacterial toxins found in Africa: MCs, NODs, CYNs, STXs, BMAA and ATXs (Table 2). Various cyanobacterial toxins have been reported in inland rivers, and in lakes in Eastern Africa, as well as coastal areas in the north. South Africa and Nigeria have also been threatened by cyanobacterial toxins. MCs were reported in Ethiopia [158], Algeria [159], Tanzania [160], Kenya [161], South Africa [162], Tunisia [163], Zimbabwe [164], Morocco [165], Uganda [166], Egypt [167], Nigeria [168] and Ghana [169]. CYNs were found in Egypt [170] and Nigeria [171]. NODs were reported in South Africa [172] and Nigeria [171]. Besides, MCs, NODs and CYNs were also found in several riverside countries along the Limpopo River [173]. STXs were reported in Morocco [174] and Nigeria [171]. ATXs were distributed in Kenya [175] and Nigeria [171]. And BMAA was only reported in South Africa [176].

### 3.3. North America


Six common cyanobacterial toxins were distributed in North America: MCs, NODs, CYNs, STXs, BMAA and ATXs (Table 2). These cyanobacterial toxins were widely distributed in inland freshwater lakes and coastal areas in the US [177–181] and Canada [182–184]. MCs [185], CYNs [186] and STXs [187] have been reported in Mexico. Also, MCs and NODs have been found in Greenland and Alaska in the arctic region [188,189].

### 3.4. South America

Although South America has had fewer reports of cyanobacterial toxins compared with other continents, five common types have been reported: MCs, CYNs, NODs, STXs, and ATXs (Table 2). In this region, Brazil is the most affected country, and all five of these toxins have been reported to be found in freshwater [190–193]. In addition, MCs were reported in Argentina [194], Chile [195] and Uruguay [196]. CYNs were also found in Argentina [194] and Uruguay [196]. Presently, BMAA has only been reported in Peru [197].

### 3.5. Antarctica

Four common cyanobacterial toxins have been found in Antarctica (Table 2). The cold climate in this area of the world is not suitable for animal survival. However, in recent years, MCs [198,199], NODs [6], CYNs [200] and STXs [201] have been described in the Antarctic. This illustrates the strong adaptability and wide distribution of cyanobacterial toxins. However, ATXs and BMAA have not been reported in Antarctica.



**Figure 6.** Distribution of cyanotoxins.

**Table 2.** The diversity of cyanotoxins in various countries.

Regions	Countries	MCs	CYNs,	NODs	STXs	ATXs	BMAA
Asia (13 countries)	China	✓	✓	✓	✓	✓	✓
	Turkey	✓	✓	✓	✓	✓	
	Japan	✓	✓			✓	✓
	India and Korea	✓			✓	✓	
	Qatar	✓				✓	✓
	Bangladesh and Singapore	✓			✓		
	Israel, Saudi Arabia,						
	Thailand and Vietnam	✓	✓				
	Philippines	✓					
Africa (13 countries)	Nigeria	✓	✓	✓	✓	✓	
	South Africa	✓		✓			✓
	Egypt	✓	✓				
	Kenya	✓				✓	
	Morocco	✓			✓		
	Algeria, Ethiopia, Ghana, Namibia, Tanzania, Tunisia, Uganda and Zimbabwe	✓					
North America (3 countries)	Canada and the US	✓	✓	✓	✓	✓	✓
	Mexico	✓	✓		✓		
South America (5 countries)	Brazil	✓	✓	✓	✓	✓	
	Argentina and Uruguay	✓	✓				
	Chile	✓					
	Peru						✓
Antarctica	-	✓	✓	✓	✓		
Europe (24 countries)	France	✓	✓	✓	✓	✓	✓
	Finland, Italy, Poland and Portugal	✓	✓		✓	✓	✓
	Germany and Spain	✓	✓	✓	✓	✓	
	Sweden	✓	✓			✓	✓
	the UK	✓			✓	✓	✓
	Bulgaria	✓			✓	✓	
	Czech, Greece and Serbia	✓	✓		✓		
	Ireland	✓	✓			✓	
	Denmark				✓	✓	
	Hungary	✓	✓				
	Latvia	✓			✓		
	Netherlands	✓					
	Norway and Russia	✓				✓	
	Austria, Romania and Switzerland	✓					
	Lebanon		✓				
Oceania (2 countries)	Australia and New Zealand	✓	✓	✓	✓	✓	✓

Abbreviations: MCs: microcystins, CYNs: cylindrospermopsins, NODs: nodularins, STXs: saxitoxins, ATXs: anatoxins, BMAA:  $\beta$ -N-methylamino-l-alanine. “✓” represents existence.

### 3.6. Europe

Europe is the continent with the most reports of cyanobacterial toxins, which have been found in more than 20 countries and regions (Table 2). In particular, the Baltic Sea region is highly affected by cyanobacterial toxins, and cyanobacteria form periodic blooms in the summer, covering an area of over 100,000 square kilometers [202]. A variety of cyanobacterial toxins are produced in water, posing a major threat to the survival environment of humans and animals. Six common cyanobacterial toxins were distributed in Europe. MCs were reported in Finland [203,204], Austria [205], Greece [206], Poland [207], Ireland [208], Portugal [209], Hungary [210], Norway [211], Netherlands [212], Czech [213], Germany [214], Sweden [215], Switzerland [205], Spain [216], Poland [217], the UK [218], Bulgaria [219], Romania [220], France [221,222], Serbia [223], Russia [224–226] and Latvia [227]. CYNs were distributed in Ireland [228], Germany [229], France [230], Italy [231], Czech [232], Spain [233], Lebanon [234], Greece [235], Finland [236], Poland [237], Sweden [215], Serbia [238], Portugal [239] and Hungary [240]. ATXs were found in Ireland [241], Italy [242], France [243], Netherlands [244],

Denmark [245], Bulgaria [219], Finland [204], Poland [246], Sweden [215], Portugal [220], Spain [247], the UK [248] and Germany [249]. STXs were reported in Portugal [250], Denmark [251], France [252], Germany [253], Czech [213], Italy [254], Finland [255], the UK [256], Poland [257], Bulgaria [219], Serbia [223], Greece [235], Russia [27], Norway [258] and Spain [258]. NODs were found in Latvia [227], Germany [228], Spain [259] and France [260]. BMAA was reported in Finland [261], Sweden [262], Portugal [263], Italy [264], Poland [265], the UK [266] and France [267].

### 3.7. Oceania


Six common cyanobacterial toxins have been found in Australia (Table 2), which were detected in inland lakes and rivers of Oceania: MCs, NODs, CYNs, ATXs, STXs and BMAA [30,268,269]. The distribution of these toxins were mainly in the eastern and southern regions of Australia. These toxins have also been detected in the surrounding islands and oceans of Oceania, such as New Zealand [250,270–272].

### 3.8. The oceans

More than twelve cyanobacterial toxins have been found in the oceans across the world. MCs have been found in the marine environments of the central Atlantic coast of Portugal, the Canary Islands, the Brazilian coast, the Amvrakikos Gulf, and the Indian Ocean [14]. BMAA was reported from Guam in the western Pacific, and Bermuda in the western region of North Atlantic [52]. Aplysiatoxins have been reported in the Red Sea region [34], and the coast of Japan and Singapore [36,89]. Lyngbyatoxins were found in the coast of Hawaii [83] and Campania [87]. Seven lipopeptide cyanotoxins have been reported in the Caribbean and Central Pacific.

## 4. Identification of Cyanotoxins

Various analytical methods have been available to detect and identify cyanobacterial toxins, such as bioassays, biochemical assays, chemical assays and molecular analysis (Figure 7A–D).



**Figure 7.** Several representative analysis techniques of cyanotoxins. (A) Bioassays, (B) Biochemical assays, (C) Chemical assays, (D) Molecular assays.

### 4.1. Bioassays

Bioassays, involving cells, plants, or animals, are an important method for monitoring the presences of cyanotoxins in environmental samples, and evaluating their effects [80].

The mouse bioassay is a common method for identifying cyanotoxins and the toxin efficacy, which is evaluated by the LD<sub>50</sub> obtained from the survival time of infected animals. However, alternative chemistry and functional analysis are replacing animal trials due to ethical considerations, prolonged processes, high cost, low sensitivity and non-specificity [273]. Cytotoxicity tests have overcome the shortcomings of specificity and sensitivity in animal tests, and can provide qualitative and quantitative analysis of cyanotoxins. Immortalized cell lines and primary cells from animal tissues have been used to evaluate cyanotoxins and cyanobacterial extracts [274]. Primary cells can better reflect the physiological status in vivo than immortal cell lines. Plant tests have been used to determine the toxic effects of cyanotoxins [275]. Some plants have certain sensitivities to cyanotoxins and their physiological indicators may be affected by toxicity. The phytotoxic effects of MCs have been investigated by plant bioassays, and it was found that MCs can inhibit the growth and chlorophyll content of *Solanum tuberosum* culture [276]. Bioassays can provide unique information about toxic effects, which are clearly not achievable through physicochemical analysis methods. If the two methods are combined, a clear causal link can be established between the presence of a toxin and its toxic effects.

#### 4.2. Biochemical Assays

Biochemical assays are detection methods that rely on the interactions between cyanotoxins and biological macromolecules. Such examples would include immunoassays, enzyme inhibition assays, and receptor bioassays. Immunoassays such as the ELISA are used to identify specific toxins in samples through antigen-antibody binding. The PPIA is a method for determining cyanotoxins by measuring protein phosphatase activity. The Receptor Assay is a method based on competitive binding between a class of neurotoxins and a receptor.

##### 4.2.1. Immunoassays

In recent years, besides the ELISA, a significant amount of antibody-based immunoassays have been developed for the detection of cyanotoxins, including Fluorescence Immunoassay (FIA) [146], immunochromatography (ICA) [277,278] and biosensor techniques [279].

The ELISA is a conventional method for the analysis of cyanotoxins, and has been used to detect MCs, NODs, CYNs, ATXs and STXs in samples [273,280]. ELISA assays provide semi-quantitative estimates of toxin concentrations. It is easy to perform an ELISA, which does not require highly skilled personnel, and it is relatively inexpensive. However, the individual variants of the toxins have not been able to be identified by ELISA, and the quantification of toxins may not be as accurate as physicochemical methods [15].

FIA is a sensitive technique for the measurement of drugs, hormones, proteins and other compounds [281]. It is often used for the quantification of cyanotoxins. Fluorescence Polarization Immunoassay (FPIA) is an immunochemical method based on FIA which can precisely quantify a target and meet the requirements of a reliable and economical screening technology [282]. The principle of FPIA is based on the difference in fluorescence polarization of the labeled antigen and labeled analyte-antibody complex [283]. Over the past decade, FPIA has been widely used in high-throughput screening of various chemical contaminants. Compared with UPLC-MS and ELISA, FPIA can detect MCs in water more simply and efficiently. The method can be extended to detect other types of cyanotoxins in the environment. However, FPIA is susceptible to interference from light scattering and endogenous fluorophores in biological samples and also from the tracer binding to sample matrix components [283].

As a combination of chromatography and an immunochemical reaction, ICA is known as a lateral flow device or strip test. It has been widely developed for point-of-care testing because of its user-friendly formats, low cost and short assay times [284]. Lateral Flow Enzyme Immunochromatography assay (LFICA) coupled with molecular imprinting technique was developed for rapid detection of MC-LR in water products [277]. However, compared with other detection methods, the significantly lower analytical sensitivity hinders its further application. A way to enhance the performance of LFICA is to

eliminate the interferences from complicated matrixes. Using molecularly imprinted polymers (MIPs) as adsorbents for solid-phase extraction (SPE) columns, most of the interference in the matrix can be eliminated and sensitivity can be improved [285]. This approach achieves excellent sensitivity and specificity as well as a very low limit of detection, and it can be used to analyze other toxic components in foods and aquatic products by replacing specific MIPs and antibodies.

Biosensors are broadly used in environmental monitoring and have replaced the traditional methods of detecting toxins. Biosensors are analytical devices for the detection of an analyte, and they can provide selective quantitative or semi-quantitative analytical information. Biosensors which consist of a bio-recognition element (bioreceptor) and a transducer, can transform physiochemical information into a visual signal which can be captured by a detector [279]. The bioreceptor, can be a cell receptor, an enzyme, an organelle, etc., that binds an analyte under study. The bioreceptor is combined with the physiochemical detector or “transducer” that produces a physiochemical signal when the biosensor binds to the analyte [286]. Electrochemical and optical biosensors have been shown to detect aquatic toxins. Advanced portable biosensors would allow immediate assessment of water bodies and water treatment deficiencies [287]. Due to the simple and sensitive nature of biosensors, they can be used as an early warning monitoring tool for semi-quantitative screening of possible biotoxins in water samples.

#### 4.2.2. Enzyme Inhibition Assays

PPIA is a common enzyme inhibition assay used in the detection of cyanotoxins. Protein phosphatases are highly conserved in eukaryotes and are involved in many cellular processes [288]. A number of compounds are known to inhibit protein phosphatases, including MCs and NODs from cyanobacteria [288,289]. Cyanotoxins can be detected by following the degree of inhibition of protein phosphatases. The PPIA can not only identify a toxin, but also estimate its toxicity. PPIA has good sensitivity and does not seem to be affected by sample matrices, except for very high concentration of cellular extracts [290].

#### 4.2.3. Receptor Bioassays

The Receptor Bioassay has become the most recognized method for the detection and quantification of ATXs and STXs in shellfish and water [291]. The principle of this approach is based on the mechanism of certain toxins that compete with the nicotinic acetylcholine receptor. However, nicotinic acetylcholine receptors can also bind with spiroides, gymnodimines and other marine toxins [273,292]. Therefore, this method cannot distinguish the specific type of toxin. Solving this problem will involve the designing of specific and sensitive cholinesterase receptors.

### 4.3. Chemical Assays

Chemical analysis is the most reliable technique for detecting and identifying cyanobacterial toxins in samples based on the physical and chemical properties of cyantoxins. Some advanced chemical analysis methods have been used to detect and identify cyanobacterial toxins accurately and sensitively [293], such as NMR, HPLC, CE, LC-MS, GC, etc. Another technique is matrix-assisted laser desorption-ionization-time-of-flight mass spectrometry (MALDI-TOF-MS), which can distinguish toxic from non-toxic strains at the level of a single colony or filament, without prior solvent extraction [294,295]. Currently, MS-based methods are considered to be the best approach to unambiguously identify and quantify the different variants of cyanotoxins. However, they are not suitable for site testing because these analytical methods often require harmful solvents, expensive instruments, skilled personnel, and complex operational procedures. They are also time consuming and typically subjected to high cost [273,296].

#### 4.4. Molecular Assays

Molecular assays based on the polymerase chain reaction (PCR) have been developed over the past 20 years. The sequencing of cyanobacterial genomes has led to the identification of the gene clusters responsible for cyanotoxin production, which paved the way for the use of these genes as targets for PCR [297]. Three different PCR-based methods have been used for the detection of cyanotoxins [298–300]. The first approach is based on conventional PCR targeting only one gene. This method provides qualitative results that can be used to detect potential cyanobacterial toxins at the beginning of the bloom. In order to get a quantitative estimation of cyanotoxin gene abundance, the second approach was developed, which is based on the use of quantitative PCR (qPCR) targeting a gene involved in the biosynthesis of cyanotoxins. Finally, the combination of PCR and DNA microarray can be used to detect potentially toxic cyanobacterial species or to identify specific genes involved in the biosynthesis of cyanobacterial toxins [301]. However, it may be found that the actual toxin concentration in samples is inconsistent with qPCR results, which may be caused by various technical reasons and biological problems [286]. Therefore, the better choice to evaluate cyanotoxins was combining traditional analytical methods and PCR techniques.

Recently, microarray technologies have been used to achieve simultaneous detection and semi-quantification of cyanotoxins in environmental samples [302]. A detection method based on microsphere array coupled flow cytometry was developed [303]. The process can be used as a semi-quantitative screening tool for cyanobacterial toxins in fresh or brackish water. This method can save time and greatly reduce sample size due to the simultaneous detection of multiple cyanotoxins.

Fluorescence in Situ Hybridization (FISH), uses a fluorescein-labeled oligonucleotide probe to hybridize with nucleic acid target sequences in samples, to obtain specific information on gene expression. In FISH imaging, specific rRNA probes allow the identification of microorganisms at the genus level, while mRNA probes will give information on the expression of a particular gene [304]. Therefore, FISH imaging might be a very powerful technique for the detection and identification of toxic cyanobacteria, particularly those that produce MCs in the environment [304–306]. Some species in suboptimal growth conditions may have lower mRNA levels, in which case the sensitivity of the assay can be increased by Tyramine Signal Amplification (TSA-FISH) [307,308]. This supplement to the method involves a specific oligonucleotide that is linked to horseradish peroxidase (HRP), which catalyzes the permanent deposition of many fluorescent tyramides in the probe surroundings. This technique showed 10 to 20 fold signal amplifications relative to fluorescein-monolabeled probes [308–310]. For all these reasons, TSA amplification is highly recommended for the microscopic detection of planktonic algae.

These methods look very promising, but so far their application remains limited to the research level [298]. And the cost of these methods still hampers their use in monitoring applications.

#### 5. Final Remarks

The diversity of cyanobacteria leads to a variety of cyanobacterial toxins. According to the chemical structure, cyanobacterial toxins are mainly classified into cyclic peptides, alkaloids, lipopeptides, nonprotein amino acids and lipoglycans. The producing genera of cyanotoxins have been summarized in Table 1. MCs, CYNs, ATXs, STXs and BMAA can be produced by a variety of cyanobacteria. NODs are only produced by *Anabaena*. LPS exists in the cell walls of almost all cyanobacteria [57]. Lipopeptide cyanotoxins are mainly produced by the marine cyanobacteria *Lyngbya majuscula*.

We reviewed the distribution of cyanobacterial toxins around the world (Figure 6, Table 2). MCs were found to be the most widely distributed cyanotoxins (57 out of 60 countries), followed by CYNs (31 out of 60), STXs (29 out of 60), ATXs (26 out of 60), BMAA (16 out of 60) and NODs (13 out of 60). It was found that the diversity of cyanotoxins is mainly reflected in developed countries. The reports of cyanotoxins in developed areas are far more than that in underdeveloped areas. More than four cyanobacterial toxins were reported in 18 countries (China, the US, Canada, France, Australia, New Zealand, Turkey, Nigeria, Brazil, Finland, Germany, Italy, Sweden, Poland, Portugal,

Spain, Japan and the UK). Of these 18 countries, 13 are developed regions, accounting for 72%. Furthermore, there are 15 countries where only one cyanotoxin was reported and 14 of which are developing countries (except Switzerland). However, we are more inclined to believe that cyanotoxins may have a greater threat in less-developed areas due to the possible lack of monitoring activities in these areas.

There are many available methods to detect and identify cyanobacterial toxins, such as bioassays, biochemical assays, chemical assays and molecular analysis. At present, there is no single method that is the optimal for the detection and identification of all types of cyanobacterial toxins, and each method has its applicability [311]. Detection methods are affected by the variety and abundance of cyanotoxins. The choice of method is also inevitably influenced by the availability of the analytical equipment and its applicability in a particular environment. For example, molecular methods are more suitable for detecting potential toxic cyanobacteria, and immune sensors are better at monitoring the presence of toxins in samples and bioassays are often used to evaluate toxic effects. Thus, research purpose, economic feasibility, speed of analysis, sensitivity and field applicability should be taken into account when selecting detection methods. Previous experience shows that combining multiple methods can greatly improve the detection efficiency. It is still necessary to develop new techniques for detecting and identifying cyanobacterial toxins more easily and sensitively. Currently, plants have become an area of interest in many studies. It was found that plants can respond to other plants and animals in the environment [312]. Perhaps plants can be used as alternatives to animal experiments in the future after people have a deeper understanding of them.

Cyanobacterial toxins are widely distributed and varied, and their harmfulness cannot be ignored. Epidemiological studies and animal experiments have confirmed that many cyanotoxins have multi-organ toxicity and carcinogenic effects, posing a serious threat to human health and life quality [22,313]. To reduce the threat of cyanotoxins to humans, it is essential to strengthen the monitoring of cyanobacterial toxins worldwide, especially in underdeveloped areas. In recent years, cyanotoxins with special biological activities have attracted wide attention. Most of these cyanotoxins are produced by marine cyanobacteria. It was found that the lipopeptide cyanotoxins not only have neurotoxicity and cytotoxicity but also have anticancer, antifungal and molluscicidal activities [3]. These specific types of natural toxins have unique pharmacological properties and show great potential of being developed and utilized for human disease. Marine cyanobacteria are still a rich source of untapped natural products. However, the exploration of marine cyanotoxins is just beginning.

**Author Contributions:** Conceptualization, H.Z., H.G. and X.D.; Methodology, H.L. and Y.W.; Validation, Y.M. and R.W.; Formal Analysis, X.D. and X.C.; Investigation, L.Y. and H.L.; Data Curation, Y.W. and R.W.; Writing-Original Draft Preparation, H.Z., H.G. and X.D.; Writing-Review & Editing, H.G., M.D.L., X.D., H.Z. and X.C.; Supervision, H.Z. and H.G.; Funding Acquisition, H.Z. All authors have read and approved the final manuscript.

**Funding:** This research was funded by The National Nature Science Foundation of China (grant number 81773384 and 81472948).

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Tokodi, N.; Drobac, D.; Lazić, G.; Petrović, T.; Marinović, Z.; Lujić, J.; Malešević, T.P.; Meriluoto, J.; Svirčev, Z. Screening of cyanobacterial cultures originating from different environments for cyanotoxicity and cyanotoxins. *Toxicon* **2018**, *154*, 1–6. [[CrossRef](#)]
2. Corbel, S.; Mougin, C.; Bouaicha, N. Cyanobacterial toxins: Modes of actions, fate in aquatic and soil ecosystems, phytotoxicity and bioaccumulation in agricultural crops. *Chemosphere* **2014**, *96*, 1–15. [[CrossRef](#)]
3. Giordano, D.; Costantini, M.; Coppola, D.; Lauritano, C.; Nunez Pons, L.; Ruocco, N.; di Prisco, G.; Ianora, A.; Verde, C. Biotechnological applications of bioactive peptides from marine sources. *Adv. Microb. Physiol.* **2018**, *73*, 171–220. [[CrossRef](#)] [[PubMed](#)]
4. Demay, J.; Bernard, C.; Reinhardt, A.; Marie, B. Natural products from cyanobacteria: Focus on beneficial activities. *Mar. Drugs* **2019**, *17*, 320. [[CrossRef](#)] [[PubMed](#)]

5. Ruibal-Conti, A.L.; Ruiz, M.A.; Rodriguez, M.I.; Lerda, D.; Romero, M.D. Assessment of specific antibodies as biological indicators of human chronic exposure to microcystins. *Ecotoxicol. Environ. Saf.* **2019**, *175*, 236–242. [[CrossRef](#)] [[PubMed](#)]
6. Cirés, S.; Casero, M.C.; Quesada, A. Toxicity at the edge of life: A review on cyanobacterial toxins from extreme environments. *Mar. Drugs* **2017**, *15*, 233. [[CrossRef](#)] [[PubMed](#)]
7. Dittmann, E.; Fewer, D.P.; Neilan, B.A. Cyanobacterial toxins: Biosynthetic routes and evolutionary roots. *FEMS Microbiol. Rev.* **2013**, *37*, 23–43. [[CrossRef](#)] [[PubMed](#)]
8. Catherine, A.; Bernard, C.; Spoof, L.; Bruno, M. Microcystins and nodularins. *Handb. Cyanobacterial Monit. Cyanotoxin Anal.* **2017**, *11*, 107–126. [[CrossRef](#)]
9. Harada, K.; Ogawa, K.; Matsuura, K.; Murata, H.; Suzuki, M.; Watanabe, M.F.; Itezono, Y.; Nakayama, N. Structural determination of geometrical isomers of microcystins LR and RR from cyanobacteria by two-dimensional NMR spectroscopic techniques. *Chem. Res. Toxicol.* **1990**, *3*, 473–481. [[CrossRef](#)]
10. Meriluoto, J.; Spoof, L.; Codd, G.A. Appendix 3: Tables of Microcystins and Nodularins. *Handb. Cyanobacterial Monit. Cyanotoxin Anal.* **2017**, 526–537. [[CrossRef](#)]
11. Buratti, F.M.; Manganelli, M.; Vichi, S.; Stefanelli, M.; Scardala, S.; Testai, E.; Funari, E. Cyanotoxins: Producing organisms, occurrence, toxicity, mechanism of action and human health toxicological risk evaluation. *Arch. Toxicol.* **2017**, *91*, 1049–1130. [[CrossRef](#)] [[PubMed](#)]
12. Humans, I. IARC monographs on the evaluation of carcinogenic risks to humans. *J. Clin. Pathol.* **2010**, *94*, v–vii, 1. [[CrossRef](#)]
13. Salmaso, N.; Bernard, C.; Humbert, J.F.; Akçalan, R.; Albay, M.; Ballot, A.; Catherine, A.; Fastner, J.; Häggqvist, K.; Horecká, M.; et al. Basic guide to detection and monitoring of potentially toxic cyanobacteria. *Handb. Cyanobacterial Monit. Cyanotoxin Anal.* **2017**, *6*, 46–69. [[CrossRef](#)]
14. Tamele, I.J.; Silva, M.; Vasconcelos, V. The incidence of marine toxins and the associated seafood poisoning episodes in the African countries of the Indian Ocean and the Red Sea. *Toxins* **2019**, *11*, 58. [[CrossRef](#)] [[PubMed](#)]
15. Sangolkar, L.N.; Maske, S.S.; Chakrabarti, T. Methods for determining microcystins (peptide hepatotoxins) and microcystin-producing cyanobacteria. *Water Res.* **2006**, *40*, 3485–3496. [[CrossRef](#)]
16. Shen, F.; Xu, Y.; Jiang, S.; Zhao, B.; Shi, J.; Chen, J. Determination of microcystins in algal bloom samples by ultra performance liquid chromatography-tandem mass spectrometry and elimination of their matrix effects. *Chin. J. Chromatogr. (Se Pu)* **2017**, *35*, 772–777. [[CrossRef](#)] [[PubMed](#)]
17. Turner, A.D.; Waack, J.; Lewis, A.; Edwards, C.; Lawton, L. Development and single-laboratory validation of a UHPLC-MS/MS method for quantitation of microcystins and nodularin in natural water, cyanobacteria, shellfish and algal supplement tablet powders. *J. Chromatogr. B* **2018**, *1074–1075*, 111–123. [[CrossRef](#)] [[PubMed](#)]
18. Chen, Y.; Shen, D.; Fang, D. Nodularins in poisoning. *Clin. Chim. Acta* **2013**, *425*, 18–29. [[CrossRef](#)] [[PubMed](#)]
19. McGregor, G.B.; Sendall, B.C. Iningairema pulvinus gen nov., sp nov. (Cyanobacteria, Scytonemataceae) a new nodularin producer from Edgbaston Reserve, north-eastern Australia. *Harmful Algae* **2017**, *62*, 10–19. [[CrossRef](#)]
20. Shishido, T.K.; Jokela, J.; Humisto, A.; Suurnäkki, S.; Wahlsten, M.; Alvarenga, D.O.; Sivonen, K.; Fewer, D.P. The biosynthesis of rare homo-amino acid containing variants of microcystin by a benthic cyanobacterium. *Mar. Drugs* **2019**, *17*, 271. [[CrossRef](#)]
21. Valério, E.; Chaves, S.; Tenreiro, R. Diversity and impact of prokaryotic toxins on aquatic environments: A review. *Toxins* **2010**, *2*, 2359–2410. [[CrossRef](#)] [[PubMed](#)]
22. Testai, E.; Scardala, S.; Vichi, S.; Buratti, F.M.; Funari, E. Risk to human health associated with the environmental occurrence of cyanobacterial neurotoxic alkaloids anatoxins and saxitoxins. *Crit. Rev. Toxicol.* **2016**, *46*, 385–419. [[CrossRef](#)] [[PubMed](#)]
23. Pearson, L.A.; Dittmann, E.; Mazmouz, R.; Ongley, S.E.; D'Agostino, P.M.; Neilan, B.A. The genetics, biosynthesis and regulation of toxic specialized metabolites of cyanobacteria. *Harmful Algae* **2016**, *54*, 98–111. [[CrossRef](#)] [[PubMed](#)]
24. Bittencourt-Oliveira, M.d.C.; Piccin, V.; Kujbida, P.; Moura, A. Cylindrospermopsin in water supply reservoirs in Brazil determined by immunochemical and molecular methods. *J. Water Resour. Prot.* **2011**, *3*, 349–355. [[CrossRef](#)]

25. Kokocinski, M.; Maria Cameán, A.; Carmeli, S.; Guzmán-Guillén, R.; Jos, A.; Mankiewicz-Boczek, J.; Metcalf, J.; Moreno, I.; Isabel Prieto, A.; Sukenik, A. Cylindrospermopsin and congeners. *Handb. Cyanobacterial Monit. Cyanotoxin Anal.* **2017**, *12*, 127–137. [CrossRef]
26. Batsalova, T.; Basheva, D.; Bardarov, K.; Bardarov, V.; Dzhambazov, B.; Teneva, I. Assessment of the cytotoxicity, antioxidant activity and chemical composition of extracts from the cyanobacterium Fischerella major Gomont. *Chemosphere* **2019**, *218*, 93–103. [CrossRef] [PubMed]
27. Belykh, O.I.; Tikhonova, I.V.; Kuzmin, A.V.; Sorokovikova, E.G.; Fedorova, G.A.; Khanaev, I.V.; Sherbakova, T.A.; Timoshkin, O.A. First detection of benthic cyanobacteria in Lake Baikal producing paralytic shellfish toxins. *Toxicon* **2016**, *121*, 36–40. [CrossRef] [PubMed]
28. Wiese, M.; D'Agostino, P.M.; Mihali, T.K.; Moffitt, M.C.; Neilan, B.A. Neurotoxic alkaloids: Saxitoxin and its analogs. *Mar. Drugs* **2010**, *8*, 2185–2211. [CrossRef] [PubMed]
29. Namikoshi, M.; Murakami, T.; Watanabe, M.F.; Oda, T.; Yamada, J.; Tsujimura, S.; Nagai, H.; Oishi, S. Simultaneous production of homoanatoxin-a, anatoxin-a, and a new non-toxic 4-hydroxyhomoanatoxin-a by the cyanobacterium Raphidiopsis mediterranea Skuja. *Toxicon* **2003**, *42*, 533–538. [CrossRef]
30. Wood, S.A.; Biessy, L.; Puddick, J. Anatoxins are consistently released into the water of streams with Microcoleus autumnalis-dominated (cyanobacteria) proliferations. *Harmful Algae* **2018**, *80*, 88–95. [CrossRef]
31. Tan, L.T.; Goh, B.P.; Tripathi, A.; Lim, M.G.; Dickinson, G.H.; Lee, S.S.; Teo, S.L. Natural antifoulants from the marine cyanobacterium *Lyngbya majuscula*. *Biofouling* **2010**, *26*, 685–695. [CrossRef] [PubMed]
32. Cullen, A.; Pearson, L.A.; Mazmouz, R.; Liu, T.; Soeriyadi, A.H.; Ongley, S.E.; Neilan, B.A. Heterologous expression and biochemical characterisation of cyanotoxin biosynthesis pathways. *Nat. Prod. Rep.* **2018**, *36*, 1117–1136. [CrossRef] [PubMed]
33. Rastogi, R.P.; Madamwar, D.; Incharoensakdi, A. Bloom dynamics of cyanobacteria and their toxins: Environmental health impacts and mitigation strategies. *Front. Microbiol.* **2015**, *6*, 1254. [CrossRef] [PubMed]
34. Youssef, D.T.; Shaala, L.A.; Mohamed, G.A.; Ibrahim, S.R.; Banjar, Z.M.; Badr, J.M.; McPhail, K.L.; Risinger, A.L.; Mooberry, S.L. 2,3-seco-2,3-dioxo-lyngbyatoxin A from a Red Sea strain of the marine cyanobacterium *Moorea producens*. *Nat. Prod. Res.* **2015**, *29*, 703–709. [CrossRef] [PubMed]
35. Aimi, N.; Odaka, H.; Sakai, S.I.; Fujiki, H.; Suganuma, M.; Moore, R.E.; Patterson, G.M.L. Lyngbyatoxins B and C, two new irritants from *Lyngbya majuscula*. *J. Nat. Prod.* **1990**, *53*, 1593–1596. [CrossRef] [PubMed]
36. Ding, C.Y.G.; Pang, L.M.; Liang, Z.X.; Goh, K.K.K.; Glukhov, E.; Gerwick, W.H.; Tan, L.T. MS/MS-based molecular networking approach for the detection of aplysiatoxin-related compounds in environmental marine cyanobacteria. *Mar. Drugs* **2018**, *16*, 505. [CrossRef] [PubMed]
37. Nogle, L.M.; Okino, T.; Gerwick, W.H. Antillatoxin B, a neurotoxic lipopeptide from the marine cyanobacterium *Lyngbya majuscula*. *J. Nat. Prod.* **2001**, *64*, 983–985. [CrossRef]
38. Orjala, J.; Nagle, D.G.; Hsu, V.; Gerwick, W.H. Antillatoxin: An exceptionally ichthyotoxic cyclic lipopeptide from the tropical cyanobacterium *Lyngbya majuscula*. *J. Am. Chem. Soc.* **1995**, *117*, 8281–8282. [CrossRef]
39. Boudreau, P.D.; Monroe, E.A.; Mehrotra, S.; Desfor, S.; Korobeynikov, A.; Sherman, D.H.; Murray, T.F.; Gerwick, L.; Dorrestein, P.C.; Gerwick, W.H. Expanding the described metabolome of the marine cyanobacterium *Moorea producens* jhb through orthogonal natural products workflows. *PLoS ONE* **2015**, *10*, e0133297. [CrossRef]
40. Edwards, D.J.; Marquez, B.L.; Nogle, L.M.; McPhail, K.; Goeger, D.E.; Roberts, M.A.; Gerwick, W.H. Structure and biosynthesis of the jamaicamides, new mixed polyketide-peptide neurotoxins from the marine cyanobacterium *Lyngbya majuscula*. *Chem. Biol.* **2004**, *11*, 817–833. [CrossRef]
41. Umezawa, T.; Sueda, M.; Kamura, T.; Kawahara, T.; Han, X.; Okino, T.; Matsuda, F. Synthesis and biological activity of kalkitoxin and its analogues. *J. Org. Chem.* **2012**, *77*, 357–370. [CrossRef] [PubMed]
42. Wu, M.; Okino, T.; Nogle, L.M.; Marquez, B.L.; Williamson, R.T.; Sitachitta, N.; Berman, F.W.; Murray, T.F.; McGough, K.; Jacobs, R.; et al. Structure, synthesis, and biological properties of kalkitoxin, a novel neurotoxin from the marine cyanobacterium *Lyngbya majuscula*. *J. Am. Chem. Soc.* **2000**, *122*, 12041–12042. [CrossRef]
43. Kothari, A.; Vaughn, M.; Garciapichel, F. Comparative genomic analyses of the cyanobacterium, *Lyngbya aestuarii* BL J, a powerful hydrogen producer. *Front. Microbiol.* **2013**, *4*, 363. [CrossRef] [PubMed]
44. Caro-Diaz, E.J.E.; Valeriote, F.A.; Gerwick, W.H. Highly convergent total synthesis and assignment of absolute configuration of majusculamide D, a potent and selective cytotoxic metabolite from *Moorea* sp. *Org. Lett.* **2019**, *21*, 793–796. [CrossRef] [PubMed]

45. Carter, D.C.; Moore, R.E.; Mynderse, J.S.; Niemczura, W.P.; Todd, J.S. Structure of majusculamide C, a cyclic depsipeptide from *Lyngbya majuscula*. *J. Org. Chem.* **1984**, *49*. [[CrossRef](#)]
46. Suntornchashwej, S.; Chaichit, N.; Isobe, M.; Suwanborirux, K. Hectochlorin and Morpholine Derivatives from the Thai Sea Hare, *Bursatella leachii*. *J. Nat. Prod.* **2005**, *68*, 951–955. [[CrossRef](#)] [[PubMed](#)]
47. Ramaswamy, A.V.; Sorrels, C.M.; Gerwick, W.H. Cloning and biochemical characterization of the hectochlorin biosynthetic gene cluster from the marine cyanobacterium *Lyngbya majuscula*. *J. Nat. Prod.* **2007**, *70*, 1977–1986. [[CrossRef](#)] [[PubMed](#)]
48. Burja, A.M.; Abou-Mansour, E.; Banaigs, B.; Payri, C.; Burgess, J.G.; Wright, P.C. Culture of the marine cyanobacterium, *Lyngbya majuscula* (Oscillatoriaceae), for bioprocess intensified production of cyclic and linear lipopeptides. *J. Microbiol. Methods* **2002**, *48*, 207–219. [[CrossRef](#)]
49. Yoo, H.D.; Gerwick, W.H. Curacins B and C, new antimitotic natural products from the marine cyanobacterium *Lyngbya majuscula*. *J. Nat. Prod.* **1995**, *58*, 1961–1965. [[CrossRef](#)]
50. Zhang, C.; Naman, C.B.; Engene, N.; Gerwick, W.H. Laucysteinamide A, a hybrid PKS/NRPS metabolite from a Saipan Cyanobacterium, cf. *Caldora penicillata*. *Mar. Drugs* **2017**, *15*, 121. [[CrossRef](#)]
51. Márquez, B.; Verdier-Pinard, P.; Hamel, E.; Gerwick, W.H. Curacin D, an antimitotic agent from the marine cyanobacterium *Lyngbya majuscula*. *Phytochemistry* **1998**, *49*, 2387–2389. [[CrossRef](#)]
52. Rodgers, K.J.; Main, B.J.; Samardzic, K. Cyanobacterial neurotoxins: Their occurrence and mechanisms of toxicity. *Neurotox. Res.* **2018**, *33*, 168–177. [[CrossRef](#)] [[PubMed](#)]
53. Banack, S.A.; Murch, S.J. Methods for the chemical analysis of β-N-Methylamino-L-A lanine: What is known and what remains to be determined. *Neurotox. Res.* **2018**, *33*, 184–191. [[CrossRef](#)] [[PubMed](#)]
54. Monteiro, M.; Costa, M.; Moreira, C.; Vasconcelos, V.M.; Baptista, M.S. Screening of BMAA-producing cyanobacteria in cultured isolates and in situ blooms. *J. Appl. Phycol.* **2017**, *29*, 879–888. [[CrossRef](#)]
55. Lage, S.; Burian, A.; Rasmussen, U.; Costa, P.R.; Annadotter, H.; Godhe, A.; Rydberg, S. BMAA extraction of cyanobacteria samples: Which method to choose? *Environ. Sci. Pollut. Res. Int.* **2016**, *23*, 338–350. [[CrossRef](#)]
56. Manolidi, K.; Triantis, T.M.; Kaloudis, T.; Hiskia, A. Neurotoxin BMAA and its isomeric amino acids in cyanobacteria and cyanobacteria-based food supplements. *J. Hazard. Mater.* **2019**, *365*, 346–365. [[CrossRef](#)]
57. Kehr, J.C.; Dittmann, E. Biosynthesis and function of extracellular glycans in cyanobacteria. *Life (Basel)* **2015**, *5*, 164–180. [[CrossRef](#)]
58. Hinojosa, M.G.; Gutierrez-Praena, D.; Prieto, A.I.; Guzman-Guillen, R.; Jos, A.; Camean, A.M. Neurotoxicity induced by microcystins and cylindrospermopsin: A review. *Sci. Total Environ.* **2019**, *668*, 547–565. [[CrossRef](#)]
59. Ohtani, I.; Moore, R.E.; Runnegar, M. Cylindrospermopsin: A potent hepatotoxin from the blue-green alga *Cylindrospermopsis raciborskii*. *ChemInform* **1992**, *114*, 7941–7942. [[CrossRef](#)]
60. Hinojosa, M.G.; Prieto, A.I.; Gutiérrez-Praena, D.; Moreno, F.J.; Cameán, A.M.; Jos, A. Neurotoxic assessment of Microcystin-LR, cylindrospermopsin and their combination on the human neuroblastoma SH-SY5Y cell line. *Chemosphere* **2019**, *224*, 751–764. [[CrossRef](#)]
61. Bourke, A.T.C.; Hawes, R.B.; Neilson, A.; Stallman, N.D. An outbreak of hepato-enteritis (the Palm Island mystery disease) possibly caused by algal intoxication. *Toxicon* **1983**, *21*, 45–48. [[CrossRef](#)]
62. Harada, K.I.; Ohtani, I.; Iwamoto, K.; Suzuki, M.; Watanabe, M.F.; Watanabe, M.; Terao, K. Isolation of cylindrospermopsin from a cyanobacterium *Umezakia natans* and its screening method. *Toxicon* **1994**, *32*, 73–84. [[CrossRef](#)]
63. Moreira, C.; Azevedo, J.; Antunes, A.; Vasconcelos, V. Cylindrospermopsin: Occurrence, methods of detection and toxicology. *J. Appl. Microbiol.* **2013**, *114*, 605–620. [[CrossRef](#)]
64. Guzmán-Guillén, R.; Prieto, A.I.; González, A.G.; Soria-Díaz, M.E.; Cameán, A.M. Cylindrospermopsin determination in water by LC-MS/MS: Optimization and validation of the method and application to real samples. *Environ. Toxicol. Chem.* **2012**, *31*, 2233–2238. [[CrossRef](#)]
65. Wong, J.L.; Oesterlin, R.; Rapoport, H. The structure of saxitoxin. *J. Am. Chem. Soc.* **1971**, *93*, 7344–7345. [[CrossRef](#)]
66. Ember, L. Chemical weapons convention. *Chem. Eng. News Arch.* **1998**, *76*, 9. [[CrossRef](#)]
67. Hallegraeff, G. Harmful algal blooms: A global overview. *Camb. Law J.* **1995**, *9*, 268–270. [[CrossRef](#)]
68. Humpage, A.R.; Magalhaes, V.F.; Froscio, S.M. Comparison of analytical tools and biological assays for detection of paralytic shellfish poisoning toxins. *Anal. Bioanal. Chem.* **2010**, *397*, 1655–1671. [[CrossRef](#)]
69. Cusick, K.D.; Sayler, G.S. An overview on the marine neurotoxin, saxitoxin: Genetics, molecular targets, methods of detection and ecological functions. *Mar. Drugs* **2013**, *11*, 991–1018. [[CrossRef](#)]

70. Haddad, S.P.; Bobbitt, J.M.; Taylor, R.B.; Lovin, L.M.; Conkle, J.L.; Chambliss, C.K.; Brooks, B.W. Determination of microcystins, nodularin, anatoxin-a, cylindrospermopsin, and saxitoxin in water and fish tissue using isotope dilution liquid chromatography tandem mass spectrometry. *J. Chromatogr. A* **2019**. [[CrossRef](#)]
71. Bragg, W.A.; Lemire, S.W.; Coleman, R.M.; Hamelin, E.I.; Johnson, R.C. Detection of human exposure to saxitoxin and neosaxitoxin in urine by online-solid phase extraction-liquid chromatography-tandem mass spectrometry. *Toxicon* **2015**, *99*, 118–124. [[CrossRef](#)]
72. Devlin, J.P.; Edwards, O.E.; Gorham, P.R.; Hunter, N.R.; Pike, R.K.; Stavric, B. Anatoxin-a, a toxic alkaloid from *Anabaena flos-aquae* NRC-44h. *Can. J. Chem.* **1977**, *55*, 1367–1371. [[CrossRef](#)]
73. Wonnacott, S.; Swanson, K.L.; Albuquerque, E.X.; Huby, N.J.S.; Thompson, P.; Gallagher, T. Homoanatoxin: A potent analogue of anatoxin-a. *Biochem. Pharmacol.* **1992**, *43*, 419–423. [[CrossRef](#)]
74. Bruno, M.; Ploux, O.; Metcalf, J.; Méjean, A.; Pawlik-Skowrońska, B.; Furey, A. Anatoxin-a, homoanatoxin-a, and natural analogues. *Handb. Cyanobacterial Monit. Cyanotoxin Anal.* **2017**, *13*, 138–147. [[CrossRef](#)]
75. Osswald, J.; Rellán, S.; Gago, A.; Vasconcelos, V. Toxicology and detection methods of the alkaloid neurotoxin produced by cyanobacteria, anatoxin-a. *Environ. Int.* **2007**, *33*, 1070–1089. [[CrossRef](#)]
76. Legrand, B.; Lesobre, J.; Colombet, J.; Latour, D.; Sabart, M. Molecular tools to detect anatoxin-a genes in aquatic ecosystems: Toward a new nested PCR-based method. *Harmful Algae* **2016**, *58*, 16–22. [[CrossRef](#)]
77. Van Apeldoorn, M.E.; van Egmond, H.P.; Speijers, G.J.; Bakker, G.J. Toxins of cyanobacteria. *Mol. Nutr. Food Res.* **2007**, *51*, 7–60. [[CrossRef](#)]
78. Matsunaga, S.; Moore, R.E.; Niemczura, W.P.; Carmichael, W.W. Anatoxin-a(s), a potent anticholinesterase from *Anabaena flos-aquae*. *J. Am. Chem. Soc.* **1989**, *111*, 8021–8023. [[CrossRef](#)]
79. Mahmood, N.A.; Carmichael, W.W. The pharmacology of anatoxin-a(s), a neurotoxin produced by the freshwater cyanobacterium *Anabaena flos-aquae* NRC 525-17. *Toxicon* **1986**, *24*, 425–434. [[CrossRef](#)]
80. Marie, B.; Blaha, L.; Maria Cameán, A.; Fessard, V.; Gutiérrez-Praena, D.; Jos, A.; Metcalf, J.; Pichardo, S.; Puerto, M.; Törökne, A.; et al. Bioassay use in the field of toxic cyanobacteria. *Handb. Cyanobacterial Monit. Cyanotoxin Anal.* **2016**, *27*, 272–278. [[CrossRef](#)]
81. Metcalf, J.; Bruno, M. Anatoxin-a(S). *Handb. Cyanobacterial Monit. Cyanotoxin Anal.* **2017**, *15*, 155–159. [[CrossRef](#)]
82. Villatte, F.; Schulze, H.; Schmid, R.D.; Bachmann, T.T. A disposable acetylcholinesterase-based electrode biosensor to detect anatoxin-a(s) in water. *Anal. Bioanal. Chem.* **2002**, *372*, 322–326. [[CrossRef](#)]
83. Cardellina, J.H.; Marner, F.J.; Moore, R.E. Seaweed dermatitis: Structure of lyngbyatoxin A. *Science* **1979**, *204*, 193. [[CrossRef](#)]
84. Fujiki, H.; Mori, M.; Nakayasu, M.; Terada, M.; Sugimura, T.; Moore, R.E. Indole alkaloids: Dihydrotetrocidin B, teleocidin, and lyngbyatoxin A as members of a new class of tumor promoters. *Proc. Natl. Acad. Sci. USA* **1981**, *78*, 3872–3876. [[CrossRef](#)]
85. Ito, E.; Satake, M.; Yasumoto, T. Pathological effects of lyngbyatoxin A upon mice. *Toxicon* **2002**, *40*, 551–556. [[CrossRef](#)]
86. Jiang, W.; Zhou, W.; Uchida, H.; Kikumori, M.; Irie, K.; Watanabe, R.; Suzuki, T.; Sakamoto, B.; Kamio, M.; Nagai, H. A new lyngbyatoxin from the Hawaiian cyanobacterium *Moorea producens*. *Mar. Drugs* **2014**, *12*, 2748–2759. [[CrossRef](#)]
87. Esposito, G.; Teta, R.; Marrone, R.; De Sterlich, C.; Casazza, M.; Anastasio, A.; Lega, M.; Costantino, V. A Fast Detection Strategy for Cyanobacterial blooms and associated cyanotoxins (FDSCC) reveals the occurrence of lyngbyatoxin A in campania (South Italy). *Chemosphere* **2019**, *225*, 342–351. [[CrossRef](#)]
88. Kato, Y.; Scheuer, P.J. Aplysiatoxin and debromoaplysiatoxin, constituents of the marine mollusk *Stylocheilus longicauda* (Quoy and Gaimard, 1824). *J. Am. Chem. Soc.* **1974**, *96*, 2245–2246. [[CrossRef](#)]
89. Fujiki, H.; Ikegami, K.; Hakii, H.; Saganuma, M.; Yamaizumi, Z.; Yamazato, K.; Moore, R.E.; Sugimura, T. A blue-green alga from Okinawa contains aplysiatoxins, the third class of tumor promoters. *Jpn. J. Cancer Res.* **1985**, *76*, 257–259.
90. Richard, K.; Williams, D.E.; de Silva, E.D.; Brockman, M.A.; Brumme, Z.L.; Andersen, R.J.; Tietjen, I. Identification of novel HIV-1 latency-reversing agents from a library of marine natural products. *Viruses* **2018**, *10*, 348. [[CrossRef](#)]
91. Li, W.I.; Marquez, B.L.; Okino, T.; Yokokawa, F.; Shioiri, T.; Gerwick, W.H.; Murray, T.F. Characterization of the preferred stereochemistry for the neuropharmacologic actions of antillatoxin. *J. Nat. Prod.* **2004**, *67*, 559–568. [[CrossRef](#)]

92. Jabba, S.V.; Prakash, A.; Dravid, S.M.; Gerwick, W.H.; Murray, T.F. Antillatoxin, a novel lipopeptide, enhances neurite outgrowth in immature cerebrocortical neurons through activation of voltage-gated sodium channels. *J. Pharmacol. Exp. Ther.* **2010**, *332*, 698–709. [CrossRef]
93. Okura, K.; Matsuoka, S.; Inoue, M. The bulky side chain of antillatoxin is important for potent toxicity: Rational design of photoresponsive cytotoxins based on SAR studies. *Chem. Commun.* **2013**, *49*, 8024–8026. [CrossRef]
94. Graf, K.M.; Tabor, M.G.; Brown, M.L.; Paige, M. Synthesis of (S)-jamaicamide C carboxylic acid. *Org. Lett.* **2009**, *11*, 5382–5385. [CrossRef]
95. Araoz, R.; Molgo, J.; de Marsac, N.T. Neurotoxic cyanobacterial toxins. *Toxicon* **2010**, *56*, 813–828. [CrossRef]
96. Morgan, J.B.; Liu, Y.; Coothankandaswamy, V.; Mahdi, F.; Jekabsons, M.B.; Gerwick, W.H.; Valeriote, F.A.; Zhou, Y.D.; Nagle, D.G. Kalkitoxin inhibits angiogenesis, disrupts cellular hypoxic signaling, and blocks mitochondrial electron transport in tumor cells. *Mar. Drugs* **2015**, *13*, 1552–1568. [CrossRef]
97. Orjala, J.; Gerwick, W.H. Barbamide, a chlorinated metabolite with molluscicidal activity from the Caribbean Cyanobacterium *Lyngbya majuscula*. *J. Nat. Prod.* **1996**, *59*, 427–430. [CrossRef]
98. Sitachitta, N.; Márquez, B.L.; Williamson, R.T.; Rossi, J.; Roberts, M.A.; Gerwick, W.H.; Nguyen, V.A.; Willis, C.L.J.T. Biosynthetic pathway and origin of the chlorinated methyl group in barbamide and dechlorobarbamide, metabolites from the marine cyanobacterium *Lyngbya majuscula*. *Tetrahedron* **2000**, *56*, 9103–9113. [CrossRef]
99. Cetusic, J.R.P.; Green, F.R.; Graupner, P.R.; Oliver, M.P. Total synthesis of hectochlorin. *Org. Lett.* **2002**, *4*, 1307–1310. [CrossRef]
100. Marquez, B.L.; Watts, K.S.; Yokochi, A.; Roberts, M.A.; Verdier-Pinard, P.; Jimenez, J.I.; Hamel, E.; Scheuer, P.J.; Gerwick, W.H. Structure and absolute stereochemistry of hectochlorin, a potent stimulator of actin assembly. *J. Nat. Prod.* **2002**, *65*, 866–871. [CrossRef]
101. Timmins, A.; Fowler, N.J.; Warwicker, J.; Straganz, G.D.; de Visser, S.P. Does substrate positioning affect the selectivity and reactivity in the hectochlorin biosynthesis halogenase? *Front. Chem.* **2018**, *6*, 513. [CrossRef] [PubMed]
102. Ueoka, R.; Hitora, Y.; Ito, A.; Yoshida, M.; Okada, S.; Takada, K.; Matsunaga, S. Curacin E from the Brittle Star *Ophiocoma scolopendrina*. *J. Nat. Prod.* **2016**, *79*, 2754–2757. [CrossRef] [PubMed]
103. Gerwick, W.H.; Proteau, P.J.; Nagle, D.G.; Hamel, E.; Slate, D.L.J.C. Structure of curacin A, a novel antimitotic, antiproliferative and brine shrimp toxic natural product from the marine cyanobacterium *Lyngbya majuscula*. *J. Organic Chem.* **1994**, *59*. [CrossRef]
104. Chang, Z.; Sitachitta, N.; Rossi, J.V.; Roberts, M.A.; Flatt, P.M.; Jia, J.; Sherman, D.H.; Gerwick, W.H. Biosynthetic pathway and gene cluster analysis of curacin A, an antitubulin natural product from the tropical marine cyanobacterium *Lyngbya majuscula*. *J. Nat. Prod.* **2004**, *67*, 1356–1367. [CrossRef] [PubMed]
105. Fiers, W.D.; Dodge, G.J.; Sherman, D.H.; Smith, J.L.; Aldrich, C.C. Vinyllogous dehydration by a polyketide dehydratase domain in curacin biosynthesis. *J. Am. Chem. Soc.* **2016**, *138*, 16024–16036. [CrossRef] [PubMed]
106. Vega, A.; Bell, E.A.  $\alpha$ -Amino- $\beta$ -methylaminopropionic acid, a new amino acid from seeds of *Cycas circinalis*. *Phytochemistry* **1967**, *6*, 759–762. [CrossRef]
107. Ploux, O.; Combes, A.; Eriksson, J.; Metcalf, J.  $\beta$ -N-methylamino-L-alanine and (S)-2,4-diaminobutyric acid. *Handb. Cyanobacterial Monit. Cyanotoxin Anal.* **2017**, *16*, 160–164. [CrossRef]
108. Main, B.J.; Bowling, L.C.; Padula, M.P.; Bishop, D.P.; Mitrović, S.M.; Guillemin, G.J.; Rodgers, K.J. Detection of the suspected neurotoxin  $\beta$ -methylamino-L-alanine (BMAA) in cyanobacterial blooms from multiple water bodies in Eastern Australia. *Harmful Algae* **2018**, *74*, 10–18. [CrossRef]
109. Cox, P.A.; Kostrzewska, R.M.; Guillemin, G.J. BMAA and neurodegenerative illness. *Neurotox. Res.* **2018**, *33*, 178–183. [CrossRef]
110. Banack, S.A.; Cox, P.A. Biomagnification of cycad neurotoxins in flying foxes: Implications for ALS-PDC in Guam. *Neurology* **2003**, *61*, 387–389. [CrossRef]
111. Porojan, C.; Mitrović, S.M.; Yeo, D.C.; Furey, A. Overview of the potent cyanobacterial neurotoxin  $\beta$ -methylamino-L-alanine (BMAA) and its analytical determination. *Food Addit. Contam. Part A* **2016**, *33*, 1570–1586. [CrossRef] [PubMed]
112. Ballot, A.; Bernard, C.; Fastner, J. Saxitoxin and analogues. *Handb. Cyanobacterial Monit. Cyanotoxin Anal.* **2017**, *14*, 148–154. [CrossRef]

113. Banack, S.A.; Johnson, H.E.; Cheng, R.; Cox, P.A. Production of the neurotoxin BMAA by a marine cyanobacterium. *Mar. Drugs* **2007**, *5*, 180–196. [CrossRef] [PubMed]
114. Kerrin, E.S.; White, R.L.; Quilliam, M.A. Quantitative determination of the neurotoxin beta-N-methylamino-L-alanine (BMAA) by capillary electrophoresis-tandem mass spectrometry. *Anal. Bioanal. Chem.* **2017**, *409*, 1481–1491. [CrossRef] [PubMed]
115. Wiegand, C.; Pflugmacher, S. Ecotoxicological effects of selected cyanobacterial secondary metabolites a short review. *Toxicol. Appl. Pharmacol.* **2005**, *203*, 201–218. [CrossRef] [PubMed]
116. Durai, P.; Batool, M.; Choi, S. Structure and effects of cyanobacterial lipopolysaccharides. *Mar. Drugs* **2015**, *13*, 4217–4230. [CrossRef] [PubMed]
117. Rapala, J.; Lahti, K.; Rasanen, L.A.; Esala, A.L.; Niemela, S.I.; Sivonen, K. Endotoxins associated with cyanobacteria and their removal during drinking water treatment. *Water Res.* **2002**, *36*, 2627–2635. [CrossRef]
118. Best, J.H.; Pflugmacher, S.; Wiegand, C.; Eddy, F.B.; Metcalf, J.S.; Codd, G.A. Effects of enteric bacterial and cyanobacterial lipopolysaccharides, and of microcystin-LR, on glutathione S-transferase activities in zebra fish (*Danio rerio*). *Aquat. Toxicol.* **2002**, *60*, 223–231. [CrossRef]
119. Bláhová, L.; Blaha, L. Extraction, purification, and testing of LPS from cyanobacterial samples. *Handb. Cyanobacterial Monit. Cyanotoxin Anal.* **2017**, *56*, 447–451. [CrossRef]
120. Zhang, C.; Tian, F.; Zhang, M.; Zhang, Z.; Bai, M.; Guo, G.; Zheng, W.; Wang, Q.; Shi, Y.; Wang, L. Endotoxin contamination, a potentially important inflammation factor in water and wastewater: A review. *Sci. Total Environ.* **2019**, *681*, 365–378. [CrossRef]
121. Blahova, L.; Adamovsky, O.; Kubala, L.; Svhalkova Sindlerova, L.; Zounkova, R.; Blaha, L. The isolation and characterization of lipopolysaccharides from *Microcystis aeruginosa*, a prominent toxic water bloom forming cyanobacteria. *Toxicon* **2013**, *76*, 187–196. [CrossRef] [PubMed]
122. Reta, N.; Michelmore, A.; Saint, C.P.; Prieto-Simon, B.; Voelcker, N.H. Label-free bacterial toxin detection in water supplies using porous silicon nanochannel sensors. *ACS Sens.* **2019**, *4*, 1515–1523. [CrossRef] [PubMed]
123. Liu, P.; Wei, J.; Yang, K.; Massey, I.Y.; Guo, J.; Zhang, C.; Yang, F. Isolation, molecular identification, and characterization of a unique toxic cyanobacterium *Microcystis* sp. found in Hunan Province, China. *J. Toxicol. Environ. Health Part A* **2018**, *81*, 1142–1149. [CrossRef] [PubMed]
124. Yang, F.; Guo, J.; Huang, F.; Massey, I.Y.; Huang, R.; Li, Y.; Wen, C.; Ding, P.; Zeng, W.; Liang, G. Removal of microcystin-LR by a novel native effective bacterial community designated as YFMCD4 isolated from Lake Taihu. *Toxins* **2018**, *10*. [CrossRef] [PubMed]
125. Zhang, W.; Lou, I.; Ung, W.K.; Kong, Y.; Mok, K.M. Analysis of cylindrospermopsin- and microcystin-producing genotypes and cyanotoxin concentrations in the Macau storage reservoir. *Hydrobiologia* **2014**, *741*, 51–68. [CrossRef]
126. Huang, W.J.; Lai, C.H.; Cheng, Y.L. Evaluation of extracellular products and mutagenicity in cyanobacteria cultures separated from a eutrophic reservoir. *Sci. Total Environ.* **2007**, *377*, 214–223. [CrossRef]
127. Takahashi, T.; Umebara, A.; Tsutsumi, H. Diffusion of microcystins (cyanobacteria hepatotoxins) from the reservoir of Isahaya Bay, Japan, into the marine and surrounding ecosystems as a result of large-scale drainage. *Mar. Pollut. Bull.* **2014**, *89*, 250–258. [CrossRef]
128. Joung, S.H.; Oh, H.M.; Ko, S.R.; Ahn, C.Y. Correlations between environmental factors and toxic and non-toxic *Microcystis* dynamics during bloom in Daechung Reservoir, Korea. *Harmful Algae* **2011**, *10*, 188–193. [CrossRef]
129. Welker, M.; Khan, S.; Haque, M.M.; Islam, S.; Khan, N.H.; Chorus, I.; Fastner, J. Microcystins (cyanobacterial toxins) in surface waters of rural Bangladesh: Pilot study. *J. Water Health* **2005**, *3*, 325–337. [CrossRef]
130. Shu Harn, T.; Gin, K. The dynamics of cyanobacteria and microcystin production in a tropical reservoir of Singapore. *Harmful Algae* **2010**, *10*, 1–44. [CrossRef]
131. Mohamed, Z.A.; Al Shehri, A.M. Microcystins in groundwater wells and their accumulation in vegetable plants irrigated with contaminated waters in Saudi Arabia. *J. Hazard. Mater.* **2009**, *172*, 310–315. [CrossRef]
132. Singh, S.; Asthana, R.K. Assessment of microcystin concentration in carp and catfish: A case study from Lakshmikund pond, Varanasi, India. *Bull. Environ. Contam. Toxicol.* **2014**, *92*, 687–692. [CrossRef]
133. Cuvín-Aralar, M.L.; Fastner, J.; Focken, U.; Becker, K.; Aralar, E.V. Microcystins in natural blooms and laboratory cultured *Microcystis aeruginosa* from Laguna de Bay, Philippines. *Syst. Appl. Microbiol.* **2002**, *25*, 179–182. [CrossRef]

134. Somdee, T.; Kaewsan, T.; Somdee, A. Monitoring toxic cyanobacteria and cyanotoxins (microcystins and cylindrospermopsins) in four recreational reservoirs (Khon Kaen, Thailand). *Environ. Monit. Assess.* **2013**, *185*, 9521–9529. [[CrossRef](#)]
135. Dao, T.S.; Nimptsch, J.; Wiegand, C. Dynamics of cyanobacteria and cyanobacterial toxins and their correlation with environmental parameters in Tri An Reservoir, Vietnam. *J. Water Health* **2016**, *14*, 699–712. [[CrossRef](#)]
136. Trung, B.; Dao, T.S.; Faassen, E.; Lurling, M. Cyanobacterial blooms and Microcystins in Southern Vietnam. *Toxins* **2018**, *10*, 471. [[CrossRef](#)]
137. Lifshits, M.; Carmeli, S. Metabolites of *Microcystis aeruginosa* bloom material from Lake Kinneret, Israel. *J. Nat. Prod.* **2012**, *75*, 209–219. [[CrossRef](#)]
138. Rodriguez, I.; Fraga, M.; Alfonso, A.; Guillebault, D.; Medlin, L.; Baudart, J.; Jacob, P.; Helmi, K.; Meyer, T.; Breitenbach, U.; et al. Monitoring of freshwater toxins in European environmental waters by using novel multi-detection methods. *Environ. Toxicol. Chem.* **2017**, *36*, 645–654. [[CrossRef](#)]
139. Li, R.; Carmichael, W.W.; Brittain, S.; Eaglesham, G.K.; Shaw, G.R.; Liu, Y.; Watanabe, M.M. First report of the cyanotoxins cylindrospermopsin and deoxycylindrospermopsin from *Raphidiopsis curvata* (Cyanobacteria). *J. Phycol.* **2001**, *37*, 1121–1126. [[CrossRef](#)]
140. Chonudomkul, D.; Yongmanitchai, W.; Theeragool, G.; Kawachi, M.; Kasai, F.; Kaya, K.; Watanabe, M.M. Morphology, genetic diversity, temperature tolerance and toxicity of *Cylindrospermopsis raciborskii* (Nostocales, Cyanobacteria) strains from Thailand and Japan. *FEMS Microbiol. Ecol.* **2004**, *48*, 345–355. [[CrossRef](#)]
141. Mohamed, Z.A.; Al-Shehri, A.M. Assessment of cylindrospermopsin toxin in an arid Saudi lake containing dense cyanobacterial bloom. *Environ. Monit. Assess.* **2013**, *185*, 2157–2166. [[CrossRef](#)]
142. Dao, T.S.; Ortiz-Rodriguez, R.; Do-Hong, L.C.; Wiegand, C. Non-microcystin and non-cylindrospermopsin producing cyanobacteria affect the biochemical responses and behavior of *Daphnia magna*. *Int. Rev. Hydrobiol.* **2013**, *98*, 235–244. [[CrossRef](#)]
143. Wimmer, K.M.; Strangman, W.K.; Wright, J.L.C. 7-Deoxy-desulfo-cylindrospermopsin and 7-deoxy-desulfo-12-acetylcytindrospermopsin: Two new cylindrospermopsin analogs isolated from a Thai strain of *Cylindrospermopsis raciborskii*. *Harmful Algae* **2014**, *37*, 203–206. [[CrossRef](#)]
144. Ballot, A.; Ramm, J.; Rundberget, T.; Kaplan-Levy, R.N.; Hadas, O.; Sukenik, A.; Wiedner, C. Occurrence of non-cylindrospermopsin-producing *Aphanizomenon ovalisporum* and *Anabaena bergii* in Lake Kinneret (Israel). *J. Plankton Res.* **2011**, *33*, 1736–1746. [[CrossRef](#)]
145. Akcaalan, R.; Koker, L.; Oguz, A.; Spoof, L.; Meriluoto, J.; Albay, M. First report of cylindrospermopsin production by two cyanobacteria (*Dolichospermum mendotae* and *Chrysosporum ovalisporum*) in Lake Iznik, Turkey. *Toxins* **2014**, *6*, 3173–3186. [[CrossRef](#)]
146. Zhang, H.; Yang, S.; Beier, R.C.; Beloglazova, N.V.; Lei, H.; Sun, X.; Ke, Y.; Zhang, S.; Wang, Z. Simple, high efficiency detection of microcystins and nodularin-R in water by fluorescence polarization immunoassay. *Anal. Chim. Acta* **2017**, *992*, 119–127. [[CrossRef](#)]
147. Wu, H.Y.; Luan, Q.S.; Guo, M.M.; Gu, H.F.; Zhai, Y.X.; Tan, Z.J. Phycotoxins in scallops (*Patinopecten yessoensis*) in relation to source, composition and temporal variation of phytoplankton and cysts in North Yellow Sea, China. *Mar. Pollut. Bull.* **2018**, *135*, 1198–1204. [[CrossRef](#)]
148. Shin, C.; Jo, H.; Kim, S.H.; Kang, G.J. Exposure assessment to paralytic shellfish toxins through the shellfish consumption in Korea. *Food Res. Int.* **2018**, *108*, 274–279. [[CrossRef](#)]
149. Shunmugam, S.; Gayathri, M.; Prasannabali, N.; Thajuddin, N.; Muralitharan, G. Unraveling the presence of multi-class toxins from *Trichodesmium* bloom in the Gulf of Mannar region of the Bay of Bengal. *Toxicon* **2017**, *135*, 43–50. [[CrossRef](#)]
150. James Holmes, M.; Bolch, C.; Green, D.; Cembella, A.; Lay Ming Teo, S. Singapore isolates of the dinoflagellate *Gymnodinium catenatum* (Dinophyceae) produce a unique profile of paralytic shellfish toxins. *J. Phycol.* **2002**, *38*, 96–106. [[CrossRef](#)]
151. Zaman, L.; Arakawa, O.; Shimosu, A.; Onoue, Y. Occurrence of paralytic shellfish poison in Bangladeshi freshwater puffers. *Toxicon* **1997**, *35*, 423–431. [[CrossRef](#)]
152. Yilmaz, M.; Foss, A.J.; Selwood, A.I.; Ozen, M.; Boundy, M. Paralytic shellfish toxin producing *Aphanizomenon gracile* strains isolated from Lake Iznik, Turkey. *Toxicon* **2018**, *148*, 132–142. [[CrossRef](#)]
153. Srivastava, A.; Ahn, C.Y.; Asthana, R.K.; Lee, H.G.; Oh, H.M. Status, alert system, and prediction of cyanobacterial bloom in South Korea. *Biomed. Res. Int.* **2015**, *2015*, 584696. [[CrossRef](#)]

154. Metcalf, J.S.; Richer, R.; Cox, P.A.; Codd, G.A. Cyanotoxins in desert environments may present a risk to human health. *Sci. Total Environ.* **2012**, *421–422*, 118–123. [[CrossRef](#)]
155. Jiao, Y.; Chen, Q.; Chen, X.; Wang, X.; Liao, X.; Jiang, L.; Wu, J.; Yang, L. Occurrence and transfer of a cyanobacterial neurotoxin  $\beta$ -methylamino-L-alanine within the aquatic food webs of Gonghu Bay (Lake Taihu, China) to evaluate the potential human health risk. *Sci. Total Environ.* **2014**, *468–469*, 457–463. [[CrossRef](#)]
156. Caparros-Lefebvre, D.; Steele, J.; Kotake, Y.; Ohta, S. Geographic isolates of atypical Parkinsonism and tauopathy in the tropics: Possible synergy of neurotoxins. *Mov. Disord.* **2006**, *21*, 1769–1771. [[CrossRef](#)]
157. Craighead, D.; Metcalf, J.S.; Banack, S.A.; Amgalan, L.; Reynolds, H.V.; Batmunkh, M. Presence of the neurotoxic amino acids beta-N-methylamino-L-alanine (BMAA) and 2,4-diamino-butyric acid (DAB) in shallow springs from the Gobi Desert. *Amyotroph. Lateral Scler.* **2009**, *10* (Suppl. 2), 96–100. [[CrossRef](#)]
158. Major, Y.; Kifle, D.; Spoof, L.; Meriluoto, J. Cyanobacteria and microcystins in Koka reservoir (Ethiopia). *Environ. Sci. Pollut. Res. Int.* **2018**, *25*, 26861–26873. [[CrossRef](#)]
159. Bouhaddada, R.; Nelieu, S.; Nasri, H.; Delarue, G.; Bouaicha, N. High diversity of microcystins in a *Microcystis* bloom from an Algerian lake. *Environ. Pollut.* **2016**, *216*, 836–844. [[CrossRef](#)]
160. Kimambo, O.N.; Gumbo, J.R.; Chikoore, H. The occurrence of cyanobacteria blooms in freshwater ecosystems and their link with hydro-meteorological and environmental variations in Tanzania. *Heliyon* **2019**, *5*, e01312. [[CrossRef](#)]
161. Simiyu, B.M.; Oduor, S.O.; Rohrlack, T.; Sitoki, L.; Kurmayer, R. Microcystin content in phytoplankton and in small fish from eutrophic Nyanza Gulf, Lake Victoria, Kenya. *Toxins* **2018**, *10*, 275. [[CrossRef](#)]
162. Ballot, A.; Sandvik, M.; Rundberget, T.; Botha, C.; Miles, C. Diversity of cyanobacteria and cyanotoxins in Hartbeespoort Dam, South Africa. *Mar. Freshw. Res.* **2014**, *65*, 175–189. [[CrossRef](#)]
163. Herry, S.; Fathallii, A.; Jerhani, A.; Bouaicha, N. Seasonal occurrence and toxicity of *Microcystis* spp. and *Oscillatoria tenuis* in the Lebna Dam, Tunisia. *Water Res.* **2008**, *42*, 1263–1273. [[CrossRef](#)]
164. Ndebele, M.R.; Magadza, C.H.D. The occurrence of microcystin-LR in Lake Chivero, Zimbabwe. *Lakes Reserv. Res. Manag.* **2006**, *11*, 57–62. [[CrossRef](#)]
165. Douma, M.; Ouahid, Y.; Loudiki, M.; Del Campo, F.F.; Oudra, B. The first detection of potentially toxic *Microcystis* strains in two Middle Atlas Mountains natural lakes (Morocco). *Environ. Monit. Assess.* **2017**, *189*, 39. [[CrossRef](#)]
166. Okello, W.; Kurmayer, R. Seasonal development of cyanobacteria and microcystin production in Ugandan freshwater lakes. *Lakes Reserv. Res. Manag.* **2011**, *16*, 123–135. [[CrossRef](#)]
167. Mohamed, Z.A.; Deyab, M.A.; Abou-Dobara, M.I.; El-Sayed, A.K.; El-Raghi, W.M. Occurrence of cyanobacteria and microcystin toxins in raw and treated waters of the Nile River, Egypt: Implication for water treatment and human health. *Environ. Sci. Pollut. Res.* **2015**, *22*, 11716–11727. [[CrossRef](#)]
168. Chia, A.M.; Abolude, D.S.; Ladan, Z.; Akanbi, O.; Kalaboms, A. The presence of microcystins in aquatic ecosystems in Northern Nigeria: Zaria as a case study. *Res. J. Environ. Toxicol.* **2009**, *3*, 170–178. [[CrossRef](#)]
169. Addico, G.N.D. Toxic Cyanobacteria, Cyanotoxins and Drinking Water Production in Ghana: Implications to Human Health. Ph.D. Thesis, University of Hull, Hull, UK, 2008.
170. Mohamed, Z.A.; Bakr, A. Concentrations of cylindrospermopsin toxin in water and tilapia fish of tropical fishponds in Egypt, and assessing their potential risk to human health. *Environ. Sci. Pollut. Res. Int.* **2018**, *25*, 36287–36297. [[CrossRef](#)]
171. Odokuma, L.O.; Isirima, J.C. Distribution of cyanotoxins in aquatic environments in the Niger Delta. *Afr. J. Biotechnol.* **2007**, *6*, 2375–2385.
172. Harding, W.R.; Rowe, N.; Wessels, J.C.; Beattie, K.A.; Codd, G.A. Death of a dog attributed to the cyanobacterial (blue-green algal) hepatotoxin nodularin in South Africa. *J. S. Afr. Vet. Assoc.* **1995**, *66*, 256–259.
173. Magonono, M.; Oberholster, P.J.; Addmore, S.; Stanley, M.; Gumbo, J.R. The presence of toxic and non-toxic Cyanobacteria in the sediments of the Limpopo River Basin: Implications for human health. *Toxins* **2018**, *10*, 269. [[CrossRef](#)]
174. Naouli, J.; Abouabdellah, R.; Bennouna, A.; Laissaoui, A.; Swarzenski, P.W.; Ait Bouh, H.; Mesfioui, A.; Benbrahim, M.S.; Dechraoui Bottein, M.Y. Using the radioligand-receptor binding assay for paralytic shellfish toxins: A case study on shellfish from Morocco. *J. Environ. Radioact.* **2018**, *192*, 485–490. [[CrossRef](#)]
175. Mussagy, A.; Annadotter, H.; Cronberg, G. An experimental study of toxin production in *Arthospira fusiformis* (Cyanophyceae) isolated from African waters. *Toxicon* **2006**, *48*, 1027–1034. [[CrossRef](#)]

176. Esterhuizen, M.; Downing, T.G. Beta-N-methylamino-L-alanine (BMAA) in novel South African cyanobacterial isolates. *Ecotoxicol. Environ. Saf.* **2008**, *71*, 309–313. [[CrossRef](#)]
177. Tatters, A.O.; Howard, M.D.; Nagoda, C.; Busse, L.; Gellene, A.G.; Caron, D.A. Multiple stressors at the land-sea interface: Cyanotoxins at the land-sea interface in the Southern California Bight. *Toxins* **2017**, *9*. [[CrossRef](#)]
178. Wiltsie, D.; Schnetzer, A.; Green, J.; Vander Borgh, M.; Fensin, E. Algal blooms and Cyanotoxins in Jordan Lake, North Carolina. *Toxins* **2018**, *10*, 92. [[CrossRef](#)]
179. Foss, A.J.; Butt, J.; Aubel, M.T. Benthic periphyton from Pennsylvania, USA is a source for both hepatotoxins (microcystins/nodularin) and neurotoxins (anatoxin-a/homoanatoxin-a). *Toxicon* **2018**, *150*, 13–16. [[CrossRef](#)]
180. Beversdorf, L.J.; Weirich, C.A.; Bartlett, S.L.; Miller, T.R. Variable cyanobacterial toxin and metabolite profiles across six eutrophic lakes of differing physiochemical characteristics. *Toxins* **2017**, *9*, 62. [[CrossRef](#)]
181. Banack, S.A.; Metcalf, J.S.; Bradley, W.G.; Cox, P.A. Detection of cyanobacterial neurotoxin beta-N-methylamino-L-alanine within shellfish in the diet of an ALS patient in Florida. *Toxicon* **2014**, *90*, 167–173. [[CrossRef](#)]
182. Zastepa, A.; Taranu, Z.E.; Kimpe, L.E.; Blais, J.M.; Gregory-Eaves, I.; Zurawell, R.W.; Pick, F.R. Reconstructing a long-term record of microcystins from the analysis of lake sediments. *Sci. Total Environ.* **2017**, *579*, 893–901. [[CrossRef](#)]
183. Roy-Lachapelle, A.; Solliec, M.; Sauvé, S. Determination of BMAA and three alkaloid cyanotoxins in lake water using dansyl chloride derivatization and high-resolution mass spectrometry. *Anal. Bioanal. Chem.* **2015**, *407*, 5487–5501. [[CrossRef](#)]
184. Lajeunesse, A.; Segura, P.A.; Gelinas, M.; Hudon, C.; Thomas, K.; Quilliam, M.A.; Gagnon, C. Detection and confirmation of saxitoxin analogues in freshwater benthic Lyngbya wollei algae collected in the St. Lawrence River (Canada) by liquid chromatography-tandem mass spectrometry. *J. Chromatogr. A* **2012**, *1219*, 93–103. [[CrossRef](#)]
185. Zamora-Barrios, C.A.; Nandini, S.; Sarma, S.S.S. Bioaccumulation of microcystins in seston, zooplankton and fish: A case study in Lake Zumpango, Mexico. *Environ. Pollut.* **2019**, *249*, 267–276. [[CrossRef](#)]
186. Berry, J.P.; Jaja-Chimedza, A.; Davalos-Lind, L.; Lind, O. Apparent bioaccumulation of cylindrospermopsin and paralytic shellfish toxins by finfish in Lake Catemaco (Veracruz, Mexico). *Food Addit. Contam. Part A* **2012**, *29*, 314–321. [[CrossRef](#)]
187. Vasconcelos, V.; Martins, A.; Vale, M.; Antunes, A.; Azevedo, J.; Welker, M.; Lopez, O.; Montejano, G. First report on the occurrence of microcystins in planktonic cyanobacteria from Central Mexico. *Toxicon* **2010**, *56*, 425–431. [[CrossRef](#)]
188. Trout-Haney, J.V.; Wood, Z.T.; Cottingham, K.L. Presence of the cyanotoxin microcystin in Arctic Lakes of Southwestern Greenland. *Toxins* **2016**, *8*, 256. [[CrossRef](#)]
189. Kleinteich, J.; Wood, S.A.; Puddick, J.; Schleheck, D.; Kupper, F.C.; Dietrich, D. Potent toxins in Arctic environments—Presence of saxitoxins and an unusual microcystin variant in Arctic freshwater ecosystems. *Chem.-Biol. Interact.* **2013**, *206*, 423–431. [[CrossRef](#)]
190. Jokela, J.; Heinila, L.M.P.; Shishido, T.K.; Wahlsten, M.; Fewer, D.P.; Fiore, M.F.; Wang, H.; Haapaniemi, E.; Permi, P.; Sivonen, K. Production of high amounts of hepatotoxin nodularin and new protease inhibitors pseudospumigins by the Brazilian benthic *Nostoc* sp. CENA543. *Front. Microbiol.* **2017**, *8*, 1963. [[CrossRef](#)]
191. Walter, J.M.; Lopes, F.A.C.; Lopes-Ferreira, M.; Vidal, L.M.; Leomil, L.; Melo, F.; de Azevedo, G.S.; Oliveira, R.M.S.; Medeiros, A.J.; Melo, A.S.O.; et al. Occurrence of harmful cyanobacteria in drinking water from a severely drought-impacted semi-arid region. *Front. Microbiol.* **2018**, *9*, 176. [[CrossRef](#)]
192. Casali, S.P.; Dos Santos, A.C.A.; de Falco, P.B.; Calijuri, M.D.C. Influence of environmental variables on saxitoxin yields by *Cylindrospermopsis raciborskii* in a mesotrophic subtropical reservoir. *J. Water Health* **2017**, *15*, 509–518. [[CrossRef](#)]
193. Boopathi, T.; Ki, J.S. Impact of environmental factors on the regulation of cyanotoxin production. *Toxins* **2014**, *6*, 1951–1978. [[CrossRef](#)]
194. Barón-Sola, Á.; Ouahid, Y.; del Campo, F.F. Detection of potentially producing cylindrospermopsin and microcystin strains in mixed populations of cyanobacteria by simultaneous amplification of cylindrospermopsin and microcystin gene regions. *Ecotoxicol. Env. Saf.* **2012**, *75*, 102–108. [[CrossRef](#)]

195. Nimptsch, J.; Woelfl, S.; Osorio Ruiz, S.; Valenzuela, J.; Moreira, C.; Ramos, V.; Castelo Branco, R.; Leao, P.; Vasconcelos, V. First record of toxins associated with cyanobacterial blooms in oligotrophic North Patagonian lakes of Chile—A genomic approach. *Int. Rev. Hydrobiol.* **2016**, *101*, 57–168. [[CrossRef](#)]
196. Vidal, L.; Kruk, C. *Cylindrospermopsis raciborskii* (Cyanobacteria) extends its distribution to Latitude 34°53' S: Taxonomical and ecological features in Uruguayan eutrophic lakes. *Pan-Am. J. Aquat. Sci.* **2008**, *3*, 142–151.
197. Johnson, H.E.; King, S.R.; Banack, S.A.; Webster, C.; Callanaupa, W.J.; Cox, P.A. Cyanobacteria (*Nostoc* commune) used as a dietary item in the Peruvian highlands produce the neurotoxic amino acid BMAA. *J. Ethnopharmacol.* **2008**, *118*, 159–165. [[CrossRef](#)]
198. Puddick, J.; Prinsep, M.R.; Wood, S.A.; Cary, S.C.; Hamilton, D.P.; Holland, P.T. Further characterization of glycine-containing microcystins from the McMurdo dry Valleys of Antarctica. *Toxins* **2015**, *7*, 493–515. [[CrossRef](#)]
199. Hitzfeld, B.C.; Lampert, C.S.; Spaeth, N.; Mountfort, D.; Kaspar, H.; Dietrich, D.R. Toxin production in cyanobacterial mats from ponds on the McMurdo ice shelf, Antarctica. *Toxicon* **2000**, *38*, 1731–1748. [[CrossRef](#)]
200. Kleinteich, J.; Hildebrand, F.; AWood, S.; Cirés, S.; Agha, R.; Quesada, A.; Pearce, D.; Convey, P.; Küpper, F.; Dietrich, D. Diversity of toxin and non-toxin containing cyanobacterial mats of meltwater ponds on the Antarctic Peninsula: A pyrosequencing approach. *Antarct. Sci.* **2014**, *26*, 521–532. [[CrossRef](#)]
201. Genuario, D.B.; Correa, D.M.; Komarek, J.; Fiore, M.F. Characterization of freshwater benthic biofilm-forming *Hydrocoryne* (Cyanobacteria) isolates from Antarctica. *J. Phycol.* **2013**, *49*, 1142–1153. [[CrossRef](#)]
202. Ceglowska, M.; Torunska-Sitarz, A.; Kowalewska, G.; Mazur-Marzec, H. Specific chemical and genetic markers revealed a thousands-year presence of toxic *nodularia spumigena* in the Baltic Sea. *Mar. Drugs* **2018**, *16*, 116. [[CrossRef](#)]
203. Sivonen, K.; Namikoshi, M.; Evans, W.R.; Carmichael, W.W.; Sun, F.; Rouhiainen, L.; Luukkainen, R.; Rinehart, K.L. Isolation and characterization of a variety of microcystins from seven strains of the cyanobacterial genus *Anabaena*. *Appl. Environ. Microbiol.* **1992**, *58*, 2495–2500. [[CrossRef](#)]
204. Chernova, E.; Sidelev, S.; Russikh, I.; Voyakina, E.; Zhakovskaya, Z. First observation of microcystin- and anatoxin-a-producing cyanobacteria in the easternmost part of the Gulf of Finland (the Baltic Sea). *Toxicon* **2019**, *157*, 18–24. [[CrossRef](#)]
205. Kurmayer, R.; Christiansen, G.; Fastner, J.; Borner, T. Abundance of active and inactive microcystin genotypes in populations of the toxic cyanobacterium *Planktothrix* spp. *Environ. Microbiol.* **2004**, *6*, 831–841. [[CrossRef](#)]
206. Gkelis, S.; Lanaras, T.; Sivonen, K. Cyanobacterial toxic and bioactive peptides in freshwater bodies of Greece: Concentrations, occurrence patterns, and implications for human health. *Mar. Drugs* **2015**, *13*, 6319–6335. [[CrossRef](#)]
207. Grabowska, M.; Kobos, J.; Toruńska-Sitarz, A.; Mazur-Marzec, H. Non-ribosomal peptides produced by *Planktothrix* agardhii from Siemianówka Dam Reservoir SDR (northeast Poland). *Arch. Microbiol.* **2014**, *196*, 697–707. [[CrossRef](#)]
208. Touzet, N.; McCarthy, D.; Gill, A.; Fleming, G.T.A. Comparative summer dynamics of surface cyanobacterial communities in two connected lakes from the west of Ireland. *Sci. Total Environ.* **2016**, *553*, 416–428. [[CrossRef](#)]
209. Lopes, V.R.; Ramos, V.; Martins, A.; Sousa, M.; Welker, M.; Antunes, A.; Vasconcelos, V.M. Phylogenetic, chemical and morphological diversity of cyanobacteria from Portuguese temperate estuaries. *Mar. Environ. Res.* **2012**, *73*, 7–16. [[CrossRef](#)]
210. Farkas, O.; Gyemant, G.; Hajdu, G.; Gonda, S.; Parizsa, P.; Horgos, T.; Mosolygo, A.; Vasas, G. Variability of microcystins and its synthetase gene cluster in *Microcystis* and *Planktothrix* waterblooms in shallow lakes of Hungary. *Acta Biol. Hung.* **2014**, *65*, 227–239. [[CrossRef](#)]
211. Handeland, K.; Ostensvik, O. Microcystin poisoning in roe deer (*Capreolus capreolus*). *Toxicon* **2010**, *56*, 1076–1078. [[CrossRef](#)]
212. Faassen, E.J.; Lurling, M. Occurrence of the microcystins MC-LW and MC-LF in Dutch surface waters and their contribution to total microcystin toxicity. *Mar. Drugs* **2013**, *11*, 2643–2654. [[CrossRef](#)]
213. Jancula, D.; Strakova, L.; Sadilek, J.; Marsalek, B.; Babica, P. Survey of cyanobacterial toxins in Czech water reservoirs—the first observation of neurotoxic saxitoxins. *Env. Sci. Pollut. Res. Int.* **2014**, *21*, 8006–8015. [[CrossRef](#)]
214. Rücker, J.; Wiedner, C.; Zippel, P. Factors controlling the dominance of *Planktothrix* agardhii and *Limnothrix* redekei in eutrophic shallow lakes. *Hydrobiologia* **1997**, *342*, 107–115. [[CrossRef](#)]

215. Pekar, H.; Westerberg, E.; Bruno, O.; Laane, A.; Persson, K.M.; Sundstrom, L.F.; Thim, A.M. Fast, rugged and sensitive ultra high pressure liquid chromatography tandem mass spectrometry method for analysis of cyanotoxins in raw water and drinking water—First findings of anatoxins, cylindrospermopsins and microcystin variants in Swedish source waters and infiltration ponds. *J. Chromatogr. A* **2016**, *1429*, 265–276. [[CrossRef](#)]
216. Aboal, M. Are we underestimating benthic cyanotoxins? Extensive sampling results from Spain. *Toxins* **2017**, *9*, 385. [[CrossRef](#)]
217. Krzton, W.; Pudas, K.; Pociecha, A.; Strzesak, M.; Kosiba, J.; Walusiak, E.; Szarek-Gwiazda, E.; Wilk-Wozniak, E. Microcystins affect zooplankton biodiversity in oxbow lakes. *Environ. Toxicol. Chem.* **2017**, *36*, 165–174. [[CrossRef](#)]
218. Hunter, P.D.; Tyler, A.N.; Carvalho, L.; Codd, G.A.; Maberly, S.C. Hyperspectral remote sensing of cyanobacterial pigments as indicators for cell populations and toxins in eutrophic lakes. *Remote Sens. Environ.* **2010**, *114*, 2705–2718. [[CrossRef](#)]
219. Teneva, I.; Mladenov, R.; Belkinova, D.; Dimitrova-Dyulgerova, I.; Dzhambazov, B. Phytoplankton community of the drinking water supply reservoir Borovitsa (South Bulgaria) with an emphasis on cyanotoxins and water quality. *Cent. Eur. J. Biol.* **2010**, *5*, 231–239. [[CrossRef](#)]
220. Osswald, J.; Rellan, S.; Gago-Martinez, A.; Vasconcelos, V. Production of anatoxin-a by cyanobacterial strains isolated from Portuguese fresh water systems. *Ecotoxicology* **2009**, *18*, 1110–1115. [[CrossRef](#)]
221. Sabart, M.; Misson, B.; Descroix, A.; Duffaud, E.; Combourieu, B.; Salencon, M.J.; Latour, D. The importance of small colonies in sustaining *Microcystis* population exposed to mixing conditions: An exploration through colony size, genotypic composition and toxic potential. *Environ. Microbiol. Rep.* **2013**, *5*, 747–756. [[CrossRef](#)]
222. Pitois, F.; Fastner, J.; Pagotto, C.; Dechesne, M. Multi-toxin occurrences in ten french water resource reservoirs. *Toxins* **2018**, *10*, 283. [[CrossRef](#)]
223. Drobac, D.; Tokodi, N.; Lujic, J.; Marinovic, Z.; Subakov-Simic, G.; Dulic, T.; Vazic, T.; Nybom, S.; Meriluoto, J.; Codd, G.A.; et al. Cyanobacteria and cyanotoxins in fishponds and their effects on fish tissue. *Harmful Algae* **2016**, *55*, 66–76. [[CrossRef](#)]
224. Belykh, O.I.; Gladkikh, A.S.; Sorokovikova, E.G.; Tikhonova, I.V.; Butina, T.V. Identification of toxic Cyanobacteria in Lake Baikal. *Dokl. Biochem. Biophys.* **2015**, *463*, 220–224. [[CrossRef](#)]
225. Sulcius, S.; Pilkaityte, R.; Mazur-Marzec, H.; Kasperoviciene, J.; Ezhova, E.; Blaszczyk, A.; Paskauskas, R. Increased risk of exposure to microcystins in the scum of the filamentous cyanobacterium *Aphanizomenon flos-aquae* accumulated on the western shoreline of the Curonian Lagoon. *Mar. Pollut. Bull.* **2015**, *99*, 264–270. [[CrossRef](#)]
226. Sidelev, S.I.; Golokolenova, T.B.; Chernova, E.N.; Russkikh, I.V. Analysis of phytoplankton in Tsimlyansk Reservoir (RUSSIA) for the presence of cyanobacterial hepato- and neurotoxins. *Mikrobiologiya* **2015**, *84*, 732–742. [[CrossRef](#)]
227. Barda, I.; Kankaanaa, H.; Purina, I.; Balode, M.; Sjovall, O.; Meriluoto, J. Bioaccumulation of hepatotoxins—A considerable risk in the Latvian environment. *Environ. Pollut.* **2015**, *196*, 313–320. [[CrossRef](#)]
228. Greer, B.; McNamee, S.E.; Boots, B.; Cimarelli, L.; Guillebault, D.; Helmi, K.; Marcheggiani, S.; Panaiotov, S.; Breitenbach, U.; Akcaalan, R.; et al. A validated UPLC-MS/MS method for the surveillance of ten aquatic biotoxins in European brackish and freshwater systems. *Harmful Algae* **2016**, *55*, 31–40. [[CrossRef](#)]
229. Preussel, K.; Stuken, A.; Wiedner, C.; Chorus, I.; Fastner, J. First report on cylindrospermopsin producing *Aphanizomenon flos-aquae* (Cyanobacteria) isolated from two German lakes. *Toxicon* **2006**, *47*, 156–162. [[CrossRef](#)]
230. Brient, L.; Lengronne, M.; Bormans, M.; Fastner, J. First occurrence of cylindrospermopsin in freshwater in France. *Environ. Toxicol.* **2009**, *24*, 415–420. [[CrossRef](#)]
231. Messineo, V.; Melchiorre, S.; Di Corcia, A.; Gallo, P.; Bruno, M. Seasonal succession of *Cylindrospermopsis raciborskii* and *Aphanizomenon ovalisporum* blooms with cylindrospermopsin occurrence in the volcanic Lake Albano, Central Italy. *Environ. Toxicol.* **2010**, *25*, 18–27. [[CrossRef](#)]
232. Bláhová, L.; Babica, P.; Adamovský, O.; Kohoutek, J.; Maršálek, B.; Bláha, L. Analyses of cyanobacterial toxins (microcystins, cylindrospermopsin) in the reservoirs of the Czech Republic and evaluation of health risks. *Environ. Chem. Lett.* **2008**, *6*, 223–227. [[CrossRef](#)]

233. Cirés, S.; Wörmer, L.; Ballot, A.; Agha, R.; Wiedner, C.; Velázquez, D.; Casero, M.C.; Quesada, A. Phylogeography of cylindrospermopsin and paralytic shellfish toxin-producing Nostocales cyanobacteria from Mediterranean Europe (Spain). *Appl. Environ. Microbiol.* **2014**, *80*, 1359–1370. [[CrossRef](#)]
234. Cires, S.; Ballot, A. A review of the phylogeny, ecology and toxin production of bloom-forming *Aphanizomenon* spp. and related species within the Nostocales (cyanobacteria). *Harmful Algae* **2016**, *54*, 21–43. [[CrossRef](#)]
235. Gkelis, S.; Zaoutsos, N. Cyanotoxin occurrence and potentially toxin producing cyanobacteria in freshwaters of Greece: A multi-disciplinary approach. *Toxicon* **2014**, *78*, 1–9. [[CrossRef](#)]
236. Spoof, L.; Berg, K.A.; Rapala, J.; Lahti, K.; Lepistö, L.; Metcalf, J.S.; Codd, G.A.; Meriluoto, J. First observation of cylindrospermopsin in *Anabaena lapponica* isolated from the boreal environment (Finland). *Environ. Toxicol.* **2006**, *21*, 552–560. [[CrossRef](#)]
237. Kokociński, M.; Mankiewicz-Boczek, J.; Jurczak, T.; Spoof, L.; Meriluoto, J.; Rejmonczyk, E.; Hautala, H.; Vehniäinen, M.; Pawełczyk, J.; Soininen, J. *Aphanizomenon gracile* (Nostocales), a cylindrospermopsin-producing cyanobacterium in Polish lakes. *Environ. Sci. Pollut. Res.* **2013**, *20*, 5243–5264. [[CrossRef](#)]
238. Đorđević, N.; Simić, S.; Cirić, A. First identification of the cylindrospermopsin (CYN) producing by cyanobacterium *Cylindrospermopsis raciborskii* (Woloszynska) Seenayya & Subba Raju in Serbia. *Fresenius Environ. Bull.* **2015**, *24*, 3736–3742.
239. Moreira, C.; Mendes, R.; Azevedo, J.; Vasconcelos, V.; Antunes, A. First occurrence of cylindrospermopsin in Portugal: A contribution to its continuous global dispersal. *Toxicon* **2017**, *130*, 87–90. [[CrossRef](#)]
240. Antal, O.; Karisztl-Gácsi, M.; Farkas, A.; Kovács, A.; Ács, A.; Törő, N.; Kiss, G.; Saker, M.L.; Győri, J.; Bánfalvi, G.; et al. Screening the toxic potential of *Cylindrospermopsis raciborskii* strains isolated from Lake Balaton, Hungary. *Toxicon* **2011**, *57*, 831–840. [[CrossRef](#)]
241. James, K.J.; Sherlock, I.R.; Stack, M.A. Anatoxin-a in Irish freshwater and cyanobacteria, determined using a new fluorimetric liquid chromatographic method. *Toxicon* **1997**, *35*, 963–971. [[CrossRef](#)]
242. Shams, S.; Capelli, C.; Cerasino, L.; Ballot, A.; Dietrich, D.R.; Sivonen, K.; Salmaso, N. Anatoxin-a producing *Tychonema* (Cyanobacteria) in European waterbodies. *Water Res.* **2015**, *69*, 68–79. [[CrossRef](#)]
243. Echenique-Subiabre, I.; Tenon, M.; Humbert, J.F.; Quiblier, C. Spatial and Temporal Variability in the Development and Potential Toxicity of *Phormidium* Biofilms in the Tarn River, France. *Toxins* **2018**, *10*, 418. [[CrossRef](#)]
244. Faassen, E.J.; Harkema, L.; Begeman, L.; Lurling, M. First report of (homo)anatoxin-a and dog neurotoxicosis after ingestion of benthic cyanobacteria in The Netherlands. *Toxicon* **2012**, *60*, 378–384. [[CrossRef](#)]
245. Onodera, H.; Oshima, Y.; Henriksen, P.; Yasumoto, T. Confirmation of anatoxin-a(s), in the cyanobacterium *Anabaena lemmermannii*, as the cause of bird kills in Danish Lakes. *Toxicon* **1997**, *35*, 1645–1648. [[CrossRef](#)]
246. Sierolsawska, A.; Rymuszka, A.; Kalinowska, R.; Skowronski, T.; Bownik, A.; Pawlik-Skowronska, B. Toxicity of cyanobacterial bloom in the eutrophic dam reservoir (Southeast Poland). *Environ. Toxicol. Chem.* **2010**, *29*, 556–560. [[CrossRef](#)]
247. Rivetti, C.; Gomez-Canela, C.; Lacorte, S.; Diez, S.; Lazaro, W.L.; Barata, C. Identification of compounds bound to suspended solids causing sub-lethal toxic effects in *Daphnia magna*. A field study on re-suspended particles during river floods in Ebro River. *Aquat. Toxicol.* **2015**, *161*, 41–50. [[CrossRef](#)]
248. Edwards, C.; Beattie, K.A.; Scrimgeour, C.M.; Codd, G.A. Identification of anatoxin-A in benthic cyanobacteria (blue-green algae) and in associated dog poisonings at Loch Insh, Scotland. *Toxicon* **1992**, *30*, 1165–1175. [[CrossRef](#)]
249. Ballot, A.; Fastner, J.; Lentz, M.; Wiedner, C. First report of anatoxin-a-producing cyanobacterium *Aphanizomenon issatschenkoi* in northeastern Germany. *Toxicon* **2010**, *56*, 964–971. [[CrossRef](#)]
250. Ballot, A.; Scherer, P.I.; Wood, S.A. Variability in the anatoxin gene clusters of *Cuspidothrix issatschenkoi* from Germany, New Zealand, China and Japan. *PLoS ONE* **2018**, *13*, e0200774. [[CrossRef](#)]
251. Kaas, H.; Henriksen, P. Saxitoxins (PSP toxins) in Danish Lakes. *Water Res.* **2000**, *34*, 2089–2097. [[CrossRef](#)]
252. Ledreux, A.; Thomazeau, S.; Catherine, A.; Duval, C.; Yéprémian, C.; Marie, A.; Bernard, C. Evidence for saxitoxins production by the cyanobacterium *Aphanizomenon gracile* in a French recreational water body. *Harmful Algae* **2010**, *10*, 88–97. [[CrossRef](#)]
253. Ballot, A.; Fastner, J.; Wiedner, C. Paralytic shellfish poisoning toxin-producing cyanobacterium *Aphanizomenon gracile* in northeast Germany. *Appl. Environ. Microbiol.* **2010**, *76*, 1173–1180. [[CrossRef](#)]

254. Pomati, F.; Sacchi, S.; Rossetti, C.; Giovannardi, S.; Onodera, H.; Oshima, Y.; Neilan, B.A. The freshwater cyanobacterium *Planktothrix* sp. fp1: Molecular identification and detection of paralytic shellfish poisoning toxins. *J. Phycol.* **2000**, *36*, 553–562. [[CrossRef](#)]
255. Rapala, J.; Robertson, A.; Negri, A.P.; Berg, K.A.; Tuomi, P.; Lyra, C.; Erkomaa, K.; Lahti, K.; Hoppu, K.; Lepisto, L. First report of saxitoxin in Finnish lakes and possible associated effects on human health. *Environ. Toxicol.* **2005**, *20*, 331–340. [[CrossRef](#)]
256. Usleber, E.; Donald, M.; Straka, M.; Märtlbauer, E. Comparison of enzyme immunoassay and mouse bioassay for determining paralytic shellfish poisoning toxins in shellfish. *Food Addit. Contam.* **1997**, *14*, 193–198. [[CrossRef](#)]
257. Savela, H.; Spoof, L.; Perälä, N.; Vehniäinen, M.; Mankiewicz-Boczek, J.; Jurczak, T.; Kokociński, M.; Meriluoto, J. First report of cyanobacterial paralytic shellfish toxin biosynthesis genes and paralytic shellfish toxin production in Polish freshwater lakes. *Adv. Oceanogr. Limnol.* **2017**, *8*. [[CrossRef](#)]
258. Ballot, A.; Cerasino, L.; Hostyeva, V.; Cires, S. Variability in the sxt gene clusters of PSP toxin producing *Aphanizomenon gracile* strains from Norway, Spain, Germany and North America. *PLoS ONE* **2016**, *11*, e0167552. [[CrossRef](#)]
259. Beltran, E.; Ibanez, M.; Sancho, J.V.; Hernandez, F. Determination of six microcystins and nodularin in surface and drinking waters by on-line solid phase extraction-ultra high pressure liquid chromatography tandem mass spectrometry. *J. Chromatogr. A* **2012**, *1266*, 61–68. [[CrossRef](#)]
260. Beattie, K.A.; Kaya, K.; Codd, G.A. The cyanobacterium *Nodularia* PCC 7804, of freshwater origin, produces [L-Har<sup>2</sup>]nodularin. *Phytochemistry* **2000**, *54*, 57–61. [[CrossRef](#)]
261. Eiser, A.R. Why does Finland have the highest dementia mortality rate? Environmental factors may be generalizable. *Brain Res.* **2017**, *1671*, 14–17. [[CrossRef](#)]
262. Rosen, J.; Westerberg, E.; Hellénas, K.E.; Salomonsson, M.L. A new method for analysis of underivatized free β-methylamino-alanine: Validation and method comparison. *Toxicon* **2016**, *121*, 105–108. [[CrossRef](#)]
263. Lage, S.; Costa, P.R.; Moita, T.; Eriksson, J.; Rasmussen, U.; Rydberg, S.J. BMAA in shellfish from two Portuguese transitional water bodies suggests the marine dinoflagellate *Gymnodinium catenatum* as a potential BMAA source. *Aquat. Toxicol.* **2014**, *152*, 131–138. [[CrossRef](#)]
264. Luglie, A.; Giacobbe, M.G.; Riccardi, E.; Bruno, M.; Pigozzi, S.; Mariani, M.A.; Satta, C.T.; Stacca, D.; Bazzoni, A.M.; Caddeo, T.; et al. Paralytic shellfish toxins and cyanotoxins in the Mediterranean: New data from Sardinia and Sicily (Italy). *Microorganisms* **2017**, *5*, 72. [[CrossRef](#)]
265. Rzymski, P.; Poniedziałek, B.; Mankiewicz-Boczek, J.; Faassen, E.; Jurczak, T.; Gała, I.; Ballot, A.; Lurling, M.; Kokociński, M. Polyphasic toxicological screening of *Cylindrospermopsis raciborskii* and *Aphanizomenon gracile* isolated in Poland. *Algal Res.* **2017**, *24*, 72–80. [[CrossRef](#)]
266. Metcalf, J.; Anne Banack, S.; Lindsay, J.; FMorrison, L.; Alan Cox, P.; ACodd, G. Co-occurrence of β-N-methylamino-L-alanine, a neurotoxic amino acid with other cyanobacterial toxins in British waterbodies, 1990–2004. *Environ. Microbiol.* **2008**, *10*, 702–708. [[CrossRef](#)]
267. Masseret, E.; Banack, S.; Boumédiène, F.; Abadie, E.; Brent, L.; Pernet, F.; Juntas-Morales, R.; Pageot, N.; Metcalf, J.; Cox, P.; et al. Dietary BMAA exposure in an amyotrophic lateral sclerosis cluster from southern France. *PLoS ONE* **2013**, *8*, e83406. [[CrossRef](#)]
268. Violi, J.P.; Mitrovic, S.M.; Colville, A.; Main, B.J.; Rodgers, K.J. Prevalence of β-methylamino-L-alanine (BMAA) and its isomers in freshwater cyanobacteria isolated from eastern Australia. *Ecotoxicol. Environ. Saf.* **2019**, *172*, 72–81. [[CrossRef](#)]
269. Gaget, V.; Humpage, A.R.; Huang, Q.; Monis, P.; Brookes, J.D. Benthic cyanobacteria: A source of cylindrospermopsin and microcystin in Australian drinking water reservoirs. *Water Res.* **2017**, *124*, 454–464. [[CrossRef](#)]
270. Wood, S.A.; Maier, M.Y.; Puddick, J.; Pochon, X.; Zaiko, A.; Dietrich, D.R.; Hamilton, D.P. Trophic state and geographic gradients influence planktonic cyanobacterial diversity and distribution in New Zealand lakes. *FEMS Microbiol. Ecol.* **2017**, *93*. [[CrossRef](#)]
271. Wood, S.A.; Stirling, D.J. First identification of the cylindrospermopsin-producing cyanobacterium *Cylindrospermopsis raciborskii* in New Zealand. *N. Z. J. Mar. Freshw. Res.* **2003**, *37*, 821–828. [[CrossRef](#)]
272. Smith, F.M.; Wood, S.A.; van Ginkel, R.; Broady, P.A.; Gaw, S. First report of saxitoxin production by a species of the freshwater benthic cyanobacterium, *Scytonema Agardh*. *Toxicon* **2011**, *57*, 566–573. [[CrossRef](#)]

273. Vogiazi, V.; de la Cruz, A.; Mishra, S.; Shanov, V.; Heineman, W.R.; Dionysiou, D.D. A comprehensive review: Development of electrochemical biosensors for detection of cyanotoxins in freshwater. *ACS Sens.* **2019**, *4*, 1151–1173. [[CrossRef](#)]
274. Straser, A.; Filipic, M.; Zegura, B. Cylindrospermopsin induced transcriptional responses in human hepatoma HepG2 cells. *Toxicol. In Vitro* **2013**, *27*, 1809–1819. [[CrossRef](#)]
275. Vasas, G.; Gaspar, A.; Suranyi, G.; Batta, G.; Gyemant, G.; M, M.H.; Mathe, C.; Grigorszky, I.; Molnar, E.; Borbely, G. Capillary electrophoretic assay and purification of cylindrospermopsin, a cyanobacterial toxin from *Aphanizomenon ovalisporum*, by plant test (blue-green Sinapis test). *Anal. Biochem.* **2002**, *302*, 95–103. [[CrossRef](#)]
276. McElhinney, J.; Lawton, L.A.; Leifert, C. Investigations into the inhibitory effects of microcystins on plant growth, and the toxicity of plant tissues following exposure. *Toxicon* **2001**, *39*, 1411–1420. [[CrossRef](#)]
277. Wu, Z.; He, D.; Cui, B.; Jin, Z. Ultrasensitive detection of microcystin-LR with gold immunochromatographic assay assisted by a molecular imprinting technique. *Food Chem.* **2019**, *283*, 517–521. [[CrossRef](#)]
278. Zhang, H.; Luo, J.; Beloglazova, N.; Yang, S.; De Saeger, S.; Mari, G.M.; Zhang, S.; Shen, J.; Wang, Z.; Yu, X. Portable multiplex immunochromatographic assay for quantitation of two typical algae toxins based on dual-color fluorescence microspheres. *J. Agric. Food Chem.* **2019**, *67*, 6041–6047. [[CrossRef](#)]
279. Cunha, I.; Biltés, R.; Sales, M.; Vasconcelos, V. Aptamer-based biosensors to detect aquatic phycotoxins and cyanotoxins. *Sensors* **2018**, *18*, 2367. [[CrossRef](#)]
280. Wharton, R.E.; Feyereisen, M.C.; Gonzalez, A.L.; Abbott, N.L.; Hamelin, E.I.; Johnson, R.C. Quantification of saxitoxin in human blood by ELISA. *Toxicon* **2017**, *133*, 110–115. [[CrossRef](#)]
281. Hicks, J.M. Fluorescence immunoassay. *Hum. Pathol.* **1984**, *15*, 112–116. [[CrossRef](#)]
282. Rossi, A.M.; Taylor, C.W. Analysis of protein-ligand interactions by fluorescence polarization. *Nat. Protoc.* **2011**, *6*, 365–387. [[CrossRef](#)] [[PubMed](#)]
283. Smith, D.S.; Eremin, S.A. Fluorescence polarization immunoassays and related methods for simple, high-throughput screening of small molecules. *Anal. Bioanal. Chem.* **2008**, *391*, 1499–1507. [[CrossRef](#)] [[PubMed](#)]
284. Bahadir, E.B.; Sezgintürk, M.K. Lateral flow assays: Principles, designs and labels. *TrAC Trends Anal. Chem.* **2016**, *82*, 286–306. [[CrossRef](#)]
285. Xing, C.; Liu, L.; Song, S.; Feng, M.; Kuang, H.; Xu, C. Ultrasensitive immunochromatographic assay for the simultaneous detection of five chemicals in drinking water. *Biosens. Bioelectron.* **2015**, *66*, 445–453. [[CrossRef](#)] [[PubMed](#)]
286. Singh, S.; Srivastava, A.; Oh, H.M.; Ahn, C.Y.; Choi, G.G.; Asthana, R.K. Recent trends in development of biosensors for detection of microcystin. *Toxicon* **2012**, *60*, 878–894. [[CrossRef](#)]
287. Bickman, S.R.; Campbell, K.; Elliott, C.; Murphy, C.; O’Kennedy, R.; Papst, P.; Lochhead, M.J. An innovative portable biosensor system for the rapid detection of freshwater cyanobacterial algal bloom toxins. *Environ. Sci. Technol.* **2018**, *52*, 11691–11698. [[CrossRef](#)]
288. Metcalf, J.; Hiskia, A.; Kaloudis, T. Protein Phosphatase Inhibition Assays. *Handb. Cyanobacterial Monit. Cyanotoxin Anal.* **2017**, *26*, 267–271. [[CrossRef](#)]
289. Pereira, S.R.; Vasconcelos, V.M.; Antunes, A. The phosphoprotein phosphatase family of Ser/Thr phosphatases as principal targets of naturally occurring toxins. *Crit. Rev. Toxicol.* **2011**, *41*, 83–110. [[CrossRef](#)]
290. Hereszty, T.; Nicholson, B.C. Determination of cyanobacterial hepatotoxins directly in water using a protein phosphatase inhibition assay. *Water Res.* **2001**, *35*, 3049–3056. [[CrossRef](#)]
291. Ruberu, S.R.; Langlois, G.W.; Masuda, M.; Kittredge, C.; Perera, S.K.; Kudela, R.M. Receptor binding assay for the detection of paralytic shellfish poisoning toxins: Comparison to the mouse bioassay and applicability under regulatory use. *Food Addit. Contam. Part A* **2018**, *35*, 144–158. [[CrossRef](#)]
292. Araoz, R.; Vilarino, N.; Botana, L.M.; Molgo, J. Ligand-binding assays for cyanobacterial neurotoxins targeting cholinergic receptors. *Anal. Bioanal. Chem.* **2010**, *397*, 1695–1704. [[CrossRef](#)] [[PubMed](#)]
293. Moreira, C.; Ramos, V.; Azevedo, J.; Vasconcelos, V. Methods to detect cyanobacteria and their toxins in the environment. *Appl. Microbiol. Biotechnol.* **2014**, *98*, 8073–8082. [[CrossRef](#)]
294. Sun, L.W.; Jiang, W.J.; Zhang, J.Y.; Wang, W.Q.; Du, Y.; Sato, H.; Kawachi, M.; Yu, R. Identification and detection sensitivity of *Microcystis aeruginosa* from mixed and field samples using MALDI-TOF MS. *Environ. Monit. Assess.* **2018**, *190*, 712. [[CrossRef](#)] [[PubMed](#)]

295. Nicolau, R.; Leloup, M.; Lachassagne, D.; Pinault, E.; Feuillade-Cathalifaud, G. Matrix-assisted laser desorption/ionization time-of-flight mass spectrometry (MALDI-TOF-MS) coupled to XAD fractionation: Method to algal organic matter characterization. *Talanta* **2015**, *136*, 102–107. [CrossRef] [PubMed]
296. He, D.; Wu, Z.; Cui, B.; Jin, Z. A novel SERS-based aptasensor for ultrasensitive sensing of microcystin-LR. *Food Chem.* **2019**, *278*, 197–202. [CrossRef] [PubMed]
297. Pacheco, A.B.; Guedes, I.A.; Azevedo, S.M. Is qPCR a reliable indicator of cyanotoxin risk in freshwater? *Toxins* **2016**, *8*, 172. [CrossRef] [PubMed]
298. Humbert, J.F.; Quiblier, C.; Gugger, M. Molecular approaches for monitoring potentially toxic marine and freshwater phytoplankton species. *Anal. Bioanal. Chem.* **2010**, *397*, 1723–1732. [CrossRef] [PubMed]
299. Pearson, L.A.; Neilan, B.A. The molecular genetics of cyanobacterial toxicity as a basis for monitoring water quality and public health risk. *Curr. Opin. Biotechnol.* **2008**, *19*, 281–288. [CrossRef] [PubMed]
300. Srivastava, A.; Singh, S.; Ahn, C.Y.; Oh, H.M.; Asthana, R.K. Monitoring approaches for a toxic cyanobacterial bloom. *Environ. Sci. Technol.* **2013**, *47*, 8999–9013. [CrossRef]
301. Humbert, J.F. Molecular tools for the detection of toxicogenic cyanobacteria in natural ecosystems. *Handb. Cyanobacterial Monit. Cyanotoxin Anal.* **2017**, *28*, 280–283. [CrossRef]
302. McNamee, S.E.; Elliott, C.T.; Greer, B.; Lochhead, M.; Campbell, K. Development of a planar waveguide microarray for the monitoring and early detection of five harmful algal toxins in water and cultures. *Environ. Sci. Technol.* **2014**, *48*, 13340–13349. [CrossRef] [PubMed]
303. Fraga, M.; Vilariño, N.; Louzao, M.C.; Rodríguez, L.P.; Alfonso, A.; Campbell, K.; Elliott, C.T.; Taylor, P.; Ramos, V.; Vasconcelos, V.; et al. Multi-detection method for five common microalgal toxins based on the use of microspheres coupled to a flow-cytometry system. *Anal. Chim. Acta* **2014**, *850*, 57–64. [CrossRef]
304. Zeller, P.; Mejean, A.; Biegala, I.; Contremoulin, V.; Ploux, O. Fluorescence in situ hybridization of *Microcystis* strains producing microcystin using specific mRNA probes. *Lett. Appl. Microbiol.* **2016**, *63*, 376–383. [CrossRef]
305. Johnson, B.N.; Mutharasan, R. A cantilever biosensor-based assay for toxin-producing cyanobacteria *Microcystis aeruginosa* using 16S rRNA. *Environ. Sci. Technol.* **2013**, *47*, 12333–12341. [CrossRef] [PubMed]
306. Gan, N.; Huang, Q.; Zheng, L.; Song, L. Quantitative assessment of toxic and nontoxic *Microcystis* colonies in natural environments using fluorescence in situ hybridization and flow cytometry. *Sci. China Life Sci.* **2010**, *53*, 973–980. [CrossRef] [PubMed]
307. Pernthaler, A.; Pernthaler, J.; Amann, R. Fluorescence in situ hybridization and catalyzed reporter deposition for the identification of marine bacteria. *Appl. Environ. Microbiol.* **2002**, *68*, 3094–3101. [CrossRef]
308. Schonhuber, W.; Fuchs, B.; Juretschko, S.; Amann, R. Improved sensitivity of whole-cell hybridization by the combination of horseradish peroxidase-labeled oligonucleotides and tyramide signal amplification. *Appl. Env. Microbiol.* **1997**, *63*, 3268–3273. [CrossRef]
309. Ramm, J.; Lupu, A.; Hadas, O.; Ballot, A.; Rucker, J.; Wiedner, C.; Sukenik, A. A CARD-FISH protocol for the identification and enumeration of cyanobacterial akinetes in lake sediments. *FEMS Microbiol. Ecol.* **2012**, *82*, 23–36. [CrossRef]
310. Riou, V.; Périot, M.; Biegala, I.C. Specificity re-evaluation of oligonucleotide probes for the detection of marine picoplankton by tyramide signal amplification-fluorescent in situ hybridization. *Front. Microbiol.* **2017**, *8*, 854. [CrossRef]
311. Meriluoto, J.; Metcalf, J.; Codd, G.A. Selection of analytical methodology for cyanotoxin analysis. *Biochemistry* **2017**, *32*, 309–312. [CrossRef]
312. Khait, I.; Obolski, U.; Yovel, Y.; Hadany, L. Sound perception in plants. *Semin. Cell Dev. Biol.* **2019**, *92*, 134–138. [CrossRef] [PubMed]
313. Zheng, C.; Zeng, H.; Lin, H.; Wang, J.; Feng, X.; Qiu, Z.; Chen, J.A.; Luo, J.; Luo, Y.; Huang, Y.; et al. Serum microcystin levels positively linked with risk of hepatocellular carcinoma: A case-control study in southwest China. *Hepatology* **2017**, *66*, 1519–1528. [CrossRef] [PubMed]

