

Article

# Tetanus Toxin Synthesis is Under the Control of A Complex Network of Regulatory Genes in *Clostridium tetani*

Diana Chapeton-Montes <sup>1</sup>, Lucile Plourde <sup>2</sup>, Cecile Deneve <sup>1</sup>, Dominique Garnier <sup>2</sup>, Fabien Barbirato <sup>2</sup>, Vincent Colombié <sup>2</sup>, Sandy Demay <sup>2</sup>, Georges Haustant <sup>1</sup>, Olivier Gorgette <sup>3</sup>, Christine Schmitt <sup>3</sup>, Catherine Thouvenot <sup>3</sup>, Holger Brüggemann <sup>4</sup> and Michel R. Popoff <sup>1,\*</sup>

<sup>1</sup> Bactéries anaérobies et Toxines, Institut Pasteur, 75724 Paris, France; dianajoanne@gmail.com (D.C.-M.); cecile.deneve-larrazet@u-psud.fr (C.D.); georges.haustant@pasteur.fr (G.H.)

<sup>2</sup> Sanofi-Pasteur, 69280 Marcy l'Etoile, France; Lucile.Plourde@sanofi.com (L.P.); Dominique.Garnier@sanofi.com (D.G.); Fabien.Barbirato@sanofi.com (F.B.); Vincent.Colombie@sanofi.com (V.C.); Sandy.Demay@sanofi.com (S.D.)

<sup>3</sup> Unité Technologie et Service Bioimagerie Ultrastructurale, Institut Pasteur, 75724 Paris, France; olivier.gorgette@pasteur.fr (O.G.); christine.schmitt@pasteur.fr (C.S.); catherine.thouvenot@pasteur.fr (C.T.)

<sup>4</sup> Department of Biomedicine, Aarhus University, 8000 Aarhus, Denmark; brueggemann@biomed.au.dk

\* Correspondence: popoff2m@gmail.com

Received: 20 April 2020; Accepted: 8 May 2020; Published: 15 May 2020



**Abstract:** *Clostridium tetani* produces a potent neurotoxin, the tetanus toxin (TeNT), which is responsible for an often-fatal neurological disease (tetanus) characterized by spastic paralysis. Prevention is efficiently acquired by vaccination with the TeNT toxoid, which is obtained by *C. tetani* fermentation and subsequent purification and chemical inactivation. *C. tetani* synthesizes TeNT in a regulated manner. Indeed, the TeNT gene (*tent*) is mainly expressed in the late exponential and early stationary growth phases. The gene *tetR* (tetanus regulatory gene), located immediately upstream of *tent*, encodes an alternative sigma factor which was previously identified as a positive regulator of *tent*. In addition, the genome of *C. tetani* encodes more than 127 putative regulators, including 30 two-component systems (TCSs). Here, we investigated the impact of 12 regulators on TeNT synthesis which were selected based on their homology with related regulatory elements involved in toxin production in other clostridial species. Among nine TCSs tested, three of them impact TeNT production, including two positive regulators that indirectly stimulate *tent* and *tetR* transcription. One negative regulator was identified that interacts with both *tent* and *tetR* promoters. Two other TCSs showed a moderate effect: one binds to the *tent* promoter and weakly increases the extracellular TeNT level, and another one has a weak inverse effect. In addition, CodY (control of *dcIA* (decoyinine induced operon) Y) but not Spo0A (sporulation stage 0) or the DNA repair protein Mfd (mutation frequency decline) positively controls TeNT synthesis by interacting with the *tent* promoter. Moreover, we found that inorganic phosphate and carbonate are among the environmental factors that control TeNT production. Our data show that TeNT synthesis is under the control of a complex network of regulators that are largely distinct from those involved in the control of toxin production in *Clostridium botulinum* or *Clostridium difficile*.

**Keywords:** *Clostridium tetani*; *Clostridium botulinum*; tetanus toxin; two-component system; gene transcription

**Key Contribution:** Tetanus toxin synthesis is under the control of a complex network of regulation linked to the metabolism in *Clostridium tetani* including at least two positive and one negative two-component system regulators and the master regulator of metabolism CodY. Inorganic phosphate and carbonate are environmental factors involved in the control of tetanus toxin synthesis.

## 1. Introduction

*Clostridium tetani* is an environmental Gram-positive, spore-forming and anaerobic rod-shaped bacterium which synthesizes a potent neurotoxin, the tetanus toxin (TeNT) [1]. *C. tetani* spores can enter an organism through an open wound. Spores germinate and *C. tetani* grows in anaerobic conditions in necrotic tissues. TeNT is synthesized at the end of the exponential growth phase and is released in the surrounding tissues. TeNT specifically recognizes nidogens at the neuromuscular junctions [2], and enters motorneurons. TeNT is retrogradely transported to the central nervous system. Then, TeNT enters inhibitory interneurons and blocks the release of neurotransmitters (glycine, GABA (gamma-aminobutyric acid)) upon proteolytic cleavage of the SNARE (soluble N-ethylmaleimide-sensitive factor attachment protein receptor) protein VAMP2 (vesicle associated membrane protein) [3,4].

Vaccination based on the TeNT toxoid is a very efficient prevention measure against tetanus. It is noteworthy that in addition to the use as a single-antigen vaccine for specific prevention of tetanus, the TeNT toxoid is combined with other vaccine antigens for protection against other infectious diseases [5]. Industrial TeNT production is obtained by fermentation of *C. tetani* clinical isolates. TeNT is then extracted from culture supernatants and inactivated with formalin. *C. tetani* fermentation is a critical and complex step in vaccine production as it is performed in a rich growth medium under a strictly controlled environment. The essential medium components such as amino acids that result in a high TeNT yield are partially identified. Thus, the toxin yield varies with different media and charges [6].

The TeNT gene (*tent*) is located on a large plasmid in *C. tetani* [7,8]. The first complete genome sequence of a toxigenic *C. tetani* strain (E88) was determined in 2003 [9]. The genome contains a chromosome of approximately 2.8 Mb and a TeNT-encoding plasmid of 74 kb. The genomes of additional *C. tetani* strains have been sequenced and showed that the plasmid-encoded *tent* is highly conserved [10–12]. A conserved gene (*tetR*, tetanus toxin regulatory gene) just upstream of *tent* encodes for an alternative sigma factor which positively regulates the transcription of *tent* [13]. TetR is homologous to BotR (botulinum toxin regulator) which controls the synthesis of the botulinum neurotoxin (BoNT) in *Clostridium botulinum* A, B, C and D [14]. TetR and BotR belong with *Clostridium difficile* TcdR (or TxeR) (*Clostridium difficile* toxin regulator) and *Clostridium perfringens* UviA (UV-inducible gene A) to a sub-group of the sigma 70 family of RNA polymerase sigma factors which control clostridial toxin gene syntheses [15,16]. However, the regulatory network governing TeNT synthesis is still poorly understood. A better knowledge of the regulatory mechanisms of TeNT synthesis in *C. tetani* and determination of the environmental factors controlling this regulation are required for improving toxin production.

To successfully respond to changes in different environmental conditions and to regulate virulence, many bacteria use a complex regulatory network involving diverse types of molecules, including RNA, DNA, proteins and metabolites. Among these are global regulators like two-component systems (TCS) and CodY to sense the relevant environmental signals [17,18]. TCSs are ubiquitous among bacteria. They consist of a membrane-bound or cytosolic sensor histidine kinase (SHK) that senses a stimulus and its cytoplasmic cognate response regulator (RR) that mediates the cellular response. Following a specific stimulus, the SHK autophosphorylates at a conserved histidine residue. The phosphoryl group is then transferred from the histidine to a conserved aspartate residue in the RR, which upon phosphorylation is able to control the expression of its target genes [18].

Depending on the availability of nutrients, bacteria have to adjust their gene expression. The global regulator CodY has been shown to be an important regulatory link between metabolism and virulence factor synthesis in many low G + C Gram-positive bacteria. CodY displays enhanced affinity for its DNA target when bound to GTP and/or branched-chain amino acids [19]. In *C. botulinum* ATCC 3502, CodY has been shown to positively regulate *bont* expression, and to bind to the promoter of the *ntnh-bontA* operon in a GTP-dependent manner [20]. In addition, the sporulation regulator Spo0A has been found to positively regulate BoNT synthesis in *C. botulinum* type E, which is part of the group II

of *C. botulinum* and lacks BotR. Spo0A binds to the promoter of the *bont/E* operon and enhances its transcription [21]. Spo0A is expressed in the exponential growth phase of *C. botulinum* and initiates the cascade of alternative sigma factors involved in sporulation [22]. However, Spo0A targets several hundred other genes and has pleiotropic effects [23]. Moreover, the mutation frequency decline (*mfd*) gene, which encodes a transcription-coupled repair factor, also shows pleiotropic effects: it increases the expression of both Toxin A and Toxin B-encoding genes in *C. difficile* [24]. These genes *codY*, *spo0A* and *mfd* are conserved in the *C. tetani* genome.

Genomic analysis of the *C. tetani* strain E88 reveals the presence of numerous putative regulatory genes, some of them being homologous to regulatory genes in other clostridia or bacteria. The aim of this work was to determine the role of 12 putative regulatory systems in TeNT synthesis, nine two-component systems as well as the global regulators CodY, Spo0A and Mfd.

## 2. Results

### 2.1. Genomic Analysis of Regulatory Genes in *C. tetani*

Genome sequencing of strain E88 showed that the large *tent*-containing plasmid also harbors regulatory elements: *tetR* located immediately upstream of *tent*, a two-component system (TCS) (CTC\_RS13805/CTC\_RS13810, previously named *ctp21/ctp22*) homologous to systems in other Gram-positive bacteria including *C. botulinum* and *Bacillus* sp., and three additional putative sigma factors [9]. In addition, refined genomic analysis showed that the chromosome of strain E88 possesses 30 TCSs based on the conserved motifs of histidine kinase as part of the SHK and a receiver domain Rec as part of the adjacent RR. Most of the 30 TCSs belong to the OmpR (outer membrane protein regulator) family (19 TCSs), two to (lytic gene regulator/alginate biosynthesis regulator) LytR/AlgR, two to (nitrate reductase regulator) NarL, two to (factor for inversion stimulation) Fis, one to (L-arabinose gene regulator) AraC, one to (purine catabolic gene regulator) PucR, one to LysR (lysine synthesis regulator) and one to XRE ((prophage *Bacillus subtilis*) PBSX repressor) families (Table 1). All 30 TCSs except one share homology with putative TCSs in other clostridia species. Among them, 19 TCSs are homologous to related genes in *C. botulinum* strain Hall (identity level  $\geq 45\%$ ) (Table 1). It was previously shown that five TCSs in strain Hall positively regulate botulinum neurotoxin (BoNT) synthesis [25]. Two of them are homologous to *C. tetani* TCSs, i.e., CLC0661/CLC0663 from *C. botulinum* with 65% protein identity to CTC\_RS02080/CTC\_RS02085 of *C. tetani* and CLC0410/CLC0411 shows 68% protein identity to CTC\_RS10030/CTC\_RS10035 (Table 1). Another study identified one TCS (CB00786/CB00787) in *C. botulinum* strain ATCC3502, that negatively regulates *bont* gene expression [26]. This TCS shares significant identity with a *C. tetani* TCS (CTC\_RS07310/CTC\_RS07315).

The genome of *C. tetani* strain CN655 [10] shares the same TCS genes as strain E88. Since we have already characterized the *tetR* in strain CN655, a strain which is more easily transformable compared to other *C. tetani* strains under the applied conditions, the strain CN655 was selected for further genetic investigations including the analysis of nine TCS genes, *mfd*, *spo0A* and *codY*.

Res, respiration; Bac, bacitracin; Vir, virulence; Spa, subtilin gene; Van, vancomycin; Gtc, gramicidin transcription; Arc, aerobic respiration control; Pho, phosphate; Feu, ferric uptake; Dcu, dicarboxylate uptake; Sin, sporulation inhibition; Eut, ethanolamine utilization; Ato, acetoacetyl-coenzyme A transferase.

**Table 1.** Two component system (TCS) genes encoded in the genome of *C. tetani* E88 and their homology with regulatory genes in other clostridia. The recombinant RNA antisense plasmids targeting TCS genes are indicated in column 1. Genetic environment indicates functional genes in close proximity of TCS in *C. tetani* E88.

Recombinant Antisense mRNA Plasmid	Gene Bank Accession Number		Other Gene Name	Role	Family (RR)	Genetic Environment	Homology	Homologs (RR) in Other Clostridia (Protein Identity > 60%)	Homolog TCS in <i>C. botulinum</i> strain Hall			Homolog TCS in <i>C. botulinum</i> Strain ATCC 3502	
	Old Locus Tag	Locus Tag							Homolog System (RR/SHK)	Protein Identity (RR)	Regulation of Botulism Neurotoxin	Homolog System (RR/SHK)	Protein Identity/Positive
<b>chromosomal localization</b>													
	CTC_00189 CTC_00191	CTC_RS00820 CTC_RS00825		RR SHK	OmpR	Periplasmic endopeptidase	ResD, respiration control	<i>C. botulinum</i> , <i>C. cochlearium</i> , <i>C. tetanomorphum</i> , <i>C. scatologenes</i> , <i>C. drakei</i> , <i>C. magnum</i> , <i>C. sporogenes</i> , <i>C. tyrobutyricum</i> , <i>C. acetireducens</i> , <i>C. ljungdahlii</i> , <i>C. novyi</i> , <i>C. pasteurianum</i> ,	CLC_3521/CLC_3520	81%	None	CBO3543 CBO3542	81/87 67/73
	CTC_00392 CTC_00393	CTC_RS01985 CTC_RS01990		RR SHK	OmpR	ABC Transporter	BacS/BacR regulation of resistance to bacitracin	<i>C. botulinum</i> , <i>C. tetanomorphum</i> , <i>C. sporogenes</i> , <i>C. lundense</i> , <i>C. cavendishii</i> , <i>C. acetobutylicum</i> , <i>C. pasteurianum</i> , <i>C. ljungdahlii</i>	CLC_0331/CLC_0332	90%	None	CBO0272 CBO0273	90/92 86/89
p1308	CTC_00411 CTC_00412	CTC_RS02080 CTC_RS02085	phoP phoR	RR SHK	OmpR	hydroxylamine reductase	VirI/VirJ regulation of toxin synthesis	<i>C. botulinum</i> , <i>C. perfringens</i> , <i>C. carboxidivorans</i> , <i>C. pasteurianum</i> , <i>C. lundense</i> , <i>C. novyi</i> , <i>C. septicum</i>	CLC_0661/CLC_0663	65%	Positive	CBO0607 CBO0608	64/72 46/55
	CTC_00455 CTC_00456	CTC_RS02330 CTC_RS14405		RR SHK		phosphomannomutase/ phosphoglucomutase	putative rRNA methylase	<i>C. tetanomorphum</i> , <i>C. ljungdahlii</i> , <i>C. homopropionicum</i>	CLC_0413	55%	unknown	CBO0355 no	54/59
	CTC_00597 CTC_00598	CTC_RS02990 CTC_RS02995		RR SHK	OmpR	Glycerine-dehydrogenase	BacR regulation of resistance to bacitracin	<i>C. lundense</i> , <i>C. tetanomorphum</i> , <i>C. ljungdahlii</i> , <i>C. argetinense</i> , <i>C. drakei</i>	CLC_2212/CLC_2211	45%	None	several no	45/66
	CTC_00628 CTC_00629	CTC_RS03125 CTC_RS03130	spaR spaK	RR SHK	OmpR	ABC Transporter SpaEFG: export subtilin	SpaR/SpaK regulation of resistance to subtilin	<i>C. botulinum</i> , <i>C. novyi</i> , <i>C. beijerinckii</i> , <i>C. saccharobutylicum</i> , <i>C. butyricum</i> , <i>C. sporogenes</i>	CLC_1615/CLC_1616	65%	unknown	CBO1585 CBO1586	65/69 37/43
	CTC_00805 CTC_00806	CTC_RS04010 CTC_RS04015		RR SHK	OmpR	Protein transport	VanR/VanS regulation of resistance to vancomycin	<i>C. botulinum</i> , <i>C. cochlearium</i> , <i>C. tetanomorphum</i> , <i>C. lundense</i> , <i>C. carboxidivorans</i> , <i>C. sporogenes</i> , <i>C. pasteurianum</i>	CLC_0423/CLC_0424	76%	None	CBO0365 CBO0366	78/82 44/50
	CTC_00848 CTC_00849	CTC_RS04235 CTC_RS04240		RR SHK	OmpR	Heat shock protein	GtcS/GtcR regulation of antibiotic synthesis (gramicidin)	<i>C. lundense</i> , <i>C. tetanomorphum</i> , <i>C. amylolyticum</i> , <i>C. perfringens</i>	CLC_1640/CLC_1639	48%	None	several no	47/55
	CTC_00872 CTC_00873	CTC_RS04365 CTC_RS04370		RR SHK	OmpR	Efflux-ATPase Copper	Arc, regulation of aerobic/anaerobic respiration	<i>C. botulinum</i> , <i>C. tetanomorphum</i> , <i>C. lundense</i> , <i>C. sporogenes</i> , <i>C. pasteurianum</i> , <i>C. beijerinckii</i> , <i>C. ljungdahlii</i>	CLC_1088/CLC_1089	68%	None	CBO1035 CBO1036	66/71 55/63
	CTC_00924 CTC_00925	CTC_RS04645 CTC_RS04650		SHK RR	PucR	Zn-dependent protease	regulation of purine catabolism, methyl-accepting chemotaxis protein		No			CBO1773 no	43/50
p1309	CTC_00934 CTC_00935 CTC_00949	CTC_RS04705 CTC_RS04710 CTC_RS04780		SHK RR SHK	AraC	amihydrolase; Pyruvate formate lyase	LytS, autolysis regulation	<i>C. botulinum</i> B str. Eklund 17B, <i>C. novyi</i>	CLC_1627	41%	unknown	no no no	
p1310	CTC_00950	CTC_RS04785	virR	RR	LytR/AlgR	transcriptional regulator, merR family; putative methyl-accepting chemotaxis protein	virulence regulation	<i>C. butyricum</i> , <i>C. septicum</i>	CLC_1105/CLC_1104	35%	None	CBO1053	35/54

Table 1. Cont.

Recombinant Antisense mRNA Plasmid	Gene Bank Accession Number		Other Gene Name	Role	Family (RR)	Genetic Environment	Homology	Homologs (RR) in Other Clostridia (Protein Identity > 60%)	Homolog TCS in <i>C. botulinum</i> strain Hall			Homolog TCS in <i>C. botulinum</i> Strain ATCC 3502		
	Old Locus Tag	Locus Tag							Homolog System (RR/SHK)	Protein Identity (RR)	Regulation of Botulism Neurotoxin	Homolog System (RR/SHK)	Protein Identity/Positive	
<b>chromosomal localization</b>														
p1311	CTC_01130 CTC_01131	CTC_RS05745 CTC_RS05750	phoR	RR SHK	OmpR	ABC Transporteur: phosphates	PhoP/PhoR, regulation of phosphate uptake	<i>C. lundense</i> , <i>C. carboxidivorans</i> , <i>C. ljungdahlii</i> , <i>C. botulinum</i> , <i>C. sporogenes</i> , <i>C. butyricum</i> , <i>C. acetobutylicum</i> , <i>C. butyricum</i> , <i>C. pasteurianum</i> , <i>C. neonatale</i> , <i>C. baratii</i>	CLC_2386/CLC_2385	73%	None	CBO2527 CBO2526	73/87 52/75	
p1312	CTC_01211 CTC_01212	CTC_RS06180 CTC_RS06185	tlpA	SHK RR	LysR	anaerobic sulfite reductase	LytR, autolysis regulation, methyl-accepting chemotaxis protein tlpA	<i>C. cochlearium</i>	CLC_3570	35%	unknown	CBO2828 several	36/61 35/57	
p1419	CTC_01420 CTC_01421	CTC_RS07310 CTC_RS07315	resE	SHK RR	OmpR	ABC Transporter	YycG/YycF, regulation of cell division	<i>C. lundense</i> , <i>C. cellulovorans</i> , <i>C. amylolyticum</i> , <i>C. botulinum</i> ,	CLC_0842/CLC_0843 CB00786/CB00787*	58%	None Negative	CBO0787 CBO0786	47/65 58/76	
	CTC_01481 CTC_01482	CTC_RS07700 CTC_RS07705		SHK RR	OmpR	Conserved proteins with transmembrane helices and 4Fe4S motif	FeuQ/FeuP, regulation of iron acquisition	<i>C. novyi</i> , <i>C. yurii</i>	CLC_3521/CLC_3520	43%	None	no several	43/60	
	CTC_01490 CTC_01491	CTC_RS07755 CTC_RS07760		RR SHK	OmpR	Heat shock protein HtpG (chaperone); Membrane protein	unknown	<i>C. lundense</i> , <i>C. tetanomorphum</i> , <i>C. kluyveri</i> , <i>C. pasteurianum</i> , <i>C. carboxidivorans</i> , <i>C. ljungdahlii</i> , <i>C. tyrobutyricum</i> , <i>C. acetobutylicum</i>	CLC_0423/CLC_0424	44%	None	several no	44/64	
	CTC_01523 CTC_01524	CTC_RS07895 CTC_RS07900	dpiB	RR SHK	NarL	Fumarate-reductase soluble flavoprotein	DcuR, regulation of fumarate anaerobic respiration through C4-dicarboxylates	<i>C. cochlearium</i>	CLC_0307/CLC_0306	48%	None	CBO0249 CBO0248	48/69 41/60	
	CTC_01804 CTC_01805	CTC_RS09305 CTC_RS09310		SHK RR	OmpR	ABC Transporter	BacS/BacR, AB-Bacitracine synthesis and regulation	<i>C. tetanomorphum</i> , <i>C. lundense</i> , <i>C. novyi</i>	CLC_2212/CLC_2211	56%	None	CBO2284 CBO2285	47/72 56/79	
	CTC_01818 CTC_01819	CTC_RS09380 CTC_RS09385	resE	SHK RR	OmpR	ABC Transporter; RNA polymerase sigma factor	unknown	<i>C. tetanomorphum</i> , <i>C. lundense</i> , <i>C. ljungdahlii</i> , <i>C. cavendishii</i>	CLC_0842/CLC_0843 CB00786/CB00787*	51%	None Negative	CBO0787 CBO0786	38/63 51/71	
	CTC_01848 CTC_01849	CTC_RS14320 CTC_RS09510	yesM	SHK RR	NarL/FixJ	Fumarate-reductase	unknown	<i>C. cochlearium</i>	CLC_2236/CLC_2235	26%	None	no no		
	CTC_01857 CTC_01858	CTC_RS09550 CTC_RS09555	sinR	SHK RR	XRE	Helicase, oleate hydratase	SinR regulation of entry in stationary phase, to nutrient depletion; Spo0A repressor	<i>C. botulinum</i> CDC_69094, <i>C. magnum</i> , <i>C. beijerinckii</i>	No			CBO0693 no	48/70	
	CTC_01905 CTC_01906	CTC_RS09790 CTC_RS09795		SHK RR	OmpR	ABC Transporter	BacR; VanR (synthèse et régulation AB)	<i>C. indolis</i> , <i>C. methoxybenzovorans</i>	CLC_0423/CLC_0424	45%	None	no CBO0365	45/66	
	CTC_01918 CTC_01919	CTC_RS09860 CTC_RS09865	resE	SHK RR	OmpR	ABC Transporter	BacR; VanR (synthèse et régulation AB)	<i>C. kluyveri</i> , <i>C. uliginosum</i> , <i>C. puniceum</i> , <i>C. scatologenes</i> , <i>C. saccharobutylicum</i> , <i>C. oryzae</i> , <i>C. ljungdahlii</i> , <i>C. lundense</i> , <i>C. drakey</i> , <i>C. botulinum</i> B2 331, <i>C. sporogenes</i> , <i>C. acetobutylicum</i> , <i>C. butyricum</i>	CLC_0842/CLC_0843 CB00786/CB00787*	42%	None Negative	CBO0787 CBO0786	32/53 42/60	

Table 1. Cont.

Recombinant Antisense mRNA Plasmid	Gene Bank Accession Number		Other Gene Name	Role	Family (RR)	Genetic Environment	Homology	Homologs (RR) in Other Clostridia (Protein Identity > 60%)	Homolog TCS in <i>C. botulinum</i> strain Hall			Homolog TCS in <i>C. botulinum</i> Strain ATCC 3502	
	Old Locus Tag	Locus Tag							Homolog System (RR/SHK)	Protein Identity (RR)	Regulation of Botulinism Neurotoxin	Homolog System (RR/SHK)	Protein Identity/Positive
<b>chromosomal localization</b>													
p1313	CTC_01951	CTC_RS10030	phoR	SHK	OmpR	Heavy metal translocating P-type aTPase	PhoP/PhoR regulation of phosphate uptake	<i>C. cochlearium</i> , <i>C. lundense</i> , <i>C. tetanomorphum</i> , <i>C. intestinale</i> , <i>C. amylolyticum</i> , <i>C. baratii</i> , <i>C. botulinum</i> , <i>C. chauvoei</i> , <i>C. sporogenes</i> , <i>C. perfringens</i> , <i>C. ljungdahlii</i>	CLC_0410/CLC_0411	68%	Positive	CBO0353	57/78
	CTC_01953	CTC_RS10035	phoP	RR								CBO0352	68/86
p1314	CTC_01978	CTC_RS10150		SHK	LytR/AlgR	carbon starvation protein A CstA	LytS/LytR autolysis regulation	<i>C. cochlearium</i> , <i>C. tetanomorphum</i> , <i>C. lundense</i>	CLC_3250/CLC_3251	55%	None	CBO3309	51/71
	CTC_01979	CTC_RS10155		RR								CBO3308	55/78
	CTC_02155	CTC_RS11115		SHK	OmpR	DNA mismatch repair protein hexA	VanS/VanR regulation of resistance to glycopeptides	<i>C. carboxidivorans</i> , <i>C. lundense</i> , <i>C. amylolyticum</i> , <i>C. botulinum</i> , <i>C. oryzae</i> , <i>C. pasteurianum</i> , <i>C. sporogenes</i> , <i>C. novyi</i> , <i>C. neonatale</i>	CLC_1640/CLC_1639	77%	None	CBO1612	48/68
	CTC_02156	CTC_RS11120		RR								CBO1613	77/86
	CTC_02178	CTC_RS11240		SHK	Fis	ethanolamine utilization protein EutA, EutP	EutS/EutR regulation of ethanolamine utilization	<i>C. argentinense</i> , <i>C. drakei</i> , <i>C. lundense</i> , <i>C. tetanomorphum</i>	No			no	no
	CTC_02179	CTC_RS11245		RR								no	no
	CTC_02322	CTC_RS11915		RR	Fis	sodium/glutamate symport carrier protein; V-Typ-ATPase-protein	AtoS/AtoC regulation of acetoacetate metabolism	<i>C. cochlearium</i> , <i>C. lundense</i> , <i>C. tetanomorphum</i>	CLC_1882	40%	unknown	several	46/63
	CTC_02323	CTC_RS11920		SHK								no	no
<b>plasmid localization</b>													
p1307	CTC_p22	CTC_RS13810		SHK	OmpR	ATP-binding protein	unknown	<i>C. lundense</i>	CLC_1431/CLC_1432	56%	None	CBO1395	39/60
	CTC_p21	CTC_RS13805		RR								CBO1394	56/74

RR, Response Regulator; SHK, Sensor Histidine Kinase, \* 100% homology with TCSs of *C. botulinum* ATCC3502.

## 2.2. Construction of TCS Anti-Sense Strains

Plasmids that were able to generate anti-sense mRNA from nine TCS genes, as well as from *mfd*, *spo0A* and *codY*, were constructed in analogy to the construction of pMRP306 which has previously been used in the investigation of *tetR* in *C. tetani* and regulatory genes in *C. botulinum* [13,14]. DNA segments for anti-sense mRNA production were designed, located in the RR gene of three TCSs (CTC\_RS04785, CTC\_RS10155, and CTC\_RS07315) and in the SHK gene in six other TCSs (CTC\_RS13810, CTC\_RS02085, CTC\_RS04705, CTC\_RS06180, CTC\_RS05750 and CTC\_RS10030) (Table 2).


**Table 2.** CN655 anti-sense strains, and primers used for the construction of recombinant plasmids generating antisense mRNA. The PCR products from CN655 genome DNA contain a 3'NcoI site and a 5'PstI site and were cloned into pMRP306, a derivative of pAT19 containing the promoter of the iota toxin gene, the cloning sites NcoI-PstI and the 3'part of the iota toxin gene [14]. The resultant antisense RNA plasmids were transformed into CN655 by electroporation. SHK: Sensor Histidine Kinase, RR: Response regulator.

Isogenic Antisense Strains	Target Gene	S/R	Primer	Nucleotide Sequence (5'→3)	Product Length (bp)
CN655/1307	CTC_p22	S	P2020-F P2021-R	CCGCTGCAGGATAATTTGGGAATGATTATTTTA GGCCATGGTTAACATATCGTCCATACTC	228
CN655/1308	CTC_00412	S	P2022-F P2023-R	CCGCTGCAGGAGGTGATTGAAAAATAG GGCCATGGTAAATCTAACATAGTAAATTTATAC	208
CN655/1310	CTC_00950	R	P2024-F P2025-R	CCGCTGCAGGGAGGGTTAAATTATGTATAATG GGCCATGGGCTACTTCTATACCATTATTTC	236
CN655/1309	CTC_00934	S	P2026-F P2027-R	CCGCTGCAGCAGGGGGTATTTTTGTGTTAAATAATAGG GGCCATGGCATTGGCATCGCAACATATGCG	236
CN655/1312	CTC_01211	S	P2028-F P2029-R	CCGCTGCAGGGGAGACAGTGGTGAAGTTGCCG GGCCATGGGGTTAAAAAATTTTCTTTTATATTTTC	223
CN655/1314	CTC_01979	R	P2030-F P2031-R	CCGCTGCAGGAGATGAATTTATGAACAAAAAT GGCCATGGGCTAATTCATGCCATTTTITAG	231
CN655/1311	CTC_01131	S	P2032-F P2033-R	CCGCTGCAGGGTGGTAAAATGAAAAAAG GGCCATGGCCTTATATCACTATCATTAA	245
CN655/1313	CTC_01951	S	P2034-F P2035-R	CCGCTGCAGGGAAGGTAGAAAATGAAAAGTATAAAG GGCCATGGCCACATTATCCATTATATTTTCTTC	243
CN655/1419	CTC_01421	R	P2291-F P2292-R	CCGCTGCAGGGGAGATTTTGTGAACAACATATT GGCCATGGTCTGATGCCCTTCTTATTTCTTAC	242
CN655/1418	CTC_01260	CodY	P2289-F P2290-R	CCGCTGCAGGAGGAGTTACAAATGTCATCATTATTA GGCCATGGACTACCTTGTCTCTTACTGTCTG	232
CN655/1472	CTC_00222	Spo0	P2361-F P2362-R	CCGCTGCAGGGAGGTATAAAAATATATGATA GGCCATGGTTATTACTCTTTAAAGGTGAA	225
CN655/1480	CTC_00194	<i>mfd</i>	P2359-F P2360-R	CCGCTGCAGGAGGTGAATTTTATTATGAGAT GGCCATGGAATATTTTTTGCTTCTATATCG	236

Although the growth kinetics of CN655 harboring the empty vector pAT18 (CN655/pAT18) was slightly lower than that of the wt CN655 during the exponential growth phase and beginning of the stationary phase, the total production of TeNT was slightly higher (Supplementary Materials Figure S1). Both CN655/pAT18 and CN655 wt showed similar growth and toxin production from 72 h to 144 h (Supplementary Materials Figure S1). No difference in bacterial size and microscopic morphology was observed between CN655 wt and CN655/pAT18.

Seven CN655 anti-sense strains targeting four TCSs (p1308 targeting CTC\_RS02085, p1310 targeting CTC\_RS04785, p1311 targeting CTC\_RS05750, p1419 targeting CTC\_RS07315), and three regulators (p1480 targeting *mfd*, p1472 targeting *spo0A*, and p1418 targeting *codY*), showed similar growth kinetics compared to the control strain CN655/pAT18 (Figure 1). However, growth of strains CN655/p1307 targeting CTC\_RS13810, CN655/p1309 targeting CTC\_RS04705, CN655/p1312 targeting CTC\_RS06180, CN655/p1313 targeting CTC\_RS10030, and CN655/p1314 targeting CTC\_RS10155 was more abundant than the control strain CN655/pAT18 in the early stationary phase (12–48 h), but the five anti-sense

strains reached a similar biomass as monitored by OD<sub>600nm</sub> than that of the other anti-sense strains at 72–144 h (Figure 1).



**Figure 1.** Growth kinetics of *C. tetani* strain CN655/pAT18 (empty vector) and CN655 antisense strains. (A) CN655/p1308, CN655/p1310, CN655/p1311, CN655/p1419, CN655/p1418, CN655/p1480 and CN655/p1472 showed a similar kinetics compared to CN655/pAT18 strain. (B) CN655/p1307, CN655/p1309, CN655/p1312, CN655/p1313 and CN655/p1314 displayed a more abundant growth than CN655/pAT18 in the early stationary phase (12–48 h). Data are mean values  $\pm$  SEM of at least three independent cultures.


### 2.3. TeNT Synthesis is Altered in Five TCS Anti-Sense Strains and in the *codY* Anti-Sense Strain

Extracellular TeNT concentrations as monitored by enzyme-linked immunosorbent assay (ELISA) in the culture supernatant were reduced in three CN655 anti-sense strains (CN655/p1307, CN655/p1314 and CN655/p1310) as well as in the *codY* anti-sense strain (CN655/p1418) compared to the control strain CN655/pAT18 (Figure 2A). CN655/p1307, which targets the TCS located on the large *tent*-containing *C. tetani* plasmid, and CN655/p1314 targeting the chromosomal CTC\_RS10155 TCS, showed a drastic decrease (65% to 75%) in the levels of secreted TeNT. In contrast, CN655/p1310 which targets CTC\_RS04785 exhibited a moderate decrease (about 25%) in secreted TeNT. The *codY* anti-sense strain (CN655/p1418) showed a significantly lower (50% to 65%) TeNT level in the culture supernatant within the first 56 h of culture and a less pronounced effect (20% to 25%) in the late growth phase (Figure 2A). The anti-sense strains CN655/p1308, CN655/p1309, CN655/p1312, CN655/p1313, CN655/p1480 and CN655/p1472 showed no significant difference in the production of extracellular TeNT compared to CN655/pAT18 (Supplementary Figure S1).


Total amounts of produced TeNT (i.e., TeNT in the culture supernatant and intracellular TeNT at the end of culture growth) were significantly decreased in CN655/p1307, CN655/p1314 and in CN655/p1418, whereas no significant difference was observed in CN655/p1310 (Figure 3A).

In contrast, the extracellular and total TeNT concentrations were significantly increased in CN655/p1419 targeting CTC\_RS07315 (Figures 2B and 3B). The anti-sense strain CN655/p1311 targeting CTC\_RS05750 showed an increase in extracellular TeNT within the first 48 h of culture, but the total TeNT amount at the end of culture was not significantly different from the control strain (Figure 2, Supplementary Materials Figure S2).

The total production of TeNT was not significantly different from that of the control strain CN655/pAT18 in CN655/p1308, CN655/p1309, CN655/p1312, CN655/p1313, CN655/p1480 and CN655/p1472 (Supplementary Materials Figure S3).



**Figure 2.** Extracellular tetanus toxin (TeNT) produced by *C. tetani* CN655/pAT18 (empty vector) and CN655 antisense strains. **(A)** Extracellular toxin was reduced in the culture supernatant of CN655/p1307, CN655/p1310, CN655/p1314 and CN655/p1418. **(B)** CN655/p1311 and CN655/p1419 showed elevated extracellular toxin kinetics compared to control strain CN655/pAT18. Statistical significance of differences between control and anti-sense strains is indicated with *p*-values (\*, *p* < 0.05; \*\*, *p* < 0.01; \*\*\*, *p* < 0.001). Data are mean values ± SEM of at least three independent cultures.




**Figure 3.** Total tetanus toxin (TeNT) produced by *C. tetani* CN655/pAT18 (empty vector) and CN655 antisense strains. (A) Total tetanus toxin production was reduced in CN655/p1307, CN655/p1314 and CN655/p1418. (B) CN655/p1419 showed elevated total toxin kinetics compared to control strain CN655/pAT18. Three independent experiments have been done. Statistical significance of differences between control strain and anti-sense strains is indicated with *p*-values (\*,  $p < 0.05$ ; \*\*,  $p < 0.01$ ).

#### 2.4. Five TCSs and *codY* Control Tent and/or Tetr at the Transcriptional Level

The transcriptional levels of *tent* and *tetR* were monitored by qRT-PCR at 8, 24, 32 and 48 h of culture, which corresponded to the exponential growth phase (8 h) and early stationary growth phase (12–48 h) (Figure 1). The limit of these analyses is based on the consideration that the copy numbers of the recombinant plasmids are similar in the recombinant strains. Among the TCSs investigated by RNA anti-sense, CN655/p1307 and CN655/p1314 showed a significant reduction of *tent* and *tetR* transcription during the 24–48 h growth phase and only a moderate effect at the 8 h exponential growth phase (Figure 4). A marked reduced transcription of *tetR* was observed in CN655/p1310 at the 8 h exponential and 24–32 h early stationary growth phase, whereas *tent* transcription was not significantly reduced (Figure 4). In contrast, the anti-sense strain CN655/p1311 showed an increase in *tent* and *tetR* transcription at the 8 h exponential growth phase and to a lower extent within the 24–48 h stationary growth phase (Figure 4). The anti-sense strain CN655/p1419 also exhibited an increased *tent* transcription level at 8 h exponential growth phase and at 48 h of culture, but a decreased transcription of *tetR* within the 24–32 h early stationary growth phase, albeit to a lower extent than in CN655/p1307

and CN655/p1314 (Figure 4). This suggests that CTC\_RS07315 (targeted in CN655/p1419) controls *tent* transcription independently of *tetR*.




**Figure 4.** Expression of (A) *tent* and (B) *tetR* in CN655/pAT 18 and CN655 antisense strains. (A) The expression of *tent* was repressed in CN655/p1307, CN655/p1310, CN655/p1314 and CN655/p1418 compared to the control strain CN655/pAT18. For strains CN655/p1311 and CN655/p1419, an increased *tent* expression was observed. (B) The expression of *tetR* was repressed in CN655/p1307, CN655/p1310, CN655/p1312, CN655/p1313, CN655/p1314, CN655/p1419 and CN655/p1418. Strain CN655/p1311 showed an increase in *tetR* expression compared to the control strain CN655/pAT18. Target gene expression was normalized to *rpoB* and *gyrA*. Three independent experiments have been done. Statistical significance of differences between control and the anti-sense strains is indicated with *p*-values (\*,  $p < 0.05$ ; \*\*,  $p < 0.01$ ).

The transcriptional levels of *tent* and *tetR* were not significantly different in CN655/p1308, and CN655/p1309 compared to the control strain CN655/pAT18 (Supplementary Materials Figure S4). Despite a slightly decreased transcription of *tetR* in CN655/p1312 and CN655/p1313 during the 24–32 h early stationary growth phase (Figure 4), no significant alteration of *tent* transcription was observed in these strains (Supplementary Materials Figure S3).

The anti-sense strain in which *codY* was targeted (CN655/p1418), showed a decrease in *tent* transcription within the 8–48 h culture and a moderate decrease in *tetR* transcription (Figure 4).

#### 2.5. The CTC\_RS07315 and CTC\_RS04785 Response Regulators as well as Cody Bind to the Tent Promoter

To identify the mode of action of the positive and negative regulators of *tent* gene transcription, we investigated their possible direct interaction with *tent* and/or *tetR* genes by testing the binding of the regulators to the promoters of *tent* and *tetR*. As shown in Figure 5, the CTC\_RS07315 (targeted in CN655/p1419) RR specifically bound to the promoters of both *tent* and *tetR* ( $P_{tent}$ ,  $P_{tetR}$ ). The RR CTC\_RS04785 (targeted in CN655/p1310) caused a shift in the mobility of  $P_{tent}$  but not in  $P_{tetR}$ , despite the significant decrease in *tetR* transcription. In contrast, no specific binding to  $P_{tent}$  or  $P_{tetR}$  was observed with the RRs of CTC\_RS10155, CTC\_RS13805 and CTC\_RS05745, albeit they influenced the TeNT synthesis and transcription of *tent*. As expected, no electrophoretic shift was observed with SHK such as CTC\_RS04710. In addition, CodY bound to  $P_{tent}$  but not to  $P_{tetR}$ . As a positive control, TetR, which has been found to positively regulate *tent*, was used [13].




**Figure 5.** Electrophoretic mobility shift assay (EMSA) showing regulatory protein binding to *tent* (A) and *tetR* (B) promoters. Biotin-labeled DNA probes corresponding to the promoter regions of *tent* ( $P_{tent}$ ) and *tetR* ( $P_{tetR}$ ) were incubated with 5  $\mu$ M of the recombinant proteins CTC\_RS13805, CTC\_RS10155, CTC\_RS07315, CTC\_RS04710, CTC\_RS05745, CTC\_RS04785, CodY and TetR. The specific binding of recombinant proteins to promoter probes resulted in an observable mobility shift when compared to the  $P_{tent}$  and  $P_{tetR}$  alone. Competition assays were performed with a 300-fold excess of unlabeled probe. Specificity of binding to  $P_{tent}$  was confirmed for CTC\_RS07315, CTC\_RS04785, CodY and TetR. CTC\_RS07315 was the only protein showing specific binding to  $P_{tetR}$ . Addition of recombinant proteins, unlabeled promoter probe as cold competitor and labeled promoter probes are indicated. Representative experiments out of three are shown.

## 2.6. Inorganic Phosphate ( $P_i$ ) Influences TeNT Production

Since two TCSs (CTC\_RS05745/CTC\_RS05750 and CTC\_RS10030/CTC\_RS10035) encoded in the *C. tetani* genome are putatively involved in the control of phosphate uptake and/or metabolism, as judged from their homologies with previously characterized TCSs, we checked the effect of inorganic phosphate ( $P_i$ ) in the culture medium on TeNT synthesis. Therefore, TGY (trypsinase/glucose/yeast extract) culture medium containing less than 2 mM  $P_i$  was supplemented with 10 to 60 mM  $P_i$ . *C. tetani* growth was similar in TGY supplemented with 10 and 20 mM  $P_i$  than that in the TGY control medium, but was slightly decreased when supplemented with 40 and 60 mM  $P_i$  (Figure 6). Extracellular and total TeNT production was increased in TGY containing 40 or 60 mM  $P_i$ , whereas no significant difference in TeNT production was observed in TGY supplemented with 10 or 20 mM  $P_i$ . The highest level of TeNT production occurred when *C. tetani* was grown in TGY medium with 40 mM  $P_i$  (at least two-fold more extracellular TeNT compared to TGY control medium) (Figure 6).

Expression of *tent* was increased in TGY medium containing 10 or 20 mM  $P_i$ , and to a higher extent when supplemented with 40 or 60 mM  $P_i$  (Figure 6D). The kinetics of increased *tent* expression in TGY supplemented with  $P_i$  is similar to that in TGY control medium, occurring at the end of the exponential and beginning of the stationary phase growth phases. Expression of *tetR* was similarly increased in cultures supplemented with  $P_i$ , but mainly within a shorter time duration between 24 and 32 h of culture (Figure 6E). This suggests that regulation of TeNT expression by inorganic phosphate is independent of *tetR*.




**Figure 6.** Effect of inorganic phosphate on tetanus toxin (TeNT) production, and *tent/tetR* expression. (A) Growth kinetics of CN655 in TGY supplemented with various concentrations of inorganic phosphate. (B) Extracellular TeNT levels. (C) Total TeNT levels. (D) Expression of *tent* and (E) *tetR*. Data are mean values  $\pm$  SEM of at least three independent cultures. \*,  $p < 0.05$ ; \*\*,  $p < 0.01$ .

### 2.7. Carbonate Stimulates TeNT Synthesis


High CO<sub>2</sub> concentrations (70%) have been found to increase the expression of *bont* and the synthesis of BoNT in *C. botulinum* group II in contrast to group I strains, where CO<sub>2</sub> has no significant stimulatory effect [27,28]. CO<sub>2</sub> also increases the toxin production in other bacteria such as *Bacillus anthracis*, *Staphylococcus aureus* and *Vibrio cholerae* [27]. However, the mechanism of the stimulatory effect of CO<sub>2</sub> is not well defined. Since *C. tetani* is phylogenetically related to *C. botulinum* [29], we tested the influence of carbonate as a nutritional requirement for TeNT synthesis. For that, the TGY medium was supplemented with various Na<sub>2</sub>CO<sub>3</sub> concentrations, and addition of 50 or 100 mM Na<sub>2</sub>CO<sub>3</sub> did not modify the growth of *C. tetani* CN655 (data not shown). We observed a significant stimulatory effect on TeNT synthesis in TGY medium supplemented with 100 mM Na<sub>2</sub>CO<sub>3</sub> (Figure 7). No synergistic or cumulative effect was detected between the addition of carbonate and P<sub>i</sub> (Figure 7). The pH at 72 h of culture was slightly higher (pH 7.6) in TGY supplemented with 100 mM Na<sub>2</sub>CO<sub>3</sub> compared to control TGY (pH 7.0) or TGY with 50 mM Na<sub>2</sub>CO<sub>3</sub> (pH 7.3). However, the pH in TGY cultures supplemented with 40 mM P<sub>i</sub> or 100 mM Na<sub>2</sub>CO<sub>3</sub>/40 mM P<sub>i</sub> was not significantly modified (pH 7.0 and 7.4, respectively), suggesting that the role of P<sub>i</sub> and Na<sub>2</sub>CO<sub>3</sub> on TeNT synthesis were not pH-dependent.

### 2.8. The TCS CTC\_RS05745/CTC\_RS05750 is Involved in Bacterial Cell Wall Organization


Since the TCS CTC\_RS05745/CTC\_RS05750 seemed to be involved in TeNT secretion without controlling the synthesis level of toxin (see above and Figure 2), we investigated the morphology of the strain CN655/p1311 in comparison to the wt CN655 by electron microscopy (Figure 8). Electron microscopic analysis of wt CN655 and strain CN655/p1311 showed a marked alteration of the bacterial wall of the mutant strain (Figure 8). In the wt CN655 strain, the layers forming the bacterial wall are well organized. In contrast, in strain CN655/p1311, the bacterial wall appeared disorganized and an abundantly detected diffuse material surrounded the bacteria. The altered bacterial wall of CN655/p1311 might account for the increased extracellular TeNT level.



**Figure 7.** Effects of sodium carbonate and inorganic phosphate on extracellular tetanus toxin (TeNT). *C. tetani* CN655 was grown in TGY supplemented with either 50 or 100 mM Na<sub>2</sub>CO<sub>3</sub>, 40 mM P<sub>i</sub>, or 100 mM Na<sub>2</sub>CO<sub>3</sub> and 40 mM P<sub>i</sub>. Extracellular toxin levels were increased in TGY medium supplemented with 40 mM P<sub>i</sub> or 100 mM Na<sub>2</sub>CO<sub>3</sub>. The addition of both P<sub>i</sub> and Na<sub>2</sub>CO<sub>3</sub> did not result in a synergistic effect on TeNT production. Data are mean values ± SEM of at least three independent assays. \*\*, *p* < 0.05.



**Figure 8.** Cont.




**Figure 8.** Ultrastructural morphology of CN655 and CN655/p1311. Bacteria from 18 h TGY culture were processed for transmission electron microscopy and scanning electron microscopy (SEM). CN655 showed well-delineated bacterial wall layers, whereas the bacterial wall of CN655/p1311 was disorganized with diffuse and enlarged wall layers. CN655/p1311 showed more abundant blebbings on the bacterial surface. About 100 bacterial cells were observed for each preparation.

### 3. Discussion

The mode of TeNT synthesis in *C. tetani* is still poorly understood. TeNT is usually produced by culturing *C. tetani* in complex media, and variability in TeNT production is a major concern in industrial processes. Carbon and nitrogen sources, notably peptides and amino acids, are important parameters for *C. tetani* metabolism and TeNT synthesis [6,30–32]. Toxin production in clostridia is typically a complex regulated process [33,34]. The gene *tetR*, which lies directly upstream of *tent*, was the first regulatory element identified in the regulation of TeNT production. The *tetR* gene encodes an alternative sigma factor that positively regulates the transcription of *tent* at the transition phase between the end of the exponential and the beginning of the stationary growth phases [13]. The genome of *C. tetani* strain E88 was found to contain at least 127 genes that encode putative transcriptional regulators, including 30 TCSs and 29 sigma factors [35]. However, their role in the regulation of TeNT synthesis is not yet known. Here, we have investigated 12 putative regulators. These have been selected based on their homology with regulators that have already been identified to be involved in the toxin regulation in other clostridia such as *C. botulinum* and *C. difficile*.

Among the nine TCSs which have been investigated by the RNA anti-sense strategy, two positively regulated the production of TeNT (CTC\_13810/CTC\_13815 and CTC\_RS10150/CTC\_RS10155) and one was identified as a negative regulator (CTC\_RS07310/CTC\_RS07315). Moreover, two other TCSs influenced the extracellular toxin level rather than the total production of TeNT, positively (CTC\_RS04780/CTC\_RS04785) and negatively (CTC\_RS05750/CTC\_RS05745) (Figure 9).



**Figure 9.** Schematic summary of the regulators of tetanus toxin (TeNT) synthesis of this study. Two two-component systems (TCSs) (RS13815 and RS10155) as well as CodY are positive regulators. They activate the transcription of *tetR* and *tent*, only CodY acts directly by interacting with the promoter of *tent* ( $P_{tent}$ ). In addition, the TCS RS04785 increases the extracellular TeNT level without or only weakly affecting the total synthesis by interacting with  $P_{tent}$ . One TCS (RS07315) is a negative regulator that interacts with ( $P_{tent}$ ) and ( $P_{tetR}$ ). The TCS RS05750 has a moderate negative effect by weakly and indirectly decreasing *tent* and *tetR* transcription. Inorganic phosphate (Pi) and carbonate are environmental factors that influence TeNT synthesis through the TCS pathway and/or through the general metabolism.

The TCSs CTC\_13810/CTC\_13805 and CTC\_RS10150/CTC\_RS10155 influenced the *tent-tetR* expression and subsequent TeNT production. However, *tent* and *tetR* transcription were not decreased in the exponential growth phase despite a reduction of TeNT production. This might result from a delayed effect of the RNA anti-sense system and/or of unproductive RNA hybridized with the anti-sense RNA. They seem to indirectly control the expression of *tent* and *tetR*, since the respective RRs bound neither to  $P_{tent}$  nor  $P_{tetR}$ .

The TCS CTC\_RS13810/CTC\_RS13805 is the only TCS located on the large *tetR-tent* operon-containing plasmid, and it is encoded about 25 kb upstream of *tent*. The CTC\_RS13805 RR belongs to the OmpR family (Table 1) and shares 91% amino acid identity with a RR of *Clostridium lundense* but has no homolog in other clostridia or bacteria.

The chromosomally encoded RR CTC\_RS10155 is related to the LytR/AlgR family of regulators and is putatively involved in autolysis. This TCS has homologs in *C. botulinum* and other clostridia. However, the TCS homolog in *C. botulinum* has no influence on BoNT production [25].

The *C. botulinum* TCS CBO0786/CBO0787 was previously found to repress BoNT synthesis [26]. This TCS has a homolog in *C. tetani* (CTC\_RS07310/CTC\_RS07315) with a 58% identity at the amino acid level of the RR, and it is predicted to be involved in cell division. This *C. tetani* RR was also found to repress TeNT synthesis. Indeed, secreted and total TeNT levels were increased in the anti-sense strain (CN655/p1419). Interestingly, the RR CTC\_RS07315 bound to  $P_{tent}$  and  $P_{tetR}$ , resulting in a significant decrease in *tent* expression at the stationary growth phase and a moderate impairment of *tetR* transcription at late exponential and early stationary growth phases. This suggests that the negative effect on *tent* transcription results from a direct interaction with  $P_{tent}$ , possibly independent of *tetR* since the effect on *tetR* transcription was weak. The mode of action of CTC\_RS07315 RR is likely similar to that of CBO0786, which binds to the *ntrh-bont/A* and *ha* operon promoters in *C. botulinum* and inhibits their transcription, in a *botR*-independent manner [26].

The TCS CTC\_RS05750 also exhibits a negative effect on TeNT secretion, since the respective anti-sense strain (CN655/p1311) showed an increased extracellular TeNT level. The CTC\_RS05750 RR belongs to the OmpR family and it is supposed to be involved in the phosphate uptake. CTC\_RS05750 moderately inhibits *tent* and *tetR* transcription, probably in an indirect manner since in contrast to the RR CTC\_RS07315, no binding of CTC\_RS05750 RR to  $P_{tent}$  and  $P_{tetR}$  was observed. This TCS probably controls pleiotropic pathways including the assembly of the bacterial wall. The regulatory pathways controlled by CTC\_RS05745/CTC\_RS05750 are still unknown and this TCS probably has an indirect role in cell wall synthesis. Indeed, a marked alteration of the bacterial wall was observed in CN655/p1311 compared to wt CN655 (Figure 8). The mechanism of TeNT secretion in *C. tetani* is not yet understood. For several decades, it has been known that *C. tetani* autolysis induces TeNT accumulation in the extracellular medium [36]. Cell wall modification in CN655/p1311 was likely responsible for the increased extracellular TeNT without affecting the total level of TeNT synthesis. CTC\_RS05745/CTC\_RS05750 is homologous to related TCSs in other clostridia, including *C. botulinum*. However, the *C. botulinum* homolog CLC\_2386/CLC\_2385 (73% amino acid identity between the corresponding RRs) was not found to be involved in the control of BoNT synthesis [25].

The anti-sense strain CN655/p1310, in which CTC\_RS04785 RR was targeted, exhibited a slight decrease in the amount of extracellular TeNT and a marked decrease in *tetR* expression. However, only a slight decrease in *tent* expression was observed, albeit CTC\_RS04785 RR interacted with  $P_{tent}$  but not with  $P_{tetR}$ . CTC\_RS04785 RR has a 34% amino acid identity with the virulence factor VirR of *C. perfringens*. VirS-VirR controls 147 genes in *C. perfringens* including chromosomally-located toxin genes encoding, for example, perfringolysin and alpha-toxin, and plasmid-located toxin genes such as those encoding the Beta2 toxin and collagen-adhesin [37,38]. The VirS-VirR system of *C. perfringens* can positively or negatively regulate its target genes. Some of the target genes contain a VirR-binding box on their promoter and are directly controlled, whereas a regulatory cascade including a non-coding RNA (VR-RNA) is used in the regulation of other genes [39]. The CTC\_RS04780/CTC\_RS04785 TCS of *C. tetani* is possibly also involved in a complex network of positive and negative regulatory pathways in *C. tetani*. Our results suggest that this TCS mainly modulates the secretion, and moderately the synthesis, of TeNT, although CTC\_RS04785 was found to bind to the *tent* promoter and to stimulate *tetR* transcription.

CodY is a conserved regulator in Gram-positive bacteria, which notably controls metabolism and virulence [17]. In *C. botulinum* A, CodY positively regulates BoNT/A synthesis through binding to the *ntrh-bontA* operon promoter in a GTP-dependent manner [20]. CodY also activates toxin production and virulence in other Gram-positive bacteria such as *Bacillus anthracis*, *Bacillus cereus*, *Listeria monocytogenes* and several *Streptococcus* species [40–44]. However, CodY indirectly represses toxin gene expression in *C. difficile* by interacting with the TcdR promoter [45]. Rapidly metabolizable carbohydrates such as glucose repress toxin production in *C. difficile* [46], and CodY is a negative regulator of toxin synthesis and virulence in this pathogen [47]. In contrast to *C. difficile*, TeNT synthesis in *C. tetani* is not inhibited by glucose. TeNT production is higher when cultivation of *C. tetani* is performed in TGY compared to TY media (Supplementary Materials Figure S5). *C. tetani* CodY shares

81% amino acid identity with that of *C. botulinum* type A strain ATCC3502, in which CodY positively controls BoNT/A synthesis [20]. In analogy to its activity in *C. botulinum*, CodY of *C. tetani* bound to  $P_{tent}$  but not to  $P_{tetR}$ , and enhanced *tent* transcription and subsequent TeNT synthesis. This further supports the interaction between metabolism and toxin production in *C. tetani*. Since CodY plays a key role in the adaptation to starvation [19], it could control TeNT synthesis via sensing the availability of specific nutrients.

Spo0A is also a master regulator that has been initially identified to control the first steps of sporulation, it also regulates numerous other genes, notably those involved in adaptation to environmental conditions [48]. For example, Spo0A is involved in the switch between the acidogenesis and solventogenesis/sporulation pathways in environmental bacteria such as *Clostridium acetobutylicum* [49]. Here, Spo0A was not found to control TeNT synthesis, in contrast to its positive impact on BoNT synthesis in *C. botulinum* E through binding to the *bontE* promoter [50]. It is noteworthy that *C. botulinum* E gene cluster does not contain any *botR* or *tetR* homologs in contrast to *C. tetani* and *C. botulinum* group I neurotoxin gene clusters. It is questionable whether there is an interplay between Spo0A, BotR/TetR and toxin synthesis. Spo0A controls the sporulation in *C. botulinum* [23], but we observed no correlation between sporulation and toxin production in *C. botulinum* A [51]. *C. tetani* CN655 formed no spore in TGY, albeit this strain synthesized TeNT. CN655 sporulates only poorly (10–100 spores/ml) in the biphasic medium of Anellis et al. [52], and spores were not detected in CN655/*spo0A* cultures in the biphasic medium, indicating that the anti-sense *spo0A* was functional in CN655. In *C. difficile*, Spo0A differently modulates the toxin synthesis, which depends on the genetic background. In certain *C. difficile* strains such as ribotype 027 strains, Spo0A negatively regulates the production of Toxin A and Toxin B, whereas in other strains, Spo0A has no effect on toxin production [53].

In *C. difficile*, the protein Mfd is involved in nucleotide excision repair and transcription elongation. This protein has been identified as a positive regulator of Toxin A and Toxin B synthesis. Mfd possibly prevents the inhibitory effect of CodY and CcpA (carbon catabolite protein A) by inhibiting their binding to the toxin gene promoters [24]. Mfd is conserved in *C. tetani*. However, it had no impact on TeNT synthesis under the conditions tested. In contrast to *C. difficile*, TeNT synthesis is not repressed by rapidly fermentable carbohydrates [54].

Albeit numerous regulatory genes are conserved in clostridia, they have distinct pleiotropic effects in each species. Our data indicate that *C. tetani* retains unique regulatory pathways to control TeNT synthesis, possibly as an adaptation to specific nutrients and/or environmental conditions.

The complex network of toxin regulation in *C. tetani* raises the question of the environmental factors that trigger the different regulatory components of the network. Since some TCSs are putatively involved in  $P_i$  uptake and/or metabolism, we tested the influence of  $P_i$ . We observed that an optimum  $P_i$  concentration around 40 mM significantly increased the transcription of *tent* and subsequent production of TeNT. In *C. perfringens*, 20 to 50 mM  $P_i$  concentrations enhance the sporulation and production of *C. perfringens* enterotoxin, the gene expression of which is dependent on sporulation.  $P_i$  induces *spo0A* expression and subsequent sporulation [55,56]. However, we observed no impact of  $P_i$  supplementation on sporulation of CN655 in TGY medium.  $P_i$  was also found to increase the toxin production in *Bacillus thuringiensis* [57].  $P_i$  is a key element in bacterial metabolism; notably, it is incorporated into transcriptional regulatory proteins, and  $P_i$  homeostasis has a critical adaptive role in bacterial virulence [58]. For example,  $P_i$  availability controls motility, biofilm formation, colonization, induction of virulence factors such as C phospholipases in *Pseudomonas aeruginosa*, colonization factor and toxin synthesis in *Vibrio cholerae*, enteropathogenic and enterohemorrhagic *Escherichia coli* (review in Reference [58]). In *Bacillus anthracis*,  $P_i$  starvation enhances spore germination, invasiveness in macrophages and toxin secretion [59]. It can be speculated that  $P_i$  availability might also facilitate wound colonization by *C. tetani* and in situ TeNT production. The mechanism of  $P_i$  in the control of *tent* transcription is not yet known. It is possibly mediated by TCS such as CTC\_RS05745/CTC\_RS05750 and CTC\_RS10030/CTC\_RS10035, which are putatively involved in

phosphate uptake. Indeed, the corresponding anti-sense strains (CN655/p1311 and CN655/p1313) grown in TGY supplemented with  $P_i$  (40 mM) showed slightly increased extracellular TeNT compared to culture in TGY (Supplementary Materials Figure S6). However, these two TCSs might have pleiotropic effects and indirectly control TeNT synthesis, since they are homologs of the PhoP/PhoR TCS which has been found to control diverse metabolism pathways such as redox homeostasis and adaptation to acidic pH in *Mycobacterium*, as well as primary and secondary metabolism pathways under phosphate limitation [60,61].

$CO_2$  at high concentration (70%) in the gas phase stimulates *bont* gene expression and BoNT production, despite a reduced growth rate of *C. botulinum* group II strains, in which the regulatory *botR* gene is missing. The mode of action of  $CO_2$  is still speculative: it is supposed to dissolve and increase the bicarbonate concentration in the TGY culture medium and subsequently to enhance carboxylation reactions [62]. Increased bicarbonate concentrations were also found to enhance toxin production in *C. difficile*, possibly through biotin-dependent carboxylation [63]. In our conditions, the extracellular TeNT levels were increased in TGY media containing 100 mM  $Na_2CO_3$  in a pH-independent manner, whereas no significant difference was observed with 50 mM  $Na_2CO_3$  supplementation. The involvement of carbonate in carboxylation reactions is more likely responsible for the regulatory effects on TeNT synthesis than the buffer effect of carbonate.

In conclusion, TeNT synthesis in *C. tetani* is under the control of a complex network of regulatory elements, including TCSs and regulators of metabolism, as tested by the anti-sense RNA system. However, a more direct approach by deletion of the putative target regulatory genes is required to confirm the regulatory function on TeNT synthesis of these genes. *C. tetani* and *C. botulinum* are related clostridial species which synthesize potent neurotoxins in a regulated manner, and their genomes contain numerous TCSs genes (30 in *C. tetani* and 39 in *C. botulinum* A [25,35]). Interestingly, albeit most of TCSs genes are homologous in both species, *C. tetani* and *C. botulinum* mainly use distinct sets of TCSs in the regulation of toxin synthesis. Among the nine RRs investigated in *C. tetani*, only one system (CTC\_RS07315 in *C. tetani* and the homolog CBO0786 in *C. botulinum*) has a similar function in both species, i.e., serves as a negative regulator of toxin synthesis [26]. Four other RRs modulate TeNT synthesis (one acting as negative and three as positive regulators), whereas their homologous counterparts in *C. botulinum* A have no effect on toxin production [25]. In addition, two TCSs, which positively control BoNT production in *C. botulinum*, have homologs in *C. tetani* that do not impact on TeNT synthesis. We hypothesize that *C. tetani* has adapted TeNT synthesis to specific nutritional and environmental requirements. Indeed, *C. tetani* preferentially uses specific metabolic pathways [35]. Notably, peptides and amino acids are critical substrates for *C. tetani* and specific short peptides from casein digestion are essential for TeNT synthesis [32,35]. Here, we also found that  $P_i$  and carbonate are additional nutritional requirements for TeNT production. Thereby, TeNT synthesis is dependent of a complex network of regulation linked to *C. tetani* metabolism, in which carbohydrates and  $P_i$  are important elements. A better understanding of the regulation of TeNT synthesis and the underlying environmental factors is required to optimize the toxin production by *C. tetani* for vaccine production. Albeit tetanus is rare in the developed world, it is still an important cause of deaths in many underdeveloped countries, with 34,000 estimated neonatal tetanus deaths in 2015 [64]. Immunization against TeNT is an efficient method of tetanus prevention, and thus, a 96% reduction of this disease was achieved since 1988 [64]. Availability of a cheap and efficient tetanus vaccine is an important factor for the eradication of this disease. Unraveling the regulation of TeNT synthesis constitutes the basis for constructing hyperproductive strains.

## 4. Materials and Methods

### 4.1. Bacterial Strains and Culture Conditions

The recombinant strains used in this study are presented in Table 2. *Escherichia coli* strains were grown in Luria-Bertani (LB) broth and *C. tetani* CN655 in TGY broth (pH 7.5) containing trypticase

Peptone (BD Biosciences; 30 g/L), yeast extract (Bacto Yeast Extract, BD Biosciences; 20 g/L), glucose (5 g/L) and cysteine, HCl (0.5g/L) under anaerobic conditions (N<sub>2</sub>/CO<sub>2</sub>/H<sub>2</sub>; 90: 5:5, vol/vol.) at 37 °C. When necessary, erythromycin was added to culture media at 5 or 50 µg/ml for *C. tetani* and 300 µg/ml for *E. coli*. *C. tetani* was grown in TGY with different inorganic phosphate (P<sub>i</sub>) concentrations by the addition of Na<sub>2</sub>HPO<sub>4</sub>, and/or with sodium carbonate Na<sub>2</sub>CO<sub>3</sub> (Merck, Guyancourt, France).

Growth kinetics of *C. tetani* CN655 strains in TGY broth culture medium supplemented with erythromycin (5 µg/mL) were monitored by spectrometry at 600 nm over a 144 h period.

#### 4.2. Construction of Vectors Encoding Anti-Sense mRNA for the Different Two-Component Systems (TCS) and Others Regulators

A DNA fragment of each TCS and regulatory genes studied containing the ribosome binding site (RBS) region was amplified by PCR and inserted in reverse orientation into pAT18, as already described in Reference [14]. DNA segments for anti-sense mRNA production were designed in the RR of three TCS, in the SHK gene of six TCS, as well as in three regulatory genes (*codY*, *spo0A* and *mfd*) (Table 2). The PCR products from CN655 genome DNA contain a 3' *NcoI* site and a 5' *PstI* site and were cloned into pMRP306, a derivative of pAT19 containing the promoter of the iota toxin gene, the cloning sites *NcoI-PstI* and the 3' part of the iota toxin gene [14]. The resultant anti-sense RNA plasmids were transformed into CN655 by electroporation.

The recombinant plasmids were prepared in *Escherichia coli* Top10 strain (Dam<sup>+</sup>, Dcm<sup>+</sup>) and electroporated into *C. tetani* CN655, with an efficiency of 25–50 transformants per 1 µg DNA. Interestingly, no transformant was obtained from plasmids prepared in *E. coli* BL21 (Dam<sup>+</sup>, Dcm<sup>-</sup>) or C2925 (Dam<sup>-</sup>, Dcm<sup>-</sup>), indicating that *C. tetani* CN655 contains a functional Dcm methylase that impairs the transformation of non-DCM-methylated DNA.

#### 4.3. Tetanus Toxin Assay

At 8, 24, 32, 48, 56, 72 and 144 h of culture growth, 4 mL of the culture were removed. The cells were harvested at 10000 rpm for 10 min at 4 °C, and the supernatants corresponding to the extracellular toxin were filtered (0.22 µm) and stored at -20 °C. For intracellular toxin assay, the pellets were washed twice with distilled water and two osmotic lysis were with TGY containing 20 mg/mL NaCl and 13.3 mg/mL sodium citrate (C<sub>6</sub>H<sub>5</sub>Na<sub>3</sub>O<sub>7</sub>, 2H<sub>2</sub>O) during 24 h, at 4 °C. after centrifugation, the supernatants containing intracellular toxin were filtered and stored at -20°C.

Extracellular and intracellular toxin were monitored by an enzyme-linked immunosorbent assay (ELISA). Wells of microtiter plates (Nunc Maxisorp; Nunc, Roskilde, Denmark) were coated with 100 µL of equine anti-tetanus serum (Sanofi Pasteur, FA193727) in 0.05 M carbonate buffer pH 9.5 and incubated at 4 °C overnight. Plates were washed 3 times with phosphate buffered saline (PBS)-Tween20 0.1% with an automatic plate washer (BioTek, Washer 120, BioTek France, Colmar, France). After blocking with 20 mg/ml BSA in carbonate buffer during 30 minutes under agitation, three washes were performed, and 100 µl/well of serial two-fold dilutions in PBS-Tween20 0.1%-BSA 1% (PBS-T-BSA) of samples were added. The plates were incubated for 1 h at room temperature with shaking. Tetanus toxin (Sanofi Pasteur, Marcy l'Etoile, France) was used as standard. After three washes, the plates were incubated with 100 µL/well of rabbit anti-tetanus serum (1:6400; Sanofi Pasteur n° 7078) for 1 h at room temperature, then with 100 µl of goat anti-rabbit Ig peroxidase-linked (1:4000, 111-035-006, Jackson ImmunoResearch) for 1 h at room temperature in PBS-T-BSA. For detection, 100 µL of 1 mg/mL ortho-phenylene-diamine (OPD, Sigma) in citrate buffer (0.05 M, pH 4.5 containing 0.06% H<sub>2</sub>O<sub>2</sub>) was used. The color development was stopped after 8 min by adding 50 µL 3 M HCl. The absorbance was read on a microplate reader (Biorad, model 680) at 490 and 655 nm.

#### 4.4. Total RNA Extraction, Reverse Transcription and Quantitative Real-Time PCR Assay

Total RNA from *C. tetani* strains were extracted at 8, 24, 32 and 48 h of growth. After centrifugation at 4000 rpm for 15 min at 4 °C, the culture pellet was mechanically disrupted in the presence of

silica beads (Lysing Matrix B, MP Biomedicals, Illkirch, France) and buffer RLT (MP Biomedicals) by shaking with a FastPrep apparatus (MP Biomedicals), and RNA total extracted using RNeasy mini kit (Qiagen, Courtaboeuf, France), according to the manufacturer's instructions. The RNA preparations were stored at  $-80^{\circ}\text{C}$ .

A DNase treatment with TURBO DNase (Ambion, ThermoFischer, Les Ulis, France) was performed following the manufacturer's instructions. The absence of DNA contamination on RNA extracts was checked by real-time PCR (RT-PCR) targeting *tent*. Total RNA amount was monitored with NanoDrop ND-100 Spectrophotometer. cDNAs were then synthesized from 1  $\mu\text{g}$  of total RNA with random primers (pDN6 5  $\mu\text{g}/\mu\text{L}$ , Roche, Meylan France) and RNase OUT™ Recombinant Ribonuclease Inhibitor (Invitrogen, ThermoFischer, Les Ulis, France) and with M-MLV Reverse Transcriptase kit (Invitrogen), according to the manufacturer's instructions.

RT-PCR was performed in 25  $\mu\text{L}$  reaction volume containing 30 ng of cDNAs, 12.5  $\mu\text{L}$  of SYBR Green Supermix (Bio-Rad, 2 $\times$ ; 1.25 U iTaq DNA polymerase, 0.4 mM each dNTP, 6 mM  $\text{MgCl}_2$ , 20 nM fluorescein, SYBR Green I) and 500 nM gene-specific primers (Table 3) in an iQ iCycler apparatus (Bio-Rad). The reaction was subjected to denaturation at  $95^{\circ}\text{C}$  for 3 min followed by 40 cycles of denaturation at  $95^{\circ}\text{C}$  for 10 s, annealing/elongation at  $61.7^{\circ}\text{C}$  for 30 s for *rpoB* and *tent* genes and  $65.1^{\circ}\text{C}$  for *gyrA* and *tetR* genes. Then, a dissociation stage of 65 to  $95^{\circ}\text{C}$  with a heating rate of  $0.5^{\circ}\text{C}$  per 10 s was performed to establish a melting curve to confirm the specificity of the RT-PCR reaction for each primer pair.

The relative cDNA quantity of each sample was determined with the threshold cycle [ $\Delta\Delta\text{CT}$ ] method (Analysis of Relative Gene Expression Data Using Real-Time Quantitative PCR and the  $2\Delta\Delta\text{CT}$  method). cDNA quantity of the *tent* and *tetR* gene was normalized to the quantity of cDNA of the *rpoB* and *gyrA* gene.

**Table 3.** Primers used for qRT-PCR, protein expression and promoter regions of *tent* and *tetR* for EMSA experiments.

Target Gene	Primer	Nucleotide Sequence (5'→3')	Product Length (bp)
<b>qRT-PCR</b>			
<i>tent</i>	P1714-F	CCAAGGTGCACAAGGAATT	146
	P1715-R	CAATGTTAATGCGGGTCCT	
<i>tetR</i>	P1726-F	GTTGCTCAAATATTAAACTTCGAA	115
	P1727-R	GCTATATCACATTCTTCATATCTTCAA	
<i>rpoB</i>	P2142-F	TTGAAGAATGTAAAGAGAGAGATGCTAC	118
	P2143-R	GGGAAGTCACCCATAAAGACA	
<i>gyrA</i>	P2146-F	AAGATGATGTAGCAGTAAGTATGGA	98
	P2147-R	CTCTGAAGCCAATGTCCTTTT	
<b>Recombinant protein expression</b>			
CTC_p21	P2349-F P2350-R	CGCCGCGATCCATGTATAAGATATTGATTGTGAA CCGCCGGAATTCCTTACACCTGAAATAAACGATAGCC	711
CTC_01979	P2351-F P2352-R	CGCCGCGATCCATGAACAAAATAAATTGTGTAATAATA CCGCCGGAATTCCTAAAAATCTAATATGTCCTTTAAGTG	792
CTC_01421	P2353-F P2354-R	CGCCGCGATCCGTGAACAACATATTGTTAGTTGAA CCGCCGGAATTCCTATTATTAATTTTCGTAGTTCCACCT	717
codY	P2355-F P2356-R	CGCCGATCCATGTCATCATTATTAGAGAAG CCGCCGGAATTCCTACTTAATTTTTTCAATTCCTC	801
CTC_00935	P2357-F P2358-R	CGCCGATCCGTGTGTAGAGTAGTGCTT CCGCCGGAATTCCTTATACTTTTTTATTATTAC	759
<b>EMSA</b>			
<i>Ptent</i>	P2365-F P2366-R	(5'-end labelled biotin) GGTGGCTCCATCATAATAATTGTAT (5'-end labelled biotin) GGTTTTAGCATTAATAAATTAGAACCTA	359
	P2363-F P2364-R	(5'-end labelled biotin) CAGTATTTTGAATGTATAATAATTACTTC (5'-end labelled biotin) CCGTTCCTTAATTTAGTAATAATCAATAT	

F, forward; R, reverse.

#### 4.5. Expression and Purification of Recombinant Proteins

The genes *ctc\_p21*, *ctc\_01979*, *ctc\_1421*, *ctc\_00935*, *codY* and *tetR* were PCR-amplified from the genome of *C. tetani* CN655, with primers (Table 3), adding a *Bam*HI site at the 5' end and an *Eco*RI at the 3' end. PCR products were digested with appropriate restriction enzymes and the resulting products were cloned into a pCR2.1 vector and subcloned into pET28a (Novagen, Merck, Guyancourt, France) for the expression of N-terminal 6-histidine proteins. The resulting constructions were transformed in *E. coli* BL21DE3, according to the manufacturer's instructions.

Recombinant proteins were purified by affinity chromatography employing the TALON cobalt affinity resin (Clontech Laboratoires, Saint Germain n Laye, France) in accordance with the standard protocol provided by the manufacturer. Briefly, to induce expression of recombinant proteins, clones were grown in 1 liter of LB supplemented with 50 µg/mL kanamycin at 37 °C to an optical density at 600 nm of 0.6–0.8. Protein expression was then induced by the addition of isopropyl-β-D-thiogalactopyranoside (IPTG) at a final concentration of 0.5 mM and the growth was continued overnight at 18 °C. The bacteria were collected by centrifugation, suspended in PBS pH 8.0 containing 10 mM imidazole and protease inhibitors (EDTA free protease inhibitor cocktail, Roche), and lysed by sonication. The recombinant proteins were eluted with 100 mM imidazole and dialyzed overnight against PBS and then frozen at –80 °C for storage. Quantification of the protein was done using the Bradford reagent (BioRad, Marnes La Coquette, France), following the manufacturer's instructions. Aliquots of fractions were analyzed on 12% sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE).

#### 4.6. Electrophoretic Mobility Shift Assay (EMSA)

A 359 bp fragment covering *tent* promoter ( $P_{tent}$  probe) and a 316 bp fragment covering *tetR* promoter ( $P_{tetR}$ ) were amplified by PCR using 5'-end biotin-labeled primers (Table 3). Binding reactions were carried out for 1 h at room temperature using 5 µM of recombinant proteins, 5'-end biotin-labeled probe at 0.2 nM for  $P_{tent}$  and 0.1 nM for  $P_{tetR}$ , 50 ng/µL of poly (dI-dC), 2.5% glycerol and 5 mM MgCl<sub>2</sub> in binding buffer. Competition assays for binding specificity were performed with a 300-fold excess of unlabeled specific probe.

Reactions were separated on a 5% native polyacrylamide gel, run in 0.5× Tris-borate buffer-EDTA (TBE) at 4 °C for 1 h at 110 V and electrotransferred to a positively charged nylon membrane (Amersham) at 380 mA for 30 min. Transferred DNAs were cross-linked to membranes with UV light. The biotin-labeled DNAs were detected with the LightShift Chemiluminescent EMSA kit (Pierce, ThermoFischer, Les Ulis, France) according the manufacturer's specifications.

#### 4.7. Electron Microscopy

Bacteria from 18 h culture were mixed V/V with a fixative solution containing 5% glutaraldehyde (GA) in 0.2 M PHEM buffer pH 7.2 (60 mM Pipes, 25 mM Hepes, 10 mM EGTA, 2 mM MgCl<sub>2</sub>) and incubated for 1 h. Samples were washed twice in PBS prior to performing a high-pressure (>2000 bar) freezing in 1-hexadecane using a BAL-TEC HPM 010 (LEICA). Freeze substitution was done with 2% OsO<sub>4</sub> and 0.5% Uranyl Acetate in acetone followed by several steps: –90 °C for 42 h, warmed up to –30 °C (5 °C/h), incubation for 12 h, warmed up to 0 °C (10 °C/h) and incubation for 1 h. Then, samples were washed with acetone on ice and incubated with increasing low-viscosity embedding media SPURR's kit (EMS ref. 14300)/acetone mixture (1:4) for 3 h, 1:1 over-night, 3:1 during the day, 9:1 over-night, SPURR resin for the day and over-night. The samples were then placed into flat tubes infiltrated with pure SPURR resin prior to polymerization at 60 °C for 48 h. Sections (60–70 nm) were obtained on a FC6/UC6 ultramicrotome (Leica, Wetzlar, Germany), transferred on 200 Mesh Square Copper grids coated with formvar and carbon (CF-200-Cu50, Delta Microscopy). Samples were stained with 4% uranyl acetate and counterstained with lead citrate. Images were recorded with

TECNAI SPIRIT 120 kv (with a bottom-mounted EAGLE 4K x 4K Camera, ThermoFisher Scientific, Waltham, MA, USA).

For scanning electron microscopy (SEM), the samples dehydrated with ethanol were dried with CO<sub>2</sub>, and then sputtered with 20 nm gold palladium with a GATAN Ion Beam Coater and were examined with a JEOL JSM 6700F field emission scanning electron microscope operating at 7 Kv. Images were acquired with the upper SE detector (SEI).

#### 4.8. Statistical Analysis

Values throughout are expressed as means ± standard error of the mean. Differences in the different anti-sense strains were assessed using unpaired Student's t-test, where statistical significance is assumed for \*  $p < 0.05$ , \*\*  $p < 0.01$  and \*\*\*  $p < 0.001$ .

**Supplementary Materials:** The following are available online at <http://www.mdpi.com/2072-6651/12/5/328/s1>: Figure S1: Growth kinetics and tetanus toxin (TeNT) production in CN655 wt and CN655/pAT18. Growth kinetics (A) of CN655/pAT18 was slightly lower than that of CN655 wt during the first 56 h, but the total production of TeNT (B) was slightly higher. In the stationary phase from 72 to 144 h, growth kinetics and total TeNT production were similar in both CN655/pAT18 and CN655 wt. Data are mean values ± SEM of at least three independent cultures. Figure S2: Extracellular tetanus toxin (TeNT) produced by CN655/pAT18 (empty vector) and CN655 anti-sense strains showing no significant difference. CN655/p1308, CN655/p1309, CN655/p1312, CN655/p1313, CN655/p1472 and CN655/p1480 showed no significant difference in extracellular levels of TeNT compared to CN655/pAT18. Data are mean values ± SEM of at least three independent cultures. Figure S3: Total tetanus toxin (TeNT) produced by CN655/pAT18 (empty vector) and CN655 anti-sense strains showing no significant difference. CN655/p1308, CN655/p1309, CN655/p1310, CN655/p1311, CN655/p1312 and CN655/p1313 showed no significant difference in total TeNT levels compared to CN655/pAT18. Data are mean values ± SEM of at least three independent cultures. Figure S4: Expression of (A) *tent* and (B) *tetR* in CN655/pAT18 and CN655 anti-sense strains showing no significant difference. (A) CN655/p1308, CN655/p1309, CN655/p1312 and CN655/p1313 showed no significant difference in *tent* expression compared to CN655/pAT18. (B) CN655/p1308 and CN655/p1309 showed no significant difference in *tetR* expression compared to CN655/pAT18. Data are mean values ± SEM of at least three independent cultures. Figure S5: Growth kinetics and tetanus toxin (TeNT) production in CN655 wt in TGY versus TY culture medium. The presence of glucose (5 g/L) in culture medium (TGY) induced a slightly increased growth and a more than two-fold extracellular TeNT compared to TY medium without supplementation in glucose. Data are mean values ± SEM of at least three independent cultures. Figure S6: Extracellular tetanus toxin (TeNT) produced by CN655/p1311 and CN655/p1313 grown in TGY supplemented with inorganic phosphate. CN655/p1311 and CN655/p1313 were grown in TGY and in TGY supplemented with 40 mM Na<sub>2</sub>HPO<sub>4</sub>. Extracellular TeNT was monitored by ELISA. Data are mean values ± SEM of at least three independent assays.

**Author Contributions:** D.C.-M., H.B., M.R.P. conceived and designed the experiments; D.C.-M., H.B., M.R.P., C.D., G.H., O.G., C.S., C.T. performed the experiments; D.C.-M., H.B., M.R.P. analyzed the data; L.P., D.G., F.B., V.C., S.D. contributed reagents/materials/analysis tools; M.R.P., H.B. wrote the manuscript. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research was funded by grant from Sanofi-Pasteur, Marcy l'Etoile and by Institut Pasteur, Paris.

**Conflicts of Interest:** LP, DG, FB, VC, and SD were employed by the company Sanofi-Pasteur. The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

#### References

1. Popoff, M.R.; Mazuet, C.; Poulain, B. Botulism and tetanus. In *The Prokaryotes: Human Microbiology*, 4th ed.; Springer: Berlin/Heidelberg, Germany, 2013; Volume 5, pp. 247–290.
2. Bercsenyi, K.; Schmiege, N.; Bryson, J.B.; Wallace, M.; Caccin, P.; Golding, M.; Zanotti, G.; Greensmith, L.; Nischt, R.; Schiavo, G. Tetanus toxin entry. Nidogens are therapeutic targets for the prevention of tetanus. *Science* **2014**, *346*, 1118–1123. [[CrossRef](#)] [[PubMed](#)]
3. Rossetto, O.; Scorzeto, M.; Megighian, A.; Montecucco, C. tetanus neurotoxin. *Toxicon* **2013**, *66*, 59–63. [[CrossRef](#)] [[PubMed](#)]
4. Schmiege, N.; Bercsenyi, K.; Schiavo, G. Uptake and transport of clostridial neurotoxins. In *The Comprehensive Sourcebook of Bacterial Protein Toxins*, 4th ed.; Alouf, J., Ladant, D., Popoff, M.R., Eds.; Elsevier: Amsterdam, The Netherlands, 2015; pp. 337–360.
5. WHO. Tetanus vaccines: WHO position paper—February 2017. *Wkly. Epidemiol. Rec.* **2017**, *92*, 53–76.

6. Licona-Cassani, C.; Steen, J.A.; Zaragoza, N.E.; Moonen, G.; Moutafis, G.; Hodson, M.P.; Power, J.; Nielsen, L.K.; Marcellin, E. Tetanus toxin production is triggered by the transition from amino acid consumption to peptides. *Anaerobe* **2016**, *41*, 113–124. [[CrossRef](#)] [[PubMed](#)]
7. Laird, W.J.; Aaronson, W.; Silver, R.P.; Habig, W.H.; Hardegree, M.C. plasmid-associated toxigenicity in *Clostridium tetani*. *J. Infect. Dis.* **1980**, *142*, 623. [[CrossRef](#)] [[PubMed](#)]
8. Finn, C.W.; Silver, R.P.; Habig, W.H.; Hardegree, M.C.; Zon, G.; Garon, C.F. The structural gene for tetanus neurotoxin is on a plasmid. *Science* **1984**, *224*, 881–884. [[CrossRef](#)]
9. Brüggemann, H.; Bäumer, S.; Fricke, W.F.; Wiezr, A.; Liesegang, H.; Decker, I.; Herzberg, C.; Martinez-Arias, R.; Henne, A.; Gottschalk, G. The genome sequence of *Clostridium tetani*, the causative agent of tetanus disease. *Proc. natl. Acad. Sci. USA* **2003**, *100*, 1316–1321. [[CrossRef](#)]
10. Brüggemann, H.; Brzuszkiewicz, E.; Chapeton-Montes, D.; Plourde, L.; Speck, D.; Popoff, M.R. Genomics of *Clostridium tetani*. *Res. Microbiol.* **2015**, *166*, 326–331. [[CrossRef](#)]
11. Chapeton-Montes, D.; Plourde, L.; Bouchier, C.; Ma, L.; Diancourt, L.; Criscuolo, A.; Popoff, M.R.; Brüggemann, H. The population structure of *Clostridium tetani* deduced from its pan-genome. *Sci. Rep.* **2019**, *9*, 11220. [[CrossRef](#)]
12. Fournier, P.E.; Levy, P.Y.; Million, M.; Croce, O.; Blanc-Taille, C.; Brouqui, P.; Raoult, D. Genome of a chronic osteitis-causing *Clostridium tetani*. *New Microbes New Infect.* **2014**, *2*, 25–26. [[CrossRef](#)]
13. Marvaud, J.C.; Eisel, U.; Binz, T.; Niemann, H.; Popoff, M.R. *tetR* is a positive regulator of the tetanus toxin gene in *Clostridium tetani* and is homologous to *botR*. *Infect. Immun.* **1998**, *66*, 5698–5702. [[CrossRef](#)] [[PubMed](#)]
14. Marvaud, J.C.; Gibert, M.; Inoue, K.; Fujinaga, V.; Oguma, K.; Popoff, M.R. *botR* is a positive regulator of botulinum neurotoxin and associated non toxic protein genes in *Clostridium botulinum* A. *Mol. Microbiol.* **1998**, *29*, 1009–1018. [[CrossRef](#)] [[PubMed](#)]
15. Dupuy, B.; Raffestin, S.; Matamouros, S.; Mani, N.; Popoff, M.R.; Sonenshein, A.L. Regulation of toxin and bacteriocin gene expression in *Clostridium* by interchangeable RNA polymerase sigma factors. *Mol. Microbiol.* **2006**, *60*, 1044–1057. [[CrossRef](#)] [[PubMed](#)]
16. Raffestin, S.; Dupuy, B.; Marvaud, J.C.; Popoff, M.R. BotR/A and TetR are alternative RNA polymerase sigma factors controlling the expression of the neurotoxin and associated protein genes in *Clostridium botulinum* type A and *Clostridium tetani*. *Mol. Microbiol.* **2005**, *55*, 235–249. [[CrossRef](#)] [[PubMed](#)]
17. Brinsmade, S.R. CodY, a master integrator of metabolism and virulence in Gram-positive bacteria. *Curr. Genet.* **2017**, *63*, 417–425. [[CrossRef](#)] [[PubMed](#)]
18. Mitrophanov, A.Y.; Groisman, E.A. Signal integration in bacterial two-component regulatory systems. *Genes Dev.* **2008**, *22*, 2601–2611. [[CrossRef](#)] [[PubMed](#)]
19. Stenz, L.; Francois, P.; Whiteson, K.; Wolz, C.; Linder, P.; Schrenzel, J. The CodY pleiotropic repressor controls virulence in gram-positive pathogens. *FEMS Immunol. Med. Microbiol.* **2011**, *62*, 123–139. [[CrossRef](#)]
20. Zhang, Z.; Dahlsten, E.; Korkeala, H.; Lindström, M. Positive regulation of botulinum neurotoxin gene expression by CodY in *Clostridium botulinum* ATCC 3502. *Appl. Environ. Microbiol.* **2014**, *80*, 7651–7658. [[CrossRef](#)]
21. Mascher, G.; Mertaola, A.; Korkeala, H.; Lindstrom, M. Neurotoxin synthesis is positively regulated by the sporulation transcription factor Spo0A in *Clostridium botulinum* type E. *Environ. Microbiol.* **2017**, *19*, 4287–4300. [[CrossRef](#)]
22. Kirk, D.G.; Palonen, E.; Korkeala, H.; Lindstrom, M. Evaluation of normalization reference genes for RT-qPCR analysis of *spo0A* and four sporulation sigma factor genes in *Clostridium botulinum* Group I strain ATCC 3502. *Anaerobe* **2014**, *26*, 14–19. [[CrossRef](#)]
23. Wörner, K.; Szurmant, H.; Chiang, C.; Hoch, J.A. Phosphorylation and functional analysis of the sporulation initiation factor Spo0A from *Clostridium botulinum*. *Mol. Microbiol.* **2006**, *59*, 1000–1012. [[CrossRef](#)] [[PubMed](#)]
24. Willing, S.E.; Richards, E.J.; Sempere, L.; Dale, A.G.; Cutting, S.M.; Fairweather, N.F. Increased toxin expression in a *Clostridium difficile* *mfid* mutant. *BMC Microbiol.* **2015**, *15*, 280. [[CrossRef](#)] [[PubMed](#)]
25. Connan, C.; Brüggemann, H.; Mazuet, C.; Raffestin, S.; Cayet, N.; Popoff, M.R. Two-component systems are involved in the regulation of botulinum neurotoxin synthesis in *Clostridium botulinum* type A strain Hall. *PLoS ONE* **2012**, *7*, e41848. [[CrossRef](#)]
26. Zhang, Z.; Korkeala, H.; Dahlsten, E.; Sahala, E.; Heap, J.T.; Minton, N.P.; Lindstrom, M. Two-component signal transduction system CBO0787/CBO0786 represses transcription from botulinum neurotoxin promoters in *Clostridium botulinum* ATCC 3502. *PLoS Pathog.* **2013**, *9*, e1003252. [[CrossRef](#)]

27. Artin, I.; Carter, A.T.; Holst, E.; Lovenklev, M.; Mason, D.R.; Peck, M.W.; Radstrom, P. Effects of carbon dioxide on neurotoxin gene expression in nonproteolytic *Clostridium botulinum* Type E. *Appl. Environ. Microbiol.* **2008**, *74*, 2391–2397. [[CrossRef](#)] [[PubMed](#)]
28. Artin, I.; Mason, D.R.; Pin, C.; Schelin, J.; Peck, M.W.; Holst, E.; Radstrom, P.; Carter, A.T. Effects of carbon dioxide on growth of proteolytic *Clostridium botulinum*, its ability to produce neurotoxin, and its transcriptome. *Appl. Environ. Microbiol.* **2010**, *76*, 1168–1172. [[CrossRef](#)]
29. Lawson, P.A.; Rainey, F.A. Proposal to restrict the genus *Clostridium* Prazmowski to *Clostridium butyricum* and related species. *Int. J. Syst. Evol. Microbiol.* **2016**, *66*, 1009–1016. [[CrossRef](#)]
30. Fratelli, F.; Siquini, T.J.; de Abreu, M.E.; Higashi, H.G.; Converti, A.; de Carvalho, J.C. Fed-batch production of tetanus toxin by *Clostridium tetani*. *Biotechnol. Prog.* **2010**, *26*, 88–92. [[CrossRef](#)]
31. Fratelli, F.; Siquini, T.J.; Prado, S.M.; Higashi, H.G.; Converti, A.; de Carvalho, J.C. Effect of medium composition on the production of tetanus toxin by *Clostridium tetani*. *Biotechnol. Prog.* **2005**, *21*, 756–761. [[CrossRef](#)]
32. Porfirio, Z.; Prado, S.M.; Vancetto, M.D.C.; Fratelli, F.; Alves, E.W.; Raw, I.; Fernandes, B.L.; Camargo, A.C.M.; Lebrun, I. Specific peptides of casein pancreatic digestion enhance the production of tetanus toxin. *J. Appl. Microbiol.* **1997**, *83*, 678–684. [[CrossRef](#)]
33. Carter, G.P.; Cheung, J.K.; Larcombe, S.; Lyras, D. Regulation of toxin production in the pathogenic clostridia. *Mol. Microbiol.* **2014**, *91*, 221–231. [[CrossRef](#)] [[PubMed](#)]
34. Connan, C.; Deneve, C.; Mazuet, C.; Popoff, M.R. Regulation of toxin synthesis in *Clostridium botulinum* and *Clostridium tetani*. *Toxicon* **2013**, *75*, 90–100. [[CrossRef](#)] [[PubMed](#)]
35. Brüggemann, H.; Gottschalk, G. Insights in metabolism and toxin production from the complete genome sequence of *Clostridium tetani*. *Anaerobe* **2004**, *10*, 53–68. [[CrossRef](#)] [[PubMed](#)]
36. Mellanby, J. The effect of glutamate on toxin production by *Clostridium tetani*. *J. Gen. Microbiol.* **1968**, *54*, 77–82. [[CrossRef](#)]
37. Ohtani, K.; Yuan, Y.; Hassan, S.; Wang, R.; Wang, Y.; Shimizu, T. Virulence gene regulation by the agr system in *Clostridium perfringens*. *J. Bacteriol.* **2009**, *191*, 3919–3927. [[CrossRef](#)]
38. Ohtani, K.; Shimizu, T. Regulation of toxin gene expression in *Clostridium perfringens*. *Res. Microbiol.* **2015**, *166*, 280–289. [[CrossRef](#)]
39. Ohtani, K.; Shimizu, T. Regulation of Toxin Production in *Clostridium perfringens*. *Toxins* **2016**, *8*, 207. [[CrossRef](#)]
40. Feng, L.; Zhu, J.; Chang, H.; Gao, X.; Gao, C.; Wei, X.; Yuan, F.; Bei, W. The CodY regulator is essential for virulence in *Streptococcus suis* serotype 2. *Sci. Rep.* **2016**, *6*, 21241. [[CrossRef](#)]
41. Lemos, J.A.; Nascimento, M.M.; Lin, V.K.; Abranches, J.; Burne, R.A. Global regulation by (p)ppGpp and CodY in *Streptococcus mutans*. *J. Bacteriol.* **2008**, *190*, 5291–5299. [[CrossRef](#)]
42. Lindback, T.; Mols, M.; Basset, C.; Granum, P.E.; Kuipers, O.P.; Kovacs, A.T. CodY, a pleiotropic regulator, influences multicellular behaviour and efficient production of virulence factors in *Bacillus cereus*. *Environ. Microbiol.* **2012**, *14*, 2233–2246. [[CrossRef](#)]
43. Lobel, L.; Sigal, N.; Borovok, I.; Ruppin, E.; Herskovits, A.A. Integrative genomic analysis identifies isoleucine and CodY as regulators of *Listeria monocytogenes* virulence. *PLoS Genet.* **2012**, *8*, e1002887. [[CrossRef](#)] [[PubMed](#)]
44. Van Schaik, W.; Chateau, A.; Dillies, M.A.; Coppee, J.Y.; Sonenshein, A.L.; Fouet, A. The global regulator CodY regulates toxin gene expression in *Bacillus anthracis* and is required for full virulence. *Infect. Immun.* **2009**, *77*, 4437–4445. [[CrossRef](#)] [[PubMed](#)]
45. Dineen, S.S.; Villapakkam, A.C.; Nordman, J.T.; Sonenshein, A.L. Repression of *Clostridium difficile* toxin gene expression by CodY. *Mol. Microbiol.* **2007**, *66*, 206–219. [[CrossRef](#)] [[PubMed](#)]
46. Martin-Verstraete, I.; Peltier, J.; Dupuy, B. The regulatory networks that control *Clostridium difficile* toxin synthesis. *Toxins* **2016**, *8*, 153. [[CrossRef](#)] [[PubMed](#)]
47. Daou, N.; Wang, Y.; Levnikov, V.M.; Nandakumar, M.; Livny, J.; Bouillaut, L.; Blagova, E.; Zhang, K.; Belitsky, B.R.; Rhee, K.; et al. Impact of CodY protein on metabolism, sporulation and virulence in *Clostridioides difficile* ribotype 027. *PLoS ONE* **2019**, *14*, e0206896. [[CrossRef](#)]
48. Kovacs, A.T. Bacterial differentiation via gradual activation of global regulators. *Curr. Genet.* **2016**, *62*, 125–128. [[CrossRef](#)]

49. Ravagnani, A.; Jennert, K.C.; Steiner, E.; Grunberg, R.; Jefferies, J.R.; Wilkinson, S.R.; Young, D.I.; Tidswell, E.C.; Brown, D.P.; Youngman, P.; et al. Spo0A directly controls the switch from acid to solvent production in solvent-forming clostridia. *Mol. Microbiol.* **2000**, *37*, 1172–1185. [[CrossRef](#)]
50. Mascher, G.; Derman, Y.; Kirk, D.G.; Palonen, E.; Lindström, M.; Korkeala, H. The CLO3403/CLO3404 two-component system of *Clostridium botulinum* E1 Beluga is important for cold shock response and growth at low temperatures. *Appl. Environ. Microbiol.* **2014**, *80*, 399–407. [[CrossRef](#)]
51. Couesnon, A.; Raffestin, S.; Popoff, M.R. Expression of botulinum neurotoxins A and E, and associated non-toxin genes, during the transition phase and stability at high temperature: Analysis by quantitative reverse transcription-PCR. *Microbiology* **2006**, *152*, 759–770. [[CrossRef](#)]
52. Anellis, A.; Berkowitz, D.; Kemper, D.; Rowley, D.B. Production of types A and B spores of *Clostridium botulinum* by the biphasic method: Effect on spore population, radiation resistance, and toxigenicity. *Appl. Microbiol.* **1972**, *23*, 734–739. [[CrossRef](#)]
53. Mackin, K.E.; Carter, G.P.; Howarth, P.; Rood, J.I.; Lyras, D. Spo0A differentially regulates toxin production in evolutionarily diverse strains of *Clostridium difficile*. *PLoS ONE* **2013**, *8*, e79666. [[CrossRef](#)] [[PubMed](#)]
54. Bouillaud, L.; Dubois, T.; Sonenshein, A.L.; Dupuy, B. Integration of metabolism and virulence in *Clostridium difficile*. *Res. Microbiol.* **2015**, *166*, 375–383. [[CrossRef](#)] [[PubMed](#)]
55. Philippe, V.A.; Mendez, M.B.; Huang, I.H.; Orsaria, L.M.; Sarker, M.R.; Grau, R.R. Inorganic phosphate induces spore morphogenesis and enterotoxin production in the intestinal pathogen *Clostridium perfringens*. *Infect. Immun.* **2006**, *74*, 3651–3656. [[CrossRef](#)]
56. Paredes-Sabja, D.; Sarker, M.R. *Clostridium perfringens* sporulation and its relevance to pathogenesis. *Future Microbiol.* **2009**, *4*, 519–525. [[CrossRef](#)] [[PubMed](#)]
57. Kurt, A.; Ozkan, M.; Ozcengiz, G. Inorganic phosphate has a crucial effect on Cry3Aa delta-endotoxin production. *Letts. Appl. Microbiol.* **2005**, *41*, 303–308. [[CrossRef](#)]
58. Chekabab, S.M.; Harel, J.; Dozois, C.M. Interplay between genetic regulation of phosphate homeostasis and bacterial virulence. *Virulence* **2014**, *5*, 786–793. [[CrossRef](#)]
59. Aggarwal, S.; Somani, V.K.; Bhatnagar, R. Phosphate starvation enhances the pathogenesis of *Bacillus anthracis*. *Int. J. Med. Microbiol.* **2015**, *305*, 523–531. [[CrossRef](#)]
60. Allenby, N.E.; Laing, E.; Bucca, G.; Kierzek, A.M.; Smith, C.P. Diverse control of metabolism and other cellular processes in *Streptomyces coelicolor* by the PhoP transcription factor: Genome-wide identification of in vivo targets. *Nucleic Acids Res.* **2012**, *40*, 9543–9556. [[CrossRef](#)]
61. Baker, J.J.; Johnson, B.K.; Abramovitch, R.B. Slow growth of *Mycobacterium tuberculosis* at acidic pH is regulated by *phoPR* and host-associated carbon sources. *Mol. Microbiol.* **2014**, *94*, 56–69. [[CrossRef](#)]
62. Lovenklev, M.; Artin, I.; Hagberg, O.; Borch, E.; Holst, E.; Radstrom, P. Quantitative interaction effects of carbon dioxide, sodium chloride, and sodium nitrite on neurotoxin gene expression in nonproteolytic *Clostridium botulinum* type B. *Appl. Environ. Microbiol.* **2004**, *70*, 2928–2934. [[CrossRef](#)]
63. Karlsson, S.; Burman, L.G.; Akerlund, T. Suppression of toxin production in *Clostridium difficile* VPI10463 by amino acids. *Microbiology* **1999**, *145*, 1683–1693. [[CrossRef](#)] [[PubMed](#)]
64. Burgess, C.; Gasse, F.; Steinglass, R.; Yakubu, A.; Raza, A.A.; Johansen, K. Eliminating maternal and neonatal tetanus and closing the immunity gap. *Lancet* **2017**, *389*, 1380–1381. [[CrossRef](#)]

