

Supplementary Materials: Influence of Endosperm Starch Composition on Maize Response to *Fusarium temperatum* Scaufl. & Munaut

Marcin Wit, Piotr Ochodzki, Roman Warzecha, Emilia Jabłońska, Ewa Mirzwa-Mróz, Elżbieta Mielniczuk, and Wojciech Wakuliński

Table S1. ANOVA analysis of the influence of the maize type, cropping season, and field localization on disease severity index I_{FER} after plant inoculation with *F. temperatum*.

Source of variation	ANOVA traits				
	SS	df	MS	F	p
Intercept	140875.1	1	140875.1	785.1575	0.000000
Maize type (A)	2865.9	2	1433.0	7.9865	0.000369
Year (B)	3847.8	3	1282.6	7.1485	0.000097
Localities (C)	8551.7	2	4275.8	23.8310	0.000000
A X B	1350.7	6	225.1	1.2547	0.276174
A X C	754.3	4	188.6	1.0510	0.379854
B X C	4428.8	6	738.1	4.1139	0.000448
A X B X C	1249.6	12	104.1	0.5804	0.859055
Error	138514.4	772	179.4		

Abbreviations: SS, sum of squares; df, degrees of freedom; MS, Mean square; F, F ratio; p, probability.

Table S2. ANOVA analysis of the influence of the maize type, cropping season, and field localization on BEA kernel contamination after plant inoculation with *F. temperatum*.

Source of variation	ANOVA traits				
	SS	df	MS	F	p
Intercept	20874.16	1	20874.16	454.6599	0.000000
Maize type (A)	780.60	2	390.30	8.5011	0.000223
Year (B)	1795.12	3	598.37	13.0332	0.000000
Localities (C)	241.12	2	120.56	2.6259	0.073039
A X B	652.21	6	108.70	2.3676	0.028426
A X C	365.52	4	91.38	1.9903	0.094177
B X C	936.68	6	156.11	3.4003	0.002569
A X B X C	211.52	12	17.63	0.3839	0.969385
Error	34479.60	751	45.91		

Abbreviations: SS, sum of squares; df, degrees of freedom; MS, Mean square; F, F ratio; p, probability.

Table S3. ANOVA analysis of the influence of the maize type, cropping season, and field localization on the ergosterol (ERG) content after plant inoculation with *F. temperatum*.

Source of variation	ANOVA traits				
	SS	df	MS	F	p
Intercept	140875.1	1	140875.1	785.1575	0.000000
Maize type (A)	2865.9	2	1433.0	7.9865	0.000369
Year (B)	3847.8	3	1282.6	7.1485	0.000097
Localities (C)	8551.7	2	4275.8	23.8310	0.000000
A X B	1350.7	6	225.1	1.2547	0.276174
A X C	754.3	4	188.6	1.0510	0.379854
B X C	4428.8	6	738.1	4.1139	0.000448
A X B X C	1249.6	12	104.1	0.5804	0.859055
Error	138514.4	772	179.4		

Abbreviations: SS, sum of squares; df, degrees of freedom; MS, Mean square; F, F ratio; p, probability.

Table S4. ANOVA analysis of the influence of the maize type, cropping season, and field localization influence on the amylose content in maize kernels.

Source of variation	ANOVA traits				
	SS	df	MS	F	p
Intercept	519879.4	1	519879.4	94644.65	0.000000
Maize type (A)	2052.4	2	1026.2	186.82	0.000000
Year (B)	1294.4	3	431.5	78.55	0.000000
Localities (C)	220.6	2	110.3	20.08	0.000000
A X B	32.6	6	5.4	0.99	0.431839
A X C	51.6	4	12.9	2.35	0.052858
B X C	145.2	6	24.2	4.40	0.000217
A X B X C	246.9	12	20.6	3.75	0.000015
Error	4251.6	774	5.5		

Abbreviations: SS, sum of squares; df, degrees of freedom; MS, Mean square; F, F ratio; p, probability.