

Combined Assessment of Preoperative Frailty and Sarcopenia Allows the Prediction of Overall Survival in Patients with Lung Cancer (NSCLC) and Surgically Treated Brain Metastasis

Inja Ilic ^{1,†,*}, Anton Faron, ^{2,†}, Muriel Heimann ¹, Anna-Laura Potthoff ¹, Niklas Schäfer ³, Christian Bode ⁴, Valeri Borger ¹, Lars Eichhorn ⁴, Frank A. Giordano ⁵, Erdem Güresir ¹, Andreas H. Jacobs ⁶, Yon-Dschun Ko ⁷, Jennifer Landsberg ⁸, Felix Lehmann ⁴, Alexander Radbruch ⁹, Ulrich Herrlinger ³, Hartmut Vatter ¹, Patrick Schuss, ^{1,‡} and Matthias Schneider ^{1,‡}

Table S1. Overview of the distribution of known molecular biomarkers in the present cohort of NSCLC patients*.

	Expression		
	n.a.	+	-
PD-L1	110 (78)	16 (11)	15 (11)
EGFR-mutant	93 (66)	4 (3)	44 (31)
ALK-rearranged	92 (65)	1 (1)	48 (34)
ROS1-rearranged	121 (86)	1 (1)	19 (13)

* Values represent number of patients unless otherwise indicated (%). ALK, anaplastic lymphoma kinase; EGFR, epidermal growth factor receptor; n.a., not applicable; NSCLC, non-small cell lung carcinoma; PD-L1, anti-programmed cell death 1 ligand 1.