

Supplementary information

Intratumoural Delivery of mRNA Loaded on a Cationic Hyper-Branched Cyclodextrin-Based Polymer Induced an Anti-Tumour Immunological Response in Melanoma

Yousef Khazaei Monfared^{1,†}, Mohammad Mahmoudian², Parvin Zakeri-Milani^{2,*}, Claudio Cecone¹, Tomoya Hayashi³, Ken J. Ishii³, João Conde⁴, Adrián Matencio¹ and Francesco Trotta^{1,*†}

¹ Department of Chemistry, University of Turin, 10125 Turin, Italy; yousef.khazaei@unito.it (Y.K.M.); claudio.cecone@unito.it (C.C.); adrian.matencioduran@unito.it (A.M.)



² Faculty of Pharmacy, Tabriz University of Medical Sciences, Tabriz 5165665931, Iran; mahmoodian.nano@gmail.com

³ Division of Vaccine Science, Department of Microbiology and Immunology, The Institute of Medical Science, The University of Tokyo (IMSUT), Tokyo 113-8654, Japan; tomoya-h@ims.u-tokyo.ac.jp (T.H.); kenishii@ims.u-tokyo.ac.jp (K.J.I.)

⁴ ToxOomics, NOVA Medical School (NMS), Faculdade de Ciências Médicas (FCM), Universidade Nova de Lisboa, 1099-085 Lisboa, Portugal; joao.conde@nms.unl.pt

* Correspondence: pzakeri@tbzmed.ac.ir (P.Z.-M.); francesco.trotta@unito.it (F.T.)

† These authors contributed equally to this work

A)**B)**

CD 8+



**C)**

Figure S1. A, Gel retardation assay for OVA, left, and EGFP-mRNA, right. **B,** Quantification of H2-Kb SIINFEKL expression and SIINFEKL+ CD8+ T cells antigen specific T cell responses in tumor tissues, and **C,** Spleen tissues following treatment with formulations. (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, # $p < 0.0001$).




Figure S2. **A**, Cytotoxicity results for different concentrations of Ppoly. **B**, Different ratios (1:1, 1:5 and 1:10) of Ppoly free and complexes with mRNA. **C**, Different volumes of Lipofectamine (μL) free and complexes with mRNA against melanoma cancer cell lines. (ns, Not Significant; * $p < 0.05$, ** $p < 0.01$)