

## **Supplementary Appendix**

This appendix has been provided by the authors to give readers additional information regarding their work.

Supplement to: Prediction of prognosis in pancreatic cancer according to methionyl-tRNA synthetase 1 levels determined by immunohistochemical staining


Sung Ill Jang, Ji Hae Nahm, See Young Lee, Jae Hee Cho, Min-Young Do, Joon Seong Park, Hye Sun Lee, Juyeon Yang, Jiwon Kong, Seunghwan Jung, Sunghoon Kim and Dong Ki Lee

### **Contents**

#### **1. Supplementary Results: page**

**Supplementary Figure S1: page 2**

**Supplementary Table S1: page 2**



Supplementary Figure S1. CA 19-9 and CEA level analyzed using Z-score.

Supplementary Table S1 Comparison of CA 19-9 and CEA level between two group.

Variables	Non-Parametric Method		
	Low MARS1 Expression Group (n=55)	High MARS1 Expression Group (n=82)	<i>p</i> -Value
CA 19-9 at adm, IU/L (median(IQR))	97.80 (25.30, 408.00)	58.80 (14.50, 373.80)	0.437
CEA at adm, IU/L (median(IQR))	3.98 (1.40, 6.80)	2.79 (1.75, 4.60)	0.2771

MARS1, methionyl-tRNA synthetase 1; se, standard error; CI, confidence intervals.